FOLFOXIRI

FOLFOXIRI combines three chemotherapy medicines with folinic acid to treat colorectal cancer that has spread. The aim is cancer control, symptom relief and, for selected people, a response that allows further local treatment after specialist review.

Metastatic colorectal cancer · NCCP 00555 v7 · Document revised 4 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

My overall treatment

Day 1: irinotecan, oxaliplatin and folinic acid, followed by a 48-hour fluorouracil pump. Repeat every 14 days while treatment is helping and side effects remain acceptable.

14-day treatment cycle

Day 1: irinotecan, oxaliplatin and folinic acid, followed by a 48-hour fluorouracil pump. Repeat every 14 days while treatment is helping and side effects remain acceptable.

Your treatment schedule

Day 1	Days 2–3	Days 4–14
Day-ward medicines; pump connected.	Complete 48-hour pump; disconnect as arranged.	Recovery, symptoms and tests; next cycle only after team review.

Your team confirms the active phase and dates. Symptoms may occur during or after treatment; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before	**DPD / baseline bloods** Required by regimen	DPD before fluoropyrimidine; blood count, kidney and liver tests.
During	**Bloods / nerves** Required by regimen	Blood count, kidney and liver tests before each cycle. Nerve review every cycle.
If indicated	**Heart** Conditional tests	ECG if cardiac function is compromised.
After / review	**Recovery / scans / markers** Team-agreed plan	Follow the individual cancer plan and any unresolved effects; scan and marker intervals are team-agreed.

Between treatments

- No guaranteed safe days: report fever, sudden illness, diarrhoea or poor intake promptly.
- Symptoms can continue after pump removal and after treatment ends.

Confirm supportive medicines, tests, working urgent contacts and pump disconnection.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Mouth / swallowing

Fluorouracil + folinic acid

Report ulcers or pain that affects eating or drinking.

3 Bowel / stoma

Irinotecan / fluorouracil

Call early for diarrhoea, increased stoma output or poor intake; severe pain needs urgent assessment.

5 Blood / infection


Chemotherapy

Fever, shivers or sudden illness: call now. Unusual bruising or marked weakness also needs advice.

7 Hands / feet

Fluorouracil

Report painful redness, peeling or changes that affect usual activity.



2 Chest

Fluorouracil

New chest pain, collapse or major breathing difficulty needs emergency help.

4 Nerves / cold

Oxaliplatin

Avoid cold triggers; report persistent numbness or difficulty with buttons or walking.

6 Hair / scalp

Irinotecan

Hair thinning or loss may occur. Ask about practical support.

8 Pump / line

Treatment delivery

Call for leakage, swelling, line pain or a loose connection. Do not reconnect it yourself.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Fever, shivers or sudden illness needs immediate oncology advice. Chest pain, collapse or severe breathing difficulty needs emergency help. Call urgently for severe or persistent diarrhoea, severe abdominal pain, inability to drink, reduced urine or a line/pump problem.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

Keep your treatment details to hand

Metastatic colorectal cancer · NCCP 00555 v7 · Document revised 4 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

FOLFOXIRI

Every 14 days · 48-hour fluorouracil pump · cycle plan confirmed by team.

Phase: _____ Cycle plan: team to confirm

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

CUT outer border

BACK / TREATMENT RECORD

My cycle record

Rolling record — follow your agreed treatment plan.

Cycle	Date / initials

My treatment information

Latest treatment date: _____

Plan changed? Ask the team to update this card.

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____

Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.