Phesgo maintenance

Phesgo on Day 1 every 21 days after the chemotherapy component is complete; continue while beneficial and tolerated. This is active metastatic maintenance, not a fixed one-year adjuvant course.

Source reconciled; independent clinical, pharmacy and patient/PPI review pending. Selected effects; not a complete medicine leaflet.

A picture of side effects

Numbered symptom boxes connect to the anatomical illustration. The medicine responsible may be uncertain; report the symptom.

1 Heart / breathing


Phesgo

Keep echoes; report breathlessness, swelling or reduced exercise tolerance.

2 Diarrhoea

Especially pertuzumab

Report watery stools early, particularly with poor intake or dizziness.



3 Injection reaction

Phesgo

Tell staff immediately about rash, chest tightness or breathing difficulty.

4 Thigh injection site

Phesgo

Report persistent or spreading redness, swelling or pain.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

The body map follows the medicines active in this phase.

> Fever, shivers or suddenly unwell: contact oncology immediately. Severe chest pain, breathing difficulty, collapse or seizure: 112 / 999.

Your treatment, at home.

Important medicine-specific effects

Pertuzumab

- Diarrhoea and dehydration
- Injection / allergic reaction
- Heart pumping-function reduction
- Rash and fatigue

Trastuzumab

- Heart pumping-function reduction / heart failure
- Injection or allergic reaction
- Diarrhoea and fatigue
- Rare serious lung effects

Surgery, radiation and monitoring: open the whole-treatment map or scan the QR.

Make the plan practical

Antiemetic plan

Phesgo alone has minimal emetic risk; routine prophylactic antiemetics are not populated.

Premedicines

No routine Phesgo premedication unless indicated by prior reaction or the individual plan.

Appointments and results

Echo before treatment and every 12 weeks (around cycle 5 if on time), including maintenance. Bloods before each cycle. Ask who books and reviews each test; report a missing appointment.

Phase transition

After chemotherapy stops, use the agreed Phesgo maintenance plan (NCCP 00785 when eligible). Keep cardiac and response checks; chemotherapy steroids, antiemetics and growth factors do not automatically continue.

Emergency: 112 / 999. Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

Contact oncology immediately for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital's instructions. If you cannot reach the team, seek urgent medical assessment.

YOUR ONCOLOGY TEAM · DAYTIME

Insert numbers here

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

Insert numbers here

Ask for medical oncology

EVENINGS / OVERNIGHT

Your out-of-hours pathway / nearest ED

Other hospitals: use the numbers your team provides.

Complete reference, visual timeline and passport

https://sactcheck.com/patient/00785/

Sources and review status are available online. Follow your own prescription and hospital instructions.