NCCP 00797 · clinician draft

Phesgo + weekly paclitaxel

Supportive prescription draft for review. This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Taxane + Phesgo phase

Active medicines: Paclitaxel + Pertuzumab + Trastuzumab

Emetogenic risk: Low

Medicine	Dose / route	When
Ondansetron	8 mg orally	On treatment day Review QT, constipation and hepatic dosing; reconcile any steroid already prescribed
Metoclopramide	10 mg orally up to three times daily as needed	Subsequent days: duration to be confirmed Assess contraindications, renal function and interactions

Additional phase-specific support

- First paclitaxel dose: dexamethasone 8 mg IV, chlorphenamine 10 mg IV and famotidine 20 mg IV, each 30 minutes before paclitaxel.
- Day 8 and thereafter: no routine dexamethasone; chlorphenamine 10 mg IV and famotidine 20 mg IV 30 minutes before. Consultant may add steroid for risk/prior reaction or omit famotidine if appropriate; monitor closely.
- Rechallenge steroid changes require consultant guidance; consider analgesia for taxane aches and G-CSF according to individual risk.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00797 v4 · p6 · NCCP antiemetics v8, Table 1 pp4–5

Sources checked 2026-10-20 · SACTCheck v0.84.12 · Regimen-level drafts; prescriber and oncology-pharmacy verification required