

Pembrolizumab + pemetrexed + carboplatin

Lung · NCCP 00568 · version 5

Your medicines

- **Pembrolizumab** - immunotherapy
- **Pemetrexed** - chemotherapy
- **Carboplatin** - chemotherapy during induction cycles 1-4

Your treatment schedule


- **Cycle length:** every 21 days
- **Cycles 1-4:** all 3 medicines on day 1
- **Maintenance:** pembrolizumab + pemetrexed every 21 days if appropriate

Pembrolizumab · immune-related effects

Pemetrexed / carboplatin · chemotherapy effects

A picture of side effects

Numbers connect each symptom box to the relevant anatomical area. Tell your team about any new or worsening symptom.



Selected effects only; the map does not show likelihood or every possible side effect.

Pembrolizumab + pemetrexed + carboplatin

Lung · NCCP 00568 · version 5

Treatment journey

Before cycle 1 Blood tests, kidney function, glucose/thyroid testing; folic acid and vitamin B12 support.	Cycles 1-4 Day 1: pembrolizumab + pemetrexed + carboplatin every 21 days.	Between cycles Watch for immune symptoms and chemotherapy effects; they can occur at any time.	After cycle 4 Carboplatin usually stops if treatment continues as planned.	Maintenance Pembrolizumab + pemetrexed every 21 days if appropriate.
---	---	--	--	--

Important medicine-specific effects

Pembrolizumab - immune effects - Immune inflammation can affect lungs, bowel, liver, hormone glands, kidneys, skin, heart, nerves or muscles. - New symptoms can begin during treatment or after treatment has stopped.	Pemetrexed - Low blood counts can increase infection, anaemia or bruising risk. - Mouth/gut inflammation and skin rash can occur. - Kidney function can deteriorate and is monitored.	Carboplatin - Low blood counts, especially neutropenia and low platelets, can occur. - Allergic infusion reactions can occur, particularly after repeated exposure. - Nausea and vomiting may occur despite preventative medicines.
--	--	--

Shared overview

The body map and these medicine boxes use the same symptom groupings so the guide, portal and passport use one consistent language.

Monitoring / medicines - Folic acid exactly as prescribed while on pemetrexed. - Vitamin B12 before pemetrexed and at protocol-defined intervals. - Blood count, kidney/liver profile and glucose before each cycle; thyroid monitoring during treatment.	Bring / track - Current medicines, including non-prescription products. - New symptoms: when they started and what they stopped you doing. - Temperature, bowel counts or other readings your team asked you to record.
--	--

WHEN TO CALL - DO NOT WAIT FOR THE NEXT APPOINTMENT

Call oncology urgently for fever, new breathlessness, persistent diarrhoea, jaundice, severe rash, marked weakness, confusion or any rapidly worsening symptom. Severe breathing difficulty, collapse, seizure or severe chest pain: call 112/999.

Consent discussion support

Use the approved consent process to document treatment aim, expected benefit, material risks, alternatives, questions and the patient's decision.

Open the regimen-specific patient portal

The QR and the direct web link both open the same live patient portal. The portal contains regimen information only; no patient-identifying data are encoded.

Open patient portal online

<https://sactcheck.com/patient/00568/>
Official NCCP 00568 v5

