Treatment scope · NCCP 00215 Lung context only. Confirm whether the platinum pathway or EGFR/erlotinib pathway applies. This guide does not supply the partner dose or replace its safety assessment.

NCCP 00215 · VERSION 7

Bevacizumab 15 mg/kg · lung treatment component

Bevacizumab 15 mg/kg is infused on Day 1 every 21 days. With platinum chemotherapy, the combined phase lasts up to six cycles, then bevacizumab may continue alone. With erlotinib, the combination continues while helpful and tolerated. Progression or unacceptable toxicity can stop either plan earlier; no fixed overall lung-treatment duration is promised.

Discussion aid: review with the patient and complete the approved local consent process.

Purpose and plan

☐ Exact indication, treatment aim and uncertainty explained

☐ Expected duration and review / switch points discussed

☐ Alternatives and the option not to proceed discussed

☐ Patient priorities, capacity and support needs explored

Practical safety

☐ Medicine, route, schedule and missed-dose instructions

☐ Food, interactions and safe handling reviewed

☐ Pregnancy, contraception / fertility and breastfeeding

☐ Monitoring, symptom reporting and contact plan

Selected material risks

☐ Raised blood pressure and protein in the urine

☐ Bleeding, blood clots and problems with wound healing

☐ Rare bowel perforation, abnormal internal connections called fistulae, or serious wound infection

☐ Rare severe headache, confusion, seizures or vision changes associated with a brain syndrome

☐ Infusion reactions and fatigue; the partner treatment adds its own risks

Confirm the chosen lung pathway, disease-control aim, alternatives and the evidence for the specific partner. Discuss blood-pressure and urine monitoring, pulmonary and other bleeding, clots, bowel perforation/fistula, wound healing, infusion reactions, fertility/pregnancy precautions and all partner risks. The chemotherapy and erlotinib trials showed different outcomes; neither predicts an individual result.

Individual benefit, alternatives and concerns

Agreed checks and next review

Before starting: blood count, kidney and liver tests, urine protein, blood pressure and a heart history/examination. Repeat blood count, kidney/liver tests and urine testing before each cycle; check blood pressure before treatment and after it. Extra checks depend on symptoms and results.

Clinician / date: _____ Patient discussion documented: _____

Consent is an ongoing conversation and may be reconsidered. This aid is not treatment authorisation.

SACTCheck · Developed by Dr Paul O'Brien · v0.84.25 · Sources checked 2026-10-29
Source reconciled; independent clinical, pharmacy and patient/PPI review pending