Weekly paclitaxel · relapsed or refractory SCLC

NCCP 00226 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

Your practical treatment plan

Paclitaxel 80 mg/m² is infused on Days 1, 8, 15 and 22 of a 28-day cycle. The consultant may instead prescribe Days 1, 8 and 15. Each planned weekly dose needs review. Continue while helping and tolerated; NCCP does not set a fixed cycle cap.

STEP 1

Before the visit

Check the next weekly appointment and whether Day 22 is included. Blood results and symptom review are needed before every dose.

STEP 2

At the unit

Paclitaxel is given through a cannula or suitable existing IV line over about one hour. The appointment also includes checks, preparation, premedication and observation and is longer than the infusion itself. This regimen has no routine take-home pump.

STEP 3

At home

Keep the contact number, medicine list and symptom record available. Ask who will arrange your next blood test, scan and review. Sedating medicines may affect driving.

Planned course

Paclitaxel 80 mg/m² is infused on Days 1, 8, 15 and 22 of a 28-day cycle. The consultant may instead prescribe Days 1, 8 and 15. Each planned weekly dose needs review. Continue while helping and tolerated; NCCP does not set a fixed cycle cap.

Sources: NCCP 00226 v9 · official regimen · HPRA conventional paclitaxel SmPC; 9 January 2026. Source reconciled; independent clinical, pharmacy and patient/PPI review pending.