NCCP 00259 · clinician draft

Vinorelbine capsules · non-small-cell lung cancer

Treatment scope · NCCP 00259 Lung monotherapy context only. The breast guide and its evidence remain separate. Discuss the exact disease setting and alternatives with the treating team.

Supportive prescription draft for review. This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Vinorelbine capsules · non-small-cell lung cancer · support plan

Active medicines: Vinorelbine

Emetogenic risk: Moderate-to-high oral risk

Medicine	Dose / route	When
One 5-HT3 antagonist · choose a single option	Ondansetron 8–16 mg total daily OR granisetron 1–2 mg total daily	Around the weekly dose; the prescriber must state the number of subsequent days NCCP Table 5 oral pathway. Do not combine both agents or assume indefinite daily treatment. Review constipation, QT risk and interacting medicines.

Additional phase-specific support

- Individual antiemetic plan differs by route; list routine and rescue medicines separately.
- Discuss constipation prevention, hydration/diet where appropriate and prescribed laxatives; regular opioid use increases the need for a bowel plan.
- No routine steroid premedication, G-CSF, antibacterial, antiviral or PJP prophylaxis is mandated by these NCCP monotherapy documents. Consider additional prevention based on individual risk/local policy; do not create a generic automatic prescription.
- Gastric protection is a consideration for high gastrointestinal-bleed risk, not a universal drug.
- Confirm strong CYP3A4 inhibitors/inducers, P-gp modulators, phenytoin, itraconazole, warfarin and live vaccines. Yellow fever vaccine is contraindicated. Avoid grapefruit per NCCP; do not self-start supplements.
- Table 5 says continue daily; for once-weekly vinorelbine the prescriber must state the number of days. Do not automatically turn this into indefinite daily treatment or import an IV moderate steroid/NK1 schedule.
- A separate locally approved as-needed plan; check constipation and QT/interactions. Avoid duplicate 5-HT3 agents. Ondansetron can worsen constipation.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00259 v6 · p5 · NCCP antiemetic guidance v6 · oral Table 5

Sources checked 2026-10-29 · SACTCheck v0.84.25 · Regimen-level drafts; prescriber and oncology-pharmacy verification required