Carry your treatment details

Adult treatment scope · NCCP 00702a / 00702b This adult suite covers entrectinib alone for ROS1-positive advanced NSCLC before any ROS1 inhibitor (00702a), or a separate NTRK gene-fusion solid-tumour indication with no previous NTRK inhibitor and no satisfactory treatment options (00702b). Molecular tests and Irish access criteria must be confirmed. The adult 600 mg schedule does not apply to children.

Print, cut out and fold. Follow your treating hospital's contact plan.

SACTCHECK · TREATMENT CARD

Entrectinib monotherapy · adult

NCCP 00702 v2

The usual adult starting dose is 600 mg by mouth once daily, every day without a planned rest week. Use your prescribed dose. Treatment continues while helping and side effects remain acceptable; there is no fixed number of cycles.

Prescriber / next review: _____

Contact plan

Oncology team: Insert numbers here · Use your hospital's opening hours.
Out of hours: Insert numbers here.
Evenings / overnight: agreed out-of-hours service or emergency department.
Emergency: 112 / 999.

Own hospital / contact: _____

Call the oncology team urgently for new or worsening breathlessness, swelling, palpitations or fainting; marked confusion or memory change; fever, jaundice, severe diarrhoea or inability to drink. Report new visual disturbance and focal bone pain promptly. Follow your unit's temperature/contact instructions. Severe breathing difficulty, chest pain or collapse needs emergency help.

https://sactcheck.com/patient/00702/ · no patient information encoded.

SACTCheck · Developed by Dr Paul O'Brien · v0.84.25 · Sources checked 2026-10-29
Source reconciled; independent clinical, pharmacy and patient/PPI review pending