Carry your treatment details

Advanced MET exon 14 skipping lung cancer NCCP 00823a covers tepotinib on its own for adults with advanced non-small cell lung cancer carrying a confirmed MET exon 14 skipping change, after immunotherapy and/or platinum chemotherapy. It is not a first-line or adjuvant treatment guide.

Print, cut out and fold. Follow your treating hospital’s contact plan.

SACTCHECK · TREATMENT CARD

Tepotinib monotherapy

NCCP 00823 v2

The usual starting dose is 450 mg once daily: two 225 mg tablets taken together with food. Treatment is continuous, without a planned rest week, while it is helping. Follow your prescribed dose; the team may use one 225 mg tablet or pause treatment for side effects.

Prescriber / next review: _____

Contact plan

Oncology team: Insert numbers here · Use your hospital’s opening hours.
Out of hours: Insert numbers here.
Evenings / overnight: agreed out-of-hours service or emergency department.
Emergency: 112 / 999.

Own hospital / contact: _____

Contact the oncology team urgently for new or worsening cough, breathlessness or fever; marked swelling or sudden weight gain; severe diarrhoea, repeated vomiting or inability to drink; yellow skin or eyes, dark urine or severe abdominal pain; or palpitations or fainting. Severe breathing difficulty, collapse or swelling of the mouth or throat needs emergency help. Use your unit’s temperature and 24-hour contact instructions.

https://sactcheck.com/patient/00823/ · no patient information encoded.

SACTCheck · Developed by Dr Paul O’Brien · v0.84.25 · Sources checked 2026-10-29
Source reconciled; independent clinical, pharmacy and patient/PPI review pending