Carry your treatment details

Advanced ALK-positive NSCLC after crizotinib Adults with ALK-positive advanced non-small-cell lung cancer previously treated with crizotinib: NCCP 00340a. This suite does not turn the broader first-line licence into an NCCP indication or funding approval. Confirm your treatment setting with the team.

Print, cut out and fold. Follow your treating hospital’s contact plan.

SACTCHECK · TREATMENT CARD

Ceritinib monotherapy

NCCP 00340 v4

The usual starting dose is 450 mg once daily with food, every day without a planned rest week. Follow your own prescription. Treatment continues while it helps and side effects remain acceptable; there is no fixed course length.

Prescriber / next review: _____

Contact plan

Oncology team: Insert numbers here · Use your hospital’s opening hours.
Out of hours: Insert numbers here.
Evenings / overnight: agreed out-of-hours service or emergency department.
Emergency: 112 / 999.

Own hospital / contact: _____

Contact oncology urgently for new or worsening cough or breathlessness, fainting, palpitations, severe dizziness, yellow skin or eyes, dark urine, severe persistent abdominal pain, persistent vomiting or diarrhoea, poor drinking or little urine, or marked thirst with frequent urination and feeling unwell. Severe breathing difficulty, collapse or severe chest pain needs emergency help.

https://sactcheck.com/patient/00340/ · no patient information encoded.