

Pemetrexed + carboplatin

Lung - NCCP 00318 - version 6

Your medicines

Pemetrexed - Carboplatin

Your treatment schedule

Pemetrexed 500 mg/m² IV followed by carboplatin AUC 5 IV.

A picture of side effects

Tell your team about new or worsening symptoms. Numbers link each symptom box to the body map.

Pemetrexed - kidney / mouth / skin

Carboplatin - blood counts / allergy / nausea

1 Blood / infection

Pemetrexed / carboplatin

Low blood counts - especially neutrophils and platelets - can cause infection or bleeding. Fever or sudden illness needs prompt contact.

3 Nausea / bowel

Carboplatin / pemetrexed

Nausea, vomiting or diarrhoea can occur. Use prescribed supportive medicines and call if you cannot maintain fluids.

5 Skin

Pemetrexed

Rash can occur; prescribed steroid prophylaxis helps reduce this risk.

2 Kidneys

Pemetrexed

Kidney function affects pemetrexed suitability and is checked before treatment. Report reduced urine or dehydration.

4 Mouth


Pemetrexed

Mouth soreness or ulcers can affect eating and drinking; report significant symptoms.

6 Infusion / allergy

Carboplatin

Rash, wheeze, swelling, dizziness or breathing difficulty during infusion needs immediate assessment.



Selected effects only. The map does not show likelihood and cannot diagnose the cause of a symptom.

Pemetrexed + carboplatin

Lung - NCCP 00318 - version 6

Your treatment journey

Day 1: Pemetrexed 500 mg/m² IV followed by carboplatin AUC 5 IV.

Every 21 days: Treatment repeats every 21 days if review and blood tests allow.

Mesothelioma: Usually up to 4-6 cycles unless progression or unacceptable toxicity occurs.

NSCLC: Usually 4-6 cycles, then maintenance pemetrexed (NCCP 00222) where appropriate.

Monitoring and support

- Folic acid and vitamin B12 support are required with pemetrexed.
- A corticosteroid is used around pemetrexed to reduce skin reactions.
- Blood count, kidney and liver profile are checked before each cycle.
- Carboplatin dosing depends on renal function.
- Follow the regimen-specific antiemetic prescription.

WHEN TO CALL

Contact oncology urgently for fever or sudden illness, significant bleeding, new breathing difficulty or infusion reaction, uncontrolled vomiting/diarrhoea, severe mouth soreness, marked reduction in urine or rapidly worsening symptoms. For a medical emergency call 112/999.

Consent discussion support

Use the approved consent process to document treatment intent, expected benefit, material risks, alternatives, questions and the patient's decision.

Scan to open the regimen-specific patient portal

<https://sactcheck.com/patient/00318/>

Official NCCP 00318 v6

SACTCheck v0.80.4 - patient information prototype for clinical review. Follow your oncology team and local emergency instructions.

