

Nivolumab + ipilimumab + pemetrexed + carboplatin

Lung - NCCP 00713 - version 4

Your medicines

Nivolumab - Ipilimumab - Pemetrexed - Carboplatin

Your treatment schedule

Nivolumab + ipilimumab + pemetrexed + carboplatin.

A picture of side effects

Tell your team about new or worsening symptoms. Numbers link each symptom box to the body map.

Nivolumab / ipilimumab - immune effects

1 Hormones / brain

Nivolumab / ipilimumab

Fatigue, dizziness, headache, thirst or passing more urine can indicate immune endocrine toxicity.

3 Heart / muscles

Nivolumab / ipilimumab

Chest pain, palpitations, fainting or marked muscle weakness needs urgent assessment.

5 Kidneys

Pemetrexed / immune treatment

Kidney function is monitored. Report reduced urine, swelling or significant dehydration.

7 Mouth / skin

Pemetrexed / immune treatment

Mouth soreness or significant rash should be reported; severe blistering skin needs urgent review.

Pemetrexed / carboplatin - chemotherapy

2 Lungs

Nivolumab / ipilimumab

New cough or breathlessness needs prompt assessment for pneumonitis and other causes.

4 Liver / bowel

Nivolumab / ipilimumab

Jaundice, dark urine, persistent diarrhoea or abdominal pain can indicate immune inflammation.

6 Blood / infection


Pemetrexed / carboplatin

Fever, shivers, bruising or sudden illness can reflect low blood counts.

8 Infusion / allergy

Carboplatin / immunotherapy

Wheeze, swelling, dizziness or an acute infusion reaction needs immediate assessment.



Selected effects only. The map does not show likelihood and cannot diagnose the cause of a symptom.

Nivolumab + ipilimumab + pemetrexed + carboplatin

Lung - NCCP 00713 - version 4

Your treatment journey

- Cycle 1 - day 1: Nivolumab + ipilimumab + pemetrexed + carboplatin.
- Cycle 1 - day 22: Nivolumab + pemetrexed + carboplatin.
- After cycle 1: Continue nivolumab on days 1 and 22 and ipilimumab on day 1 of each 42-day cycle.
- Duration: Immunotherapy continues until progression/unacceptable toxicity or up to 24 months in patients without progression, according to the NCCP source.

Monitoring and support

- Folic acid and vitamin B12 support are required with pemetrexed.
- Blood count, kidney/liver profile and endocrine monitoring support treatment safety.
- Use the regimen-specific antiemetic prescription for the chemotherapy phase.
- Immune adverse events can begin during treatment or after it stops; report new symptoms early.

WHEN TO CALL
Contact oncology urgently for fever, new breathlessness, persistent diarrhoea, jaundice, severe rash, chest pain, marked weakness, confusion, reduced urine or rapidly worsening symptoms. For severe breathing difficulty, collapse, seizure or severe chest pain call 112/999.

Consent discussion support

Use the approved consent process to document treatment intent, expected benefit, material risks, alternatives, questions and the patient's decision.