

TCHP

Breast - NCCP 00722 - version 2

Your medicines

Docetaxel - Carboplatin - Trastuzumab - Pertuzumab

Your treatment schedule

Day 1 every 21 days during chemotherapy; HER2-directed treatment then continues per plan.

A picture of side effects

Tell your team about new or worsening symptoms. Numbers link each symptom box to the body map.

Trastuzumab / pertuzumab - HER2 treatment

Docetaxel / carboplatin - chemotherapy

1 Heart

Trastuzumab / pertuzumab

Heart function is monitored. Report new breathlessness, swelling or chest symptoms.

3 Mouth / taste

Chemotherapy

Report mouth sores, difficulty drinking or significant taste changes.

5 Nerves

Docetaxel

Report tingling or numbness, especially if persistent.

7 Infusion symptoms

Any infused medicine

Tell the nurse immediately about wheeze, flushing, rash, swelling or sudden symptoms during treatment.

2 Bowel

Pertuzumab-containing treatment

Persistent or severe diarrhoea needs early advice.

4 Blood / infection

Docetaxel / carboplatin

Fever or sudden illness needs immediate oncology contact.

6 Skin / nails


Docetaxel-containing treatment

Report rash, nail changes or painful skin.

8 Swelling

Treatment-related symptoms

New leg swelling or breathlessness needs assessment; symptoms may have several causes.



Selected effects only. The map does not show likelihood and cannot diagnose the cause of a symptom.

TCHP

Breast - NCCP 00722 - version 2

Your treatment journey

Before cycle 1: Baseline blood tests and cardiac assessment are reviewed.

Cycle day 1: Docetaxel + carboplatin + trastuzumab + pertuzumab are given.

Days 2-21: Recovery, supportive medicines and symptom monitoring.

Every 21 days: The cycle repeats if review and blood tests allow.

After chemotherapy: HER2-directed treatment usually continues according to the post-operative/adjuvant treatment plan.

Monitoring and support

- Follow the prescribed steroid premedication around docetaxel.
- Growth-factor support may be part of the plan; follow the exact product and timing given by your oncology team.
- Cardiac function is monitored because trastuzumab/pertuzumab can affect the heart.
- Your current medicines, including non-prescription products.
- A note of new symptoms, when they started and what they stopped you doing.
- Any temperature, blood-pressure or bowel-count readings your team asked you to record.
- Questions about the purpose of treatment, scan/review timing or changes to the plan.

WHEN TO CALL

Call oncology urgently for fever, severe diarrhoea, new breathlessness, swelling, severe rash, uncontrolled vomiting, significant bleeding or a rapidly worsening symptom. Severe breathing difficulty or chest pain: call 112/999. If your team has given you different local thresholds or emergency instructions, follow those instructions.

Consent discussion support

Use the approved consent process to document treatment intent, expected benefit, material risks, alternatives, questions and the patient's decision.

Scan to open the regimen-specific patient portal

<https://sactcheck.com/patient/00722/>

Official NCCP 00722

SACTCheck v0.80.4 - patient information prototype for clinical review. Follow your oncology team and local emergency instructions.

