

Modified FOLFOX-6

Gastrointestinal - NCCP 00209 - version 10a

Your medicines

Oxaliplatin - Folinic acid - 5-fluorouracil (5-FU)

Your treatment schedule

14-day cycle with a 46-hour 5-FU infusion pump.

A picture of side effects

Tell your team about new or worsening symptoms. Numbers link each symptom box to the body map.

Oxaliplatin - nerve and cold effects

5-FU / chemotherapy - other effects

1 Mouth / throat

5-FU / oxaliplatin

Mouth ulcers or soreness; sudden throat tightness needs prompt assessment, including after cold exposure.

3 Blood / infection

Oxaliplatin / 5-FU

Fever, shivers or sudden illness: contact oncology immediately.

5 Hands / nerves

Oxaliplatin

Cold-triggered tingling or numbness; report symptoms that persist between cycles.

7 Skin / mouth

5-fluorouracil (5-FU)

Report significant skin changes or mouth soreness that limits eating or drinking.

2 Heart / breathing

5-fluorouracil (5-FU)

New chest pain, pressure, fainting or breathlessness needs urgent help, including during the infusion.

4 Stomach / bowel

5-FU / oxaliplatin

Persistent diarrhoea, vomiting or inability to drink needs advice.

6 Feet / nerves


Oxaliplatin

Numbness, altered sensation or difficulty walking should be reported before the next dose.

8 Line / home pump

46-hour 5-FU infusion

Call for leakage, disconnection, pain, redness, swelling or a pump alarm. Follow local pump instructions.



Selected effects only. The map does not show likelihood and cannot diagnose the cause of a symptom.

Modified FOLFOX-6

Gastrointestinal - NCCP 00209 - version 10a

Your treatment journey

Day 1: Oxaliplatin and folinic acid are given by drip, then 5-FU is started.

Days 1-3: A portable pump gives 5-FU continuously for about 46 hours.

Around day 3: Pump disconnect. Follow your local line and pump instructions.

Days 4-14: Recovery and monitoring. Symptoms can occur on any day.

Day 15: Next cycle if review and blood tests allow.

Monitoring and support

- DPD status is checked before fluoropyrimidine treatment.
- Before each cycle: blood count, kidney/liver bloods and assessment of nerve symptoms.
- Follow your own labelled anti-sickness, mouth-care and diarrhoea medicines.
- Your current medicines, including non-prescription products.
- A note of new symptoms, when they started and what they stopped you doing.
- Any temperature, blood-pressure or bowel-count readings your team asked you to record.
- Questions about the purpose of treatment, scan/review timing or changes to the plan.

WHEN TO CALL

Call your oncology SOS service for fever or shivers, sudden illness, 4 or more diarrhoea episodes in a day, vomiting despite medicines, mouth sores stopping fluids, bleeding, new chest pain or breathlessness. Severe chest pain or breathing difficulty: call 112/999. If your team has given you different local thresholds or emergency instructions, follow those instructions.

Consent discussion support

Use the approved consent process to document treatment intent, expected benefit, material risks, alternatives, questions and the patient's decision.

Scan to open the regimen-specific patient portal

<https://sactcheck.com/patient/00209/>

Official NCCP 00209 v10a

SACTCheck v0.80.4 - patient information prototype for clinical review. Follow your oncology team and local emergency instructions.

