

Trifluridine/tipiracil (Lonsurf)

Gastrointestinal - NCCP 00382 - version 4

Your medicines

Trifluridine/tipiracil (Lonsurf)

Your treatment schedule

28-day oral cycle: tablets on days 1-5 and 8-12.

A picture of side effects

Tell your team about new or worsening symptoms. Numbers link each symptom box to the body map.

Trifluridine/tipiracil - oral chemotherapy

1 Blood / infection

Trifluridine/tipiracil

Low white cells can increase infection risk. Fever, shivers or sudden illness needs immediate contact.

3 Stomach / bowel

Trifluridine/tipiracil

Nausea, vomiting or diarrhoea can cause dehydration if persistent.

5 Skin

Trifluridine/tipiracil

Report persistent or severe rash or other skin changes.

2 Mouth

Trifluridine/tipiracil

Report mouth soreness or ulcers that limit eating or drinking.

4 Energy / fatigue


Trifluridine/tipiracil

Report sudden or marked tiredness; infection, anaemia or dehydration can also contribute.

6 Tablet handling

Your prescribed tablets

Follow the pharmacy handling and storage instructions. Keep tablets in the original packaging.



Trifluridine/tipiracil (Lonsurf)

Gastrointestinal - NCCP 00382 - version 4

Your treatment journey

- Days 1-5: Take the prescribed dose twice daily.
- Days 6-7: No Lonsurf.
- Days 8-12: Take the prescribed dose twice daily.
- Days 13-28: No Lonsurf; recovery and monitoring period.
- Next cycle: A new 28-day cycle starts only if review and blood tests allow.

Monitoring and support

- Take doses after the morning and evening meals exactly as prescribed.
- Do not double a dose for a missed or vomited dose; follow your oncology/pharmacy instructions.
- Blood counts are particularly important because neutropenia can be significant.
- Your current medicines, including non-prescription products.
- A note of new symptoms, when they started and what they stopped you doing.
- Any temperature, blood-pressure or bowel-count readings your team asked you to record.
- Questions about the purpose of treatment, scan/review timing or changes to the plan.

WHEN TO CALL

Call oncology urgently for fever, shivers, sudden illness, uncontrolled vomiting, persistent diarrhoea, bleeding, inability to drink or marked weakness. Severe breathing difficulty, collapse or severe chest pain: call 112/999. If your team has given you different local thresholds or emergency instructions, follow those instructions.

Consent discussion support

Use the approved consent process to document treatment intent, expected benefit, material risks, alternatives, questions and the patient's decision.

Scan to open the regimen-specific patient portal

<https://sactcheck.com/patient/00382/>
Official NCCP 00382 v4
SACTCheck v0.80.4 - patient information prototype for clinical review. Follow your oncology team and local emergency instructions.

