

Pembrolizumab + pemetrexed + carboplatin

Lung - NCCP 00568 - version 5

Your medicines

Pembrolizumab - Pemetrexed - Carboplatin

Your treatment schedule

Cycles 1-4 every 21 days, then pembrolizumab + pemetrexed maintenance when appropriate.

A picture of side effects

Tell your team about new or worsening symptoms. Numbers link each symptom box to the body map.

Pembrolizumab - immune effects

Pemetrexed / carboplatin - chemotherapy

1 Hormones / brain

Pembrolizumab

Unusual tiredness, dizziness, headache, thirst or passing more urine.

3 Heart

Pembrolizumab

Chest pain, racing heartbeat or fainting needs urgent assessment.

5 Kidneys

Pemetrexed / pembrolizumab

Kidney function is monitored. Report less urine or new swelling.

7 Skin / nerves

Pembrolizumab / chemotherapy

Spreading rash, blisters, weakness or new numbness should be reported.

2 Lungs

Pembrolizumab

New cough or breathlessness: report promptly. Infection, cancer or a clot can cause similar symptoms.

4 Liver

Pembrolizumab

Yellow eyes or skin, dark urine or changes in blood tests.

6 Bowel


Pembrolizumab / chemotherapy

New diarrhoea, persistent vomiting or severe tummy pain: contact your team.

8 Blood / infection

Pemetrexed / carboplatin

Fever, shivers or suddenly feeling unwell: contact oncology immediately.



Selected effects only. The map does not show likelihood and cannot diagnose the cause of a symptom.

Pembrolizumab + pemetrexed + carboplatin

Lung - NCCP 00568 - version 5

Your treatment journey

Before cycle 1: Blood tests, kidney function, glucose and thyroid testing; folic acid and vitamin B12 support for pemetrexed.

Cycles 1-4 - day 1: Pembrolizumab + pemetrexed + carboplatin every 21 days.

Between cycles: Watch for immune symptoms and chemotherapy effects; they can occur at any time.

After cycle 4: Carboplatin stops if the plan is followed and disease/toxicity allow.

Maintenance: Pembrolizumab + pemetrexed every 21 days until progression or unacceptable toxicity, according to the treatment plan.

Monitoring and support

- Take folic acid exactly as prescribed while on pemetrexed and for the required period after it stops.
- Vitamin B12 is given before pemetrexed and then at protocol-defined intervals.
- Blood count, kidney/liver profile and glucose are checked before each cycle; thyroid tests are repeated during treatment.
- Your current medicines, including non-prescription products.
- A note of new symptoms, when they started and what they stopped you doing.
- Any temperature, blood-pressure or bowel-count readings your team asked you to record.
- Questions about the purpose of treatment, scan/review timing or changes to the plan.

WHEN TO CALL

Call oncology urgently for fever, new breathlessness, persistent diarrhoea, jaundice, severe rash, marked weakness, confusion or any rapidly worsening symptom. Severe breathing difficulty, collapse, seizure or severe chest pain: call 112/999. If your team has given you different local thresholds or emergency instructions, follow those instructions.

Consent discussion support

Use the approved consent process to document treatment intent, expected benefit, material risks, alternatives, questions and the patient's decision.

Scan to open the regimen-specific patient portal

<https://sactcheck.com/patient/00568/>

Official NCCP 00568 v5

SACTCheck v0.80.4 - patient information prototype for clinical review. Follow your oncology team and local emergency instructions.

