

Trifluridine/tipiracil (Lonsurf)

Oral anticancer therapy

Where changes may appear

The body systems and labels below are a memory aid. They are regimen/agent-specific and do not predict which effects you will experience.

Trifluridine/tipiracil

Neutropenia / infection	Low neutrophils are a key toxicity. Fever, shivering or sudden illness needs urgent oncology contact.
Anaemia / low platelets	Increasing breathlessness, marked fatigue, unusual bruising or bleeding should be reported.
Nausea, diarrhoea and appetite change	Persistent vomiting/diarrhoea or inability to maintain fluids needs advice.
Fatigue	Record whether fatigue is limiting normal activity and discuss worsening symptoms.

Oral-treatment safety

Dose schedule	Take only on the prescribed treatment days. Do not make up a missed or vomited dose unless your oncology team specifically tells you to.
Medication handling	Keep tablets in their original packaging and follow the handling/storage instructions supplied by your oncology pharmacy.

When to call: follow your oncology SOS instructions for fever, rapidly worsening symptoms, new breathing difficulty, persistent diarrhoea, significant bleeding, severe rash, jaundice, confusion or other concerning changes. For a medical emergency in Ireland call 112/999.



Scan for this regimen portal: <https://sactcheck.com/patient/00382/>

Clinical-review prototype. Your current oncology plan and local emergency instructions take priority.