

Pembrolizumab + pemetrexed + carboplatin

Metastatic non-squamous NSCLC

Where changes may appear

The body systems and labels below are a memory aid. They are regimen/agent-specific and do not predict which effects you will experience.

Pembrolizumab - immune effects

Lung inflammation (pneumonitis)	New cough or breathlessness needs prompt assessment.
Bowel / liver inflammation	Persistent diarrhoea, abdominal pain, yellow eyes or dark urine can signal immune inflammation.
Hormone or sugar changes	Marked fatigue, dizziness, headache, thirst or passing much more urine can reflect thyroid, adrenal, pituitary or glucose problems.
Skin, kidney, nerve or muscle inflammation	Spreading rash, blistering, reduced urine, new weakness, numbness or visual change needs assessment.

Pemetrexed

Kidney toxicity	Kidney function is checked because pemetrexed clearance depends on it. Report reduced urine, swelling or dehydration.
Mouth / gut and skin effects	Mouth soreness, diarrhoea or rash can occur; folic acid and vitamin B12 support are important.

Carboplatin + chemotherapy

Low blood counts / infection	Fever, shivering, bleeding or unusual breathlessness can reflect low blood counts and needs prompt advice.
Nausea, fatigue and appetite change	Use prescribed supportive medicines and contact the team if you cannot maintain fluids.
Hypersensitivity reaction	Rash, wheeze, swelling or dizziness during infusion needs immediate assessment.

When to call: follow your oncology SOS instructions for fever, rapidly worsening symptoms, new breathing difficulty, persistent diarrhoea, significant bleeding, severe rash, jaundice, confusion or other concerning changes. For a medical emergency in Ireland call 112/999.



Scan for this regimen portal: <https://sactcheck.com/patient/00568/>

Clinical-review prototype. Your current oncology plan and local emergency instructions take priority.