

TCHP - docetaxel + carboplatin + trastuzumab + pertuzumab

HER2-positive breast cancer

Where changes may appear

The body systems and labels below are a memory aid. They are regimen/agent-specific and do not predict which effects you will experience.

Trastuzumab + pertuzumab

Heart function change	New breathlessness, ankle swelling, rapid weight gain or palpitations should be reported; heart function is monitored.
Diarrhoea	Pertuzumab-containing treatment can cause diarrhoea. Persistent or severe diarrhoea needs advice to prevent dehydration.
Infusion reaction	Fever, chills, flushing, wheeze or dizziness during/after infusion needs immediate assessment.

Docetaxel

Peripheral neuropathy	Tingling, numbness or pain in hands/feet should be reported before it interferes with function.
Fluid retention / skin and nails	New swelling, rapid weight gain, painful rash or significant nail changes should be discussed.
Hypersensitivity reaction	Rash, wheeze, swelling or dizziness during infusion needs immediate assessment.

Carboplatin + regimen-wide

Low blood counts / infection	Fever, shivering, bleeding or sudden illness needs prompt oncology contact.
Nausea and fatigue	Use prescribed supportive medicines and seek advice if unable to maintain fluids.
Carboplatin hypersensitivity	Allergic reactions can occur, particularly after repeated exposure; tell staff immediately about symptoms during infusion.

When to call: follow your oncology SOS instructions for fever, rapidly worsening symptoms, new breathing difficulty, persistent diarrhoea, significant bleeding, severe rash, jaundice, confusion or other concerning changes. For a medical emergency in Ireland call 112/999.



Scan for this regimen portal: <https://sactcheck.com/patient/00722/>

Clinical-review prototype. Your current oncology plan and local emergency instructions take priority.