Carry your treatment details

Which treatment this guide covers NCCP 00219a is first-line EGFR-mutated advanced NSCLC; 00219b is switch-maintenance after stable disease following first-line chemotherapy; 00219c is after failure of at least one previous chemotherapy regimen. Global NCCP eligibility requires a validated activating EGFR mutation in each setting. These are advanced-disease settings, not adjuvant treatment.

Print, cut out and fold. Follow your treating hospital’s contact plan.

SACTCHECK · TREATMENT CARD

Erlotinib monotherapy

NCCP 00219 v6

The usual dose is 150 mg by mouth once daily, continuously without a planned rest week. Use the dose on your prescription. Treatment continues while it helps and side effects remain acceptable; any change is agreed with your team.

Prescriber / next review: _____

Contact plan

Oncology team: Insert numbers here · Use your hospital’s opening hours.
Out of hours: Insert numbers here.
Evenings / overnight: agreed out-of-hours service or emergency department.
Emergency: 112 / 999.

Own hospital / contact: _____

Call urgently for new or worsening breathlessness, cough or fever; severe diarrhoea, inability to drink or much less urine; severe or persistent abdominal pain; blistering or peeling skin; or yellow skin/eyes. Eye pain, light sensitivity or blurred vision needs prompt assessment. Severe breathing difficulty, collapse or swelling of the tongue/throat needs emergency help. Use your unit’s temperature and 24-hour contact instructions.

https://sactcheck.com/patient/00219/ · no patient information encoded.

SACTCheck · Developed by Dr Paul O’Brien · v0.84.26 · Sources checked 2026-10-29
Source reconciled; independent clinical, pharmacy and patient/PPI review pending