Which treatment this guide covers NCCP 00220a covers adults with locally advanced or metastatic NSCLC with an activating EGFR mutation. This is continuous treatment for advanced disease, not a fixed adjuvant course.

NCCP 00220 · VERSION 5

Gefitinib monotherapy

The usual dose is 250 mg by mouth once daily, continuously without a planned rest week. Use the dose on your prescription. Treatment continues while it helps and side effects remain acceptable; any change is agreed with your team.

Your treatment and review passport

Bring this to reviews. Record symptoms, treatment changes and questions when helpful. You do not need to complete a daily diary or calculate a toxicity grade.

The four sheets are reusable review periods, not a prescribed number of cycles. Follow the treatment and rest days agreed with your team; do not calculate a restart from this sheet.

These patient-language prompts are not a validated questionnaire or a formal CTCAE assessment. Urgent symptoms override the form.

Emergency: 112 / 999. Severe chest pain, major breathing difficulty, collapse or stroke symptoms need emergency help.

Contact your treating team promptly: Call urgently for new or worsening breathlessness, cough or fever; severe diarrhoea, inability to drink or much less urine; severe or persistent abdominal pain; blistering or peeling skin; or yellow skin/eyes. Eye pain, light sensitivity or blurred vision needs prompt assessment. Severe breathing difficulty, collapse or swelling of the tongue/throat needs emergency help. Use your unit’s temperature and 24-hour contact instructions.

YOUR ONCOLOGY TEAM · DAYTIME

Insert numbers here

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

Insert numbers here

Ask for medical oncology

EVENINGS / OVERNIGHT

Your agreed out-of-hours pathway

Other hospitals: use the contacts your own team provides.

Your complete treatment reference

https://sactcheck.com/patient/00220/

Public information only. No personal information is encoded.

Source reconciled; independent clinical, pharmacy and patient/PPI review pending

Gefitinib monotherapy

Review period / visit: _____ Next review: _____

Medicine-specific symptom	No concern	Present / manageable	Affects sleep or daily activities — contact team	When / what changed?
Diarrhoea	☐	☐	☐	
Skin / nail changes	☐	☐	☐	
Mouth soreness / ulcers	☐	☐	☐	
New cough or breathlessness	☐	☐	☐	
Eye pain or blurred vision	☐	☐	☐	
Fatigue	☐	☐	☐	

Checks: expected, booked and reviewed are different

Test / review	Due window	Who books it?	Who reviews the result?	Outcome / next action
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____

Missed tablets / injection changes / questions / support needed:

> Call urgently for new or worsening breathlessness, cough or fever; severe diarrhoea, inability to drink or much less urine; severe or persistent abdominal pain; blistering or peeling skin; or yellow skin/eyes. Eye pain, light sensitivity or blurred vision needs prompt assessment. Severe breathing difficulty, collapse or swelling of the tongue/throat needs emergency help. Use your unit’s temperature and 24-hour contact instructions. Emergency symptoms: 112 / 999.

Gefitinib monotherapy

Review period / visit: _____ Next review: _____

Medicine-specific symptom	No concern	Present / manageable	Affects sleep or daily activities — contact team	When / what changed?
Diarrhoea	☐	☐	☐	
Skin / nail changes	☐	☐	☐	
Mouth soreness / ulcers	☐	☐	☐	
New cough or breathlessness	☐	☐	☐	
Eye pain or blurred vision	☐	☐	☐	
Fatigue	☐	☐	☐	

Checks: expected, booked and reviewed are different

Test / review	Due window	Who books it?	Who reviews the result?	Outcome / next action
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____

Missed tablets / injection changes / questions / support needed:

> Call urgently for new or worsening breathlessness, cough or fever; severe diarrhoea, inability to drink or much less urine; severe or persistent abdominal pain; blistering or peeling skin; or yellow skin/eyes. Eye pain, light sensitivity or blurred vision needs prompt assessment. Severe breathing difficulty, collapse or swelling of the tongue/throat needs emergency help. Use your unit’s temperature and 24-hour contact instructions. Emergency symptoms: 112 / 999.

Gefitinib monotherapy

Review period / visit: _____ Next review: _____

Medicine-specific symptom	No concern	Present / manageable	Affects sleep or daily activities — contact team	When / what changed?
Diarrhoea	☐	☐	☐	
Skin / nail changes	☐	☐	☐	
Mouth soreness / ulcers	☐	☐	☐	
New cough or breathlessness	☐	☐	☐	
Eye pain or blurred vision	☐	☐	☐	
Fatigue	☐	☐	☐	

Checks: expected, booked and reviewed are different

Test / review	Due window	Who books it?	Who reviews the result?	Outcome / next action
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____

Missed tablets / injection changes / questions / support needed:

> Call urgently for new or worsening breathlessness, cough or fever; severe diarrhoea, inability to drink or much less urine; severe or persistent abdominal pain; blistering or peeling skin; or yellow skin/eyes. Eye pain, light sensitivity or blurred vision needs prompt assessment. Severe breathing difficulty, collapse or swelling of the tongue/throat needs emergency help. Use your unit’s temperature and 24-hour contact instructions. Emergency symptoms: 112 / 999.

Gefitinib monotherapy

Review period / visit: _____ Next review: _____

Medicine-specific symptom	No concern	Present / manageable	Affects sleep or daily activities — contact team	When / what changed?
Diarrhoea	☐	☐	☐	
Skin / nail changes	☐	☐	☐	
Mouth soreness / ulcers	☐	☐	☐	
New cough or breathlessness	☐	☐	☐	
Eye pain or blurred vision	☐	☐	☐	
Fatigue	☐	☐	☐	

Checks: expected, booked and reviewed are different

Test / review	Due window	Who books it?	Who reviews the result?	Outcome / next action
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____
_____	_____	_____	_____	_____

Missed tablets / injection changes / questions / support needed:

> Call urgently for new or worsening breathlessness, cough or fever; severe diarrhoea, inability to drink or much less urine; severe or persistent abdominal pain; blistering or peeling skin; or yellow skin/eyes. Eye pain, light sensitivity or blurred vision needs prompt assessment. Severe breathing difficulty, collapse or swelling of the tongue/throat needs emergency help. Use your unit’s temperature and 24-hour contact instructions. Emergency symptoms: 112 / 999.