Carry your treatment details

Which treatment this guide covers NCCP 00220a covers adults with locally advanced or metastatic NSCLC with an activating EGFR mutation. This is continuous treatment for advanced disease, not a fixed adjuvant course.

Print, cut out and fold. Follow your treating hospital’s contact plan.

SACTCHECK · TREATMENT CARD

Gefitinib monotherapy

NCCP 00220 v5

The usual dose is 250 mg by mouth once daily, continuously without a planned rest week. Use the dose on your prescription. Treatment continues while it helps and side effects remain acceptable; any change is agreed with your team.

Prescriber / next review: _____

Contact plan

Oncology team: Insert numbers here · Use your hospital’s opening hours.
Out of hours: Insert numbers here.
Evenings / overnight: agreed out-of-hours service or emergency department.
Emergency: 112 / 999.

Own hospital / contact: _____

Call urgently for new or worsening breathlessness, cough or fever; severe diarrhoea, inability to drink or much less urine; severe or persistent abdominal pain; blistering or peeling skin; or yellow skin/eyes. Eye pain, light sensitivity or blurred vision needs prompt assessment. Severe breathing difficulty, collapse or swelling of the tongue/throat needs emergency help. Use your unit’s temperature and 24-hour contact instructions.

https://sactcheck.com/patient/00220/ · no patient information encoded.

SACTCheck · Developed by Dr Paul O’Brien · v0.84.26 · Sources checked 2026-10-29
Source reconciled; independent clinical, pharmacy and patient/PPI review pending