# Atezolizumab 1200 mg · every 3 weeks · lung cancer

NCCP 00544 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

## Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

### Appointments

Atezolizumab 1200 mg is given by intravenous infusion on Day 1 every 3 weeks (21 days). The first infusion lasts about 60 minutes; later infusions may last 30 minutes if tolerated. The visit includes checks and preparation. This protocol also offers a separately prescribed 1875 mg injection under the skin of the thigh over about 7 minutes every 3 weeks. The IV and under-skin doses are not interchangeable milligram for milligram.

STEP 2

### Course length

Adjuvant treatment lasts up to 12 months. Advanced NSCLC and ES-SCLC maintenance continue until progression or unacceptable toxicity, with regular individual review.

STEP 3

### Checks between visits

Blood tests are due before each infusion. Thyroid tests stay every 3–6 weeks regardless of dose schedule. Ask about travel, work, transport and whom to contact if a test or visit is delayed.

STEP 4

### Home medicines and records

Keep the alert card, symptom record, current medicine list and urgent contact details accessible. Your team may prescribe steroids or hormone replacement only if needed.

STEP 5

### After the final dose

Keep reporting new symptoms. Some hormone medicines and monitoring continue after the cancer treatment finishes.

Sources: NCCP 00544 v14 · official regimen. Source reconciled; independent clinical, pharmacy and patient/PPI review pending.