Which treatment this guide covers NCCP 00220a covers adults with locally advanced or metastatic NSCLC with an activating EGFR mutation. This is continuous treatment for advanced disease, not a fixed adjuvant course.

NCCP 00220 · VERSION 5

Gefitinib monotherapy

The usual dose is 250 mg by mouth once daily, continuously without a planned rest week. Use the dose on your prescription. Treatment continues while it helps and side effects remain acceptable; any change is agreed with your team.

Discussion aid: review with the patient and complete the approved local consent process.

Purpose and plan

☐ Exact indication, treatment aim and uncertainty explained

☐ Expected duration and review / switch points discussed

☐ Alternatives and the option not to proceed discussed

☐ Patient priorities, capacity and support needs explored

Practical safety

☐ Medicine, route, schedule and missed-dose instructions

☐ Food, interactions and safe handling reviewed

☐ Pregnancy, contraception / fertility and breastfeeding

☐ Monitoring, symptom reporting and contact plan

Selected material risks

☐ Diarrhoea and dehydration

☐ Serious lung inflammation

☐ Eye inflammation

☐ Skin, nail and mouth changes

☐ Liver injury

☐ Rare bowel perforation and serious skin reactions

Discuss the aim of advanced cancer control, expected benefit and uncertainty, alternatives including not proceeding, bowel and skin effects, serious lung/liver problems, fertility and pregnancy, daily treatment, monitoring and your right to reconsider.

Individual benefit, alternatives and concerns

Agreed checks and next review

NCCP specifies blood count, kidney and liver tests throughout treatment as clinically indicated. The team sets your appointments and scan plan. Report symptoms promptly; extra checks may be needed for diarrhoea, dehydration or liver problems.

Clinician / date: _____ Patient discussion documented: _____

Consent is an ongoing conversation and may be reconsidered. This aid is not treatment authorisation.