NCCP 00689 · clinician draft

Atezolizumab + carboplatin + etoposide

Supportive prescription draft for review. This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Induction Days 2–3 · etoposide and post-treatment cover

Active medicines: etoposide

Emetogenic risk: low

Medicine	Dose / route	When
Low-risk antiemetic reference	Ondansetron 8 mg orally once daily where selected	Before IV etoposide Days 2–3; review coverage through Day 5 NCIS low-risk row is a reference, not an additional automatic dose on top of Day 1 prophylaxis. Reconcile with ongoing aprepitant/dexamethasone and any 5-HT3 doses. Oral etoposide alternative requires its explicit local prescription.
Rescue antiemetic	Individual prescription	During treatment and later days NCIS low-risk row lists metoclopramide 10 mg orally three times daily as needed; account for any already scheduled doses and clinical suitability.

Additional phase-specific support

- NCIS multi-day principle: match prophylaxis to the drugs on each treatment day and continue appropriate cover for two days after.
- Etoposide Days 2–3 may be IV or an explicitly prescribed oral twice-daily alternative; no silent conversion.
- Issue a single reconciled day-by-day plan; do not start a fresh carboplatin high-risk course on each etoposide-only day.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00689 v4 · p2–3,10 · NCCP NCIS Medical Oncology antiemetics V8

Sources checked 2026-10-29 · SACTCheck v0.84.26 · Regimen-level drafts; prescriber and oncology-pharmacy verification required