NCCP 00857 · clinician draft

Pembrolizumab + carboplatin/paclitaxel → AC

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Phase 1 · Days 8 and 15 paclitaxel

Active medicines: Paclitaxel

Emetogenic risk: Low

Medicine	Dose / route	When
Ondansetron	8 mg orally	On treatment day Review QT, constipation and hepatic dosing; reconcile any steroid already prescribed
Metoclopramide	10 mg orally up to three times daily as needed	Subsequent days: duration to be confirmed Assess contraindications, renal function and interactions

Additional phase-specific support

- Use the 00857 paclitaxel premedication / prior-reaction plan.
- Carboplatin Day 1 delayed antiemetics are not repeated automatically on these visits.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00857 v3 · NCCP antiemetics v8, Table 1 pp4–5

Sources checked 2026-10-17 · SACTCheck v0.83.8 · Regimen-level drafts; prescriber and oncology-pharmacy verification required