Adjuvant ribociclib + endocrine therapy

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Adjuvant ribociclib + endocrine therapy

Active medicines: Ribociclib + Non-steroidal aromatase inhibitor ± ovarian suppression

Emetogenic risk: Oral minimal to low

No routine prophylactic antiemetic order is populated for this phase. Agree an individual symptom-treatment plan where needed.

Additional phase-specific support

- No default QT-prolonging antiemetic is generated. Pharmacy must choose symptom treatment after reviewing ECG, electrolytes and interacting medicines.
- Ribociclib Days 1–21; Days 22–28 off. Endocrine therapy continues separately.
- Bloods every 2 weeks for cycles 1–2, then before the next four cycles; ECG baseline and around C1D14; further checks as indicated.
- The optional local D25–26 review window does not replace early checks or authorise restarting.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00892 v2 · p6 · NCCP antiemetics v8, Table 1 pp4–5

Sources checked 2026-10-17 · SACTCheck v0.83.8 · Regimen-level drafts; prescriber and oncology-pharmacy verification required