Ribociclib + endocrine therapy · metastatic

Ribociclib is taken once daily on Days 1–21, followed by seven days off, in a 28-day cycle. The endocrine partner continues according to its own prescription. Treatment continues while it is helping and tolerable.

Draft — independent clinical, pharmacy and patient/PPI review pending. Selected effects; not a complete medicine leaflet.

A picture of side effects

Numbered symptom boxes connect to the anatomical illustration. The medicine responsible may be uncertain; report the symptom.

1 Heart rhythm / QT


Ribociclib

Baseline ECG and an ECG around Cycle 1 Day 14. Fainting or palpitations need urgent advice.

2 Liver injury

Ribociclib

Keep liver blood tests. Yellow skin / eyes, dark urine or marked upper-abdominal discomfort: call promptly.



3 Low neutrophils / infection

Ribociclib

Blood counts guide whether the next cycle can start. Fever, shivers or sudden illness: contact oncology immediately.

4 Lung inflammation

Ribociclib · uncommon but important

New cough or breathlessness needs prompt assessment, even between visits.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

> Fever, shivers or suddenly unwell: contact oncology immediately. Severe chest pain, breathing difficulty, collapse or seizure: 112 / 999.

NCCP 00525 · PRACTICAL SUPPORT

Your treatment, at home.

Important medicine-specific effects

Ribociclib

- Low white cells and infection
- Liver injury / abnormal blood tests
- QT prolongation / heart-rhythm problems
- Nausea, fatigue and hair thinning
- Rare serious lung or skin reactions

AI or fulvestrant

- Hot flushes, joint pain and fatigue
- Bone loss with AI / ovarian suppression
- Injection-site discomfort with fulvestrant
- Drug-specific liver and other effects

Surgery, radiation and monitoring: open the whole-treatment map or scan the QR.

Make the plan practical

Three weeks on, one off

The ribociclib break is Days 22–28. Your endocrine partner usually continues.

Bloods and ECG

Early blood tests and a Day-14 ECG are important even if you feel well.

Medicines check

Avoid grapefruit; check new prescriptions, over-the-counter medicines and supplements for interactions.

Missed or vomited tablets

Do not take an extra dose. Take the next prescribed dose at the usual time and contact the team if uncertain.

Emergency: 112 / 999. Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

Contact oncology immediately for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital's instructions. If you cannot reach the team, seek urgent medical assessment.

CUH · MON–FRI 8AM–5PM

087 181 9645

Acute Oncology Service

CUH · WEEKENDS / BANK HOLIDAYS 9AM–5PM

021 4922000

Ask for medical oncology

EVENINGS / OVERNIGHT

Your out-of-hours pathway / nearest ED

Other hospitals: use the numbers your team provides.

Complete reference, visual timeline and passport

https://sactcheck.com/patient/00525/

Sources and review status are available online. Follow your own prescription and hospital instructions.