Adjuvant ribociclib + endocrine therapy

NCCP 00892 · Draft — independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

Your practical treatment plan

Ribociclib 400 mg once daily on Days 1–21, followed by Days 22–28 off, for up to 3 years. The aromatase inhibitor continues daily on its own schedule; ovarian suppression is added when indicated.

STEP 1

Before dispensing

Confirm blood tests, interactions, the exact tablet strengths and the plan for missed doses.

STEP 2

Taking tablets

Ribociclib 400 mg once daily on Days 1–21, followed by Days 22–28 off, for up to 3 years. The aromatase inhibitor continues daily on its own schedule; ovarian suppression is added when indicated.

STEP 3

Review before restarting

The team confirms blood results, side effects and the next prescription. Never double a missed dose or restart held treatment without advice.

Missed or vomited dose

Do not take an extra dose. Follow the medicine-specific instructions and ask the team if uncertain.

Safe handling

Swallow tablets whole, keep them in their original packaging away from children, and return unused tablets to pharmacy.

Practical arrangements

Confirm who supplies the tablets, who arranges tests, who reviews results and your urgent contact route.

Sources: NCCP 00892 v2 · official regimen · Novartis · Kisqali product information, updated September 2026. Draft — independent clinical, pharmacy and patient/PPI review pending.