

Attractor structure of Boolean networks of small canalizing depth

Gleb Pogudin,

MAX team, LIX, CNRS, École Polytechnique, Institut Polytechnique de Paris,

joint work with R. Laubenbacher, E. Paul, and W. Qin



Plan of the talk

Main question: difference in dynamics between
“a random Boolean network” and “random **canalizing** Boolean network”

Plan

- Intro and problem statement
- Simulation results
- Theoretic results
- Remarks on the computation

Boolean networks and their structure

Definition

\mathbf{F}_2 is a field with two elements 0 (false / off) and 1 (true / on) and arithmetic operations:

$$x + y := x \text{ XOR } y, \quad xy := x \text{ AND } y.$$

Boolean networks and their structure

Definition

\mathbb{F}_2 is a field with two elements 0 (false / off) and 1 (true / on) and arithmetic operations:

$$x + y := x \text{ XOR } y, \quad xy := x \text{ AND } y.$$

An n -dimensional **Boolean network** defined by n polynomials $f_1, \dots, f_n \in \mathbb{F}_2[x_1, \dots, x_n]$ is a discrete-time dynamical system such that

- states are points in \mathbb{F}_2^n
- and, the the state at time t is $\mathbf{a}_t = (a_{t,1}, \dots, a_{t,n}) \in \mathbb{F}_2^n$, then

$$\mathbf{a}_{t+1} = (f_1(\mathbf{a}_t), \dots, f_n(\mathbf{a}_t)).$$

Boolean networks and their structure

Definition

\mathbb{F}_2 is a field with two elements 0 (false / off) and 1 (true / on) and arithmetic operations:

$$x + y := x \text{ XOR } y, \quad xy := x \text{ AND } y.$$

An n -dimensional Boolean network defined by n polynomials $f_1, \dots, f_n \in \mathbb{F}_2[x_1, \dots, x_n]$ is a discrete-time dynamical system such that

- states are points in \mathbb{F}_2^n
- and, the the state at time t is $\mathbf{a}_t = (a_{t,1}, \dots, a_{t,n}) \in \mathbb{F}_2^n$, then

$$\mathbf{a}_{t+1} = (f_1(\mathbf{a}_t), \dots, f_n(\mathbf{a}_t)).$$

Example

Let $f_1(x_1, x_2) = x_1(x_2 + 1)$ and $f_2(x_1, x_2) = (x_1 + 1)x_2$:

$$(0, 1) \rightarrow (1, 0) \rightarrow (0, 1) \rightarrow \dots$$

Boolean networks and their structure

Definition

An n -dimensional **Boolean network** defined by n polynomials $f_1, \dots, f_n \in \mathbb{F}_2[x_1, \dots, x_n]$ is a discrete-time dynamical system such that

- states are points in \mathbb{F}_2^n
- and, the the state at time t is $\mathbf{a}_t = (a_{t,1}, \dots, a_{t,n}) \in \mathbb{F}_2^n$, then

$$\mathbf{a}_{t+1} = (f_1(\mathbf{a}_t), \dots, f_n(\mathbf{a}_t)).$$

Why: regulatory networks.

Boolean networks and their structure

Definition

An n -dimensional **Boolean network** defined by n polynomials $f_1, \dots, f_n \in \mathbb{F}_2[x_1, \dots, x_n]$ is a discrete-time dynamical system such that

- states are points in \mathbb{F}_2^n
- and, the the state at time t is $\mathbf{a}_t = (a_{t,1}, \dots, a_{t,n}) \in \mathbb{F}_2^n$, then

$$\mathbf{a}_{t+1} = (f_1(\mathbf{a}_t), \dots, f_n(\mathbf{a}_t)).$$

Why: regulatory networks.

Why polynomials?

\mathbb{F}^n is a finite set \implies **any** function is polynomial

Boolean networks and their structure

Definition

An n -dimensional **Boolean network** defined by n polynomials $f_1, \dots, f_n \in \mathbb{F}_2[x_1, \dots, x_n]$ is a discrete-time dynamical system such that

- states are points in \mathbb{F}_2^n
- and, the the state at time t is $\mathbf{a}_t = (a_{t,1}, \dots, a_{t,n}) \in \mathbb{F}_2^n$, then

$$\mathbf{a}_{t+1} = (f_1(\mathbf{a}_t), \dots, f_n(\mathbf{a}_t)).$$

Why: regulatory networks.

Why polynomials?

\mathbb{F}^n is a finite set \implies **any** function is polynomial

Pros: we can define any dynamics!

Boolean networks and their structure

Definition

An n -dimensional **Boolean network** defined by n polynomials $f_1, \dots, f_n \in \mathbb{F}_2[x_1, \dots, x_n]$ is a discrete-time dynamical system such that

- states are points in \mathbb{F}_2^n
- and, the the state at time t is $\mathbf{a}_t = (a_{t,1}, \dots, a_{t,n}) \in \mathbb{F}_2^n$, then

$$\mathbf{a}_{t+1} = (f_1(\mathbf{a}_t), \dots, f_n(\mathbf{a}_t)).$$

Why: regulatory networks.

Why polynomials?

\mathbb{F}^n is a finite set \implies **any** function is polynomial

Pros: we can define any dynamics!

Cons: we can define any dynamics...


Dynamics: attractors and basins

n -dimensional **Boolean network** \implies directed **graph** on 2^n vertices

Network:

$$f_1 = x_1(x_2 + 1),$$

$$f_2 = (x_1 + 1)x_2.$$



Dynamics: attractors and basins

n -dimensional **Boolean network** \implies directed **graph** on 2^n vertices


Network:

$$f_1 = x_1(x_2 + 1),$$

$$f_2 = (x_1 + 1)x_2.$$

Notions

- **Attractors** are **cycles** in the graph.



Dynamics: attractors and basins

n -dimensional **Boolean network** \implies directed **graph** on 2^n vertices


Network:

$$f_1 = x_1(x_2 + 1),$$

$$f_2 = (x_1 + 1)x_2.$$

Notions

- **Attractors** are **cycles** in the graph.
- **Steady states** are **loops** in the graph.




Dynamics: attractors and basins

n -dimensional **Boolean network** \implies directed **graph** on 2^n vertices

Network:

$$f_1 = x_1(x_2 + 1),$$

$$f_2 = (x_1 + 1)x_2.$$



Notions

- **Attractors** are **cycles** in the graph.
- **Steady states** are **loops** in the graph.
- **Basins** are **connected components**.
Each basin corresponds to an attractor and each attractor has its basin.


Dynamics: attractors and basins

n -dimensional **Boolean network** \implies directed **graph** on 2^n vertices

Network:

$$f_1 = x_1(x_2 + 1),$$

$$f_2 = (x_1 + 1)x_2.$$



Notions

- **Attractors** are **cycles** in the graph.
- **Steady states** are **loops** in the graph.
- **Basins** are **connected components**.
Each basin corresponds to an attractor and each attractor has its basin.

Intuition

Attractors \leftrightarrow Cell types

Length \leftrightarrow number of expression patterns

Canalization

One variable to rule them all

A nonconstant function $f(x_1, \dots, x_n)$ is **canalizing** with respect to a variable x_i if there exists a **canalizing value** $a \in \{0, 1\}$ such that

$$f(x_1, \dots, x_{i-1}, a, x_{i+1}, \dots, x_n) \equiv \text{const.}$$

Canalization

One variable to rule them all

A nonconstant function $f(x_1, \dots, x_n)$ is **canalizing** with respect to a variable x_i if there exists a **canalizing value** $a \in \{0, 1\}$ such that

$$f(x_1, \dots, x_{i-1}, a, x_{i+1}, \dots, x_n) \equiv \text{const.}$$

Example

$f(x_1, x_2) = x_1(x_2 + 1)$ is canalizing w.r.t

x_1 with value 0

x_2 with value 1

Canalization

One variable to rule them all

A nonconstant function $f(x_1, \dots, x_n)$ is **canalizing** with respect to a variable x_i if there exists a **canalizing value** $a \in \{0, 1\}$ such that

$$f(x_1, \dots, x_{i-1}, a, x_{i+1}, \dots, x_n) \equiv \text{const.}$$

Example

$f(x_1, x_2) = x_1(x_2 + 1)$ is canalizing w.r.t

x_1 with value 0

x_2 with value 1

Why?

- Often appear in the published models.
- Make sense.

Canalization

One variable to rule them all

A nonconstant function $f(x_1, \dots, x_n)$ is **canalizing** with respect to a variable x_i if there exists a **canalizing value** $a \in \{0, 1\}$ such that

$$f(x_1, \dots, x_{i-1}, a, x_{i+1}, \dots, x_n) \equiv \text{const.}$$

Canalizing depth (Layne, Dimitrova, Macaulay, 2012)

We define canalizing depth recursively

1. noncanalizing function \implies depth 0
2. if f is canalizing w.r.t. x_1 with value a , then

$$\text{depth}(f) = \text{depth}(f|_{x_1=1-a}) + 1.$$

Main question

Question: How does the canalization (and its depth) affect the attractor structure?

Question: How does the canalization (and its depth) affect
the attractor structure?

Related results include: no chaos (Karlsson, Hörnquist), stability (Kauffman, Peterson, Samuelsson, Troein and Layne, Dimitrova, Macauley)

Experimental results

Setup

For each $n = 4, \dots, 20$ and each $k = 0, \dots, n$ we generate random Boolean networks

- of dimension n ;
- with all the functions having canalizing depth k .

Experimental results

Setup

For each $n = 4, \dots, 20$ and each $k = 0, \dots, n$ we generate random Boolean networks

- of dimension n ;
- with all the functions having canalizing depth k .

Computed values

For each Boolean network we compute



- N:** the number of attractors
- S:** the total number of states in the attractors
- AS:** the average attractor size $= \frac{S}{N}$

Experimental results

For $n = 15$ based in 50k samples for each k .

$N = \#$ attractors

$AS = \#$ avg. attractor size





Experimental results

For $n = 15$ based in 50k samples for each k .

$N = \#$ attractors

$AS = \#$ avg. attractor size



Analysis



- Depth increases \implies number and sizes decrease

Experimental results

For $n = 15$ based in 50k samples for each k .

$N = \#$ attractors

$AS = \#$ avg. attractor size



Analysis



- Depth increases \implies number and sizes decrease
- Sizes decrease more substantially

Experimental results

For $n = 15$ based in 50k samples for each k .

$N = \#$ attractors

$AS = \#$ avg. attractor size



Analysis



- Depth increases \implies number and sizes decrease
- Sizes decrease more substantially
- Most of the change happens between $k = 0$ and $k = 3$.

Experimental results

For $n = 12$ based in 300k samples for each k .

$N = \#$ attractors

$AS = \#$ avg. attractor size





Experimental results

For $n = 12$ based in 300k samples for each k .

$N = \#$ attractors

$AS = \#$ avg. attractor size



Analysis

- Confirms earlier conclusions
- Avg. size differs dramatically for $k = 0$ and $k = 1$

Theoretical results

Quantity of interest

$A_{\ell,n}$:= expected # of attractors of length ℓ in a random network of dimension n and depth **one**.

$$A_\ell := \lim_{n \rightarrow \infty} A_{\ell,n}$$

Theoretical results

Quantity of interest

$A_{\ell,n}$:= expected # of attractors of length ℓ in a random network of dimension n and depth **one**.

$$A_\ell := \lim_{n \rightarrow \infty} A_{\ell,n}$$

What if depth = 0?

The corresponding limit will be $\frac{1}{\ell}$

Theoretical results

Quantity of interest

$A_{\ell,n}$:= expected # of attractors of length ℓ in a random network of dimension n and depth **one**.

$$A_\ell := \lim_{n \rightarrow \infty} A_{\ell,n}$$

What if depth = 0?

The corresponding limit will be $\frac{1}{\ell}$

Theorem

We give a formula for A_ℓ , in particular

- $A_\ell > \frac{1}{\ell}$ for $\ell > 0$;
- $A_1 = 1$, $A_2 = 0.66 \dots$, $A_3 = 0.3386 \dots$, $A_4 = 0.2856 \dots$

Summary

- canalization \implies fewer attractors and they are much smaller
Biologically relevant!

Summary

- canalization \implies fewer attractors and they are much smaller
Biologically relevant!
- no big difference between small (but > 0) and large canalizing depths
Applicable to a wide range of models

Remarks on the computation

Sampling:

- Relies on the structure theorem by He and Macaulay
- Complexity $\mathcal{O}(n2^n)$ almost optimal
more precisely $\mathcal{O}(k^3 + (n - k)2^{n-k} + k2^n)$

Finding attractors: DFS

Open problems

Taking into account:

- sparsity
- uneven canalization profiles
(i.e. only some of the functions are canalizing)
- collective canalization

This work was partially supported by the National Science Foundation.

