




# Subspace methods for line spectra



Per Mattsson

Systems and Control  
Department of Information Technology  
Uppsala University

2019-09-19

# Summary from last lecture

---

- ▶ Signal model for line spectra
- ▶ Estimating the spectrum  $\implies$  frequency estimation
- ▶ Nonlinear least squares
- ▶ Models of the covariance matrix and ARMA-type model
- ▶ Singular value decomposition (SVD)

# Today

---

- ▶ A recap from L5
- ▶ Subspace methods for frequency estimation
  - ▶ High order Yule-Walker
  - ▶ MUSIC
  - ▶ Min-Norm
  - ▶ ESPRIT
- ▶ Makes use of the covariance structure

# Line spectra

---

Signal model:

$$y(t) = \sum_{k=1}^n \alpha_k e^{i(\omega_k t + \varphi_k)} + e(t)$$

# Line spectra

---

Signal model:

$$y(t) = \sum_{k=1}^n \alpha_k e^{i(\omega_k t + \varphi_k)} + e(t)$$

Theoretical ACS for line spectra

$$r(k) = E\{y(t)y^*(t-k)\} = \sum_{p=1}^n \alpha_p^2 e^{i\omega_p k} + \sigma^2 \delta_{k,0}$$

# Line spectra

---

Signal model:

$$y(t) = \sum_{k=1}^n \alpha_k e^{i(\omega_k t + \varphi_k)} + e(t)$$

Theoretical ACS for line spectra

$$r(k) = E\{y(t)y^*(t-k)\} = \sum_{p=1}^n \alpha_p^2 e^{i\omega_p k} + \sigma^2 \delta_{k,0}$$

Theoretical spectrum

$$\phi(\omega) = 2\pi \sum_{p=1}^n \alpha_p^2 \delta(\omega - \omega_p) + \sigma^2$$

# Frequency estimation

---

## Problem

Determine  $\{\omega_k\}_{k=1}^n$  from observations or samples  $\{y(t)\}_{t=1}^N$

# Frequency estimation

## Problem

Determine  $\{\omega_k\}_{k=1}^n$  from observations or samples  $\{y(t)\}_{t=1}^N$

- ▶ The nonlinear least square method,

$$\min_{\omega, \alpha, \varphi} \sum_{k=1}^N \left| y(t) - \sum_{k=1}^n \alpha_k e^{i(\omega_k t + \varphi_k)} \right|^2$$



# Frequency estimation

## Problem

Determine  $\{\omega_k\}_{k=1}^n$  from observations or samples  $\{y(t)\}_{t=1}^N$

- ▶ The nonlinear least square method,

$$\min_{\omega, \alpha, \varphi} \sum_{k=1}^N \left| y(t) - \sum_{k=1}^n \alpha_k e^{i(\omega_k t + \varphi_k)} \right|^2$$

- ▶ Excellent properties if the global minimum can be found.

# Frequency estimation

## Problem

Determine  $\{\omega_k\}_{k=1}^n$  from observations or samples  $\{y(t)\}_{t=1}^N$

- ▶ The nonlinear least square method,

$$\min_{\omega, \alpha, \varphi} \sum_{k=1}^N \left| y(t) - \sum_{k=1}^n \alpha_k e^{i(\omega_k t + \varphi_k)} \right|^2$$

- ▶ Excellent properties if the global minimum can be found.
- ▶ Hard (impossible) to use in general.

# Frequency estimation

## Problem

Determine  $\{\omega_k\}_{k=1}^n$  from observations or samples  $\{y(t)\}_{t=1}^N$

- ▶ The nonlinear least square method,

$$\min_{\omega, \alpha, \varphi} \sum_{k=1}^N \left| y(t) - \sum_{k=1}^n \alpha_k e^{i(\omega_k t + \varphi_k)} \right|^2$$

- ▶ Excellent properties if the global minimum can be found.
- ▶ Hard (impossible) to use in general.
- ▶ Today we will instead use subspace methods.

# Singular value decomposition (SVD)

## Definition

Factor a matrix  $A \in \mathbb{C}^{m \times n}$  as

$$A = U \Sigma V^*$$

where  $U \in \mathbb{C}^{m \times m}$  and  $V \in \mathbb{C}^{n \times n}$  are unitary matrices, and  $\Sigma \in \mathbb{R}^{m \times n}$  is diagonal with non-negative diagonal values  $\sigma_i$  (singular values) ordered in decreasing order

(See A.4 in the book for the details)

- ▶ Low-rank approximations
- ▶ Useful to find the best fit subspaces of  $A$
- ▶ Projection onto subspace – perpendicular distance
- ▶ LS fit – vertical distance

## SVD (2)

$A \in \mathbb{C}^{m \times n}$  is of rank  $r < \min(m, n)$  then

$$A = \begin{bmatrix} U_1 & U_2 \end{bmatrix} \begin{bmatrix} \Sigma_1 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} V_1^* \\ V_2^* \end{bmatrix} = U_1 \Sigma_1 V_1^*$$

where  $\Sigma_1 \in \mathbb{R}^{r \times r}$ , then

- ▶  $U_1$  is an orthonormal basis in  $\mathcal{R}(A)$
- ▶  $U_2$  is an orthonormal basis in  $\mathcal{N}(A^*)$
- ▶  $V_1$  is an orthonormal basis in  $\mathcal{R}(A^*)$
- ▶  $V_2$  is an orthonormal basis in  $\mathcal{N}(A)$

Moreover,  $\mathcal{R}(A)$  and  $\mathcal{N}(A^*)$  are orthogonal to each other and they span  $\mathbb{C}^m$

( $\mathcal{N}(A^*)$  is the orthogonal complement to  $\mathcal{R}(A)$  and vice versa)



# High-order Yule-Walker (HOYW)

# High-order Yule-Walker (HOYW)

---

- ▶ Start from ARMA-model:

$$A(z)y(t) = A(z)e(t), \quad A(z) = \prod_{k=1}^n (1 - e^{i\omega_k} z^{-1}).$$

# High-order Yule-Walker (HOYW)

---

- ▶ Start from ARMA-model:

$$A(z)y(t) = A(z)e(t), \quad A(z) = \prod_{k=1}^n (1 - e^{i\omega_k} z^{-1}).$$

- ▶ Can't divide away  $A(z)$  due to poles on unit circle!



# High-order Yule-Walker (HOYW)

- ▶ Start from ARMA-model:

$$A(z)y(t) = A(z)e(t), \quad A(z) = \prod_{k=1}^n (1 - e^{i\omega_k} z^{-1}).$$

- ▶ Can't divide away  $A(z)$  due to poles on unit circle!
- ▶ Multiply both sides with some filter  $\tilde{A}(z)$  to get

$$B(z)y(t) = B(z)e(t),$$

where  $B(z) = A(z)\tilde{A}(z)$  has order  $L > n$  ("high order").

# High-order Yule-Walker (HOYW)

- ▶ Start from ARMA-model:

$$A(z)y(t) = A(z)e(t), \quad A(z) = \prod_{k=1}^n (1 - e^{i\omega_k} z^{-1}).$$

- ▶ Can't divide away  $A(z)$  due to poles on unit circle!
- ▶ Multiply both sides with some filter  $\tilde{A}(z)$  to get

$$B(z)y(t) = B(z)e(t),$$

where  $B(z) = A(z)\tilde{A}(z)$  has order  $L > n$  ("high order").

- ▶  $B(z)$  will have  $n$  roots  $\{e^{i\omega_k}\}_{k=1}^n$  on the unit circle corresponding to the frequencies, and  $L - n$  spurious roots.

# High-order Yule-Walker (HOYW)

- ▶ Start from ARMA-model:

$$A(z)y(t) = A(z)e(t), \quad A(z) = \prod_{k=1}^n (1 - e^{i\omega_k} z^{-1}).$$

- ▶ Can't divide away  $A(z)$  due to poles on unit circle!
- ▶ Multiply both sides with some filter  $\tilde{A}(z)$  to get

$$B(z)y(t) = B(z)e(t),$$

where  $B(z) = A(z)\tilde{A}(z)$  has order  $L > n$  ("high order").

- ▶  $B(z)$  will have  $n$  roots  $\{e^{i\omega_k}\}_{k=1}^n$  on the unit circle corresponding to the frequencies, and  $L - n$  spurious roots.
- ▶ **Idea:** Estimate  $B(z)$  and take the  $n$  roots  $\{r_k\}_{k=1}^n$  closest to the unit circle to get estimates  $\hat{\omega}_k = \arg(r_k)$ .

# High-order YW-equations

---

$$B(z)y(t) = B(z)e(t).$$

- From before we know that

$$r(k) + \sum_{i=1}^L b_i r(k-i) = 0, k > L.$$

# High-order YW-equations

$$B(z)y(t) = B(z)e(t).$$

- From before we know that

$$r(k) + \sum_{i=1}^L b_i r(k-i) = 0, k > L.$$

- For  $k = L+1, \dots, L+M$  we get

$$\underbrace{\begin{bmatrix} r(L) & \cdots & r(1) \\ \vdots & \ddots & \vdots \\ r(L+M-1) & \cdots & r(M) \end{bmatrix}}_{\Omega} b = - \underbrace{\begin{bmatrix} r(L+1) \\ \vdots \\ r(L+M) \end{bmatrix}}_{\rho}$$

# High-order YW-equations

$$B(z)y(t) = B(z)e(t).$$

- From before we know that

$$r(k) + \sum_{i=1}^L b_i r(k-i) = 0, k > L.$$

- For  $k = L+1, \dots, L+M$  we get

$$\underbrace{\begin{bmatrix} r(L) & \cdots & r(1) \\ \vdots & \ddots & \vdots \\ r(L+M-1) & \cdots & r(M) \end{bmatrix}}_{\Omega} b = - \underbrace{\begin{bmatrix} r(L+1) \\ \vdots \\ r(L+M) \end{bmatrix}}_{\rho}$$

- **Important property:**  $\text{rank}(\Omega) = n$ .

# High-order YW-equations

$$B(z)y(t) = B(z)e(t).$$

- ▶ From before we know that

$$r(k) + \sum_{i=1}^L b_i r(k-i) = 0, k > L.$$

- ▶ For  $k = L+1, \dots, L+M$  we get

$$\underbrace{\begin{bmatrix} r(L) & \cdots & r(1) \\ \vdots & \ddots & \vdots \\ r(L+M-1) & \cdots & r(M) \end{bmatrix}}_{\Omega} b = - \underbrace{\begin{bmatrix} r(L+1) \\ \vdots \\ r(L+M) \end{bmatrix}}_{\rho}$$

- ▶ **Important property:**  $\text{rank}(\Omega) = n$ .
- ▶ When  $L > n$  this rank-condition gives extra information about the structure of  $\Omega$ .

# HOYW-equations with estimated ACS

---

- Form  $\hat{\Omega}$  and  $\hat{\rho}$  by replacing  $r(k)$  with  $\hat{r}(k)$  in  $\Omega$  and  $\rho$ ,

$$\hat{\Omega}b \approx -\hat{\rho}. \quad (1)$$



# HOYW-equations with estimated ACS

---

- ▶ Form  $\hat{\Omega}$  and  $\hat{\rho}$  by replacing  $r(k)$  with  $\hat{r}(k)$  in  $\Omega$  and  $\rho$ ,

$$\hat{\Omega}b \approx -\hat{\rho}. \quad (1)$$

- ▶ Random errors in  $\hat{r}(k)$ , so we (almost surely) get a full-rank  $\hat{\Omega}$ ,

$$\text{rank}(\hat{\Omega}) = \min(M, L) > n.$$

# HOYW-equations with estimated ACS

- ▶ Form  $\hat{\Omega}$  and  $\hat{\rho}$  by replacing  $r(k)$  with  $\hat{r}(k)$  in  $\Omega$  and  $\rho$ ,

$$\hat{\Omega}b \approx -\hat{\rho}. \quad (1)$$

- ▶ Random errors in  $\hat{r}(k)$ , so we (almost surely) get a full-rank  $\hat{\Omega}$ ,

$$\text{rank}(\hat{\Omega}) = \min(M, L) > n.$$

- ▶ **Ill-conditioned:** For reasonable large  $N$ ,  $\hat{\Omega}$  will be close to a rank- $n$  matrix. Hence, solving for  $b$  in (1) directly is numerically problematic.

# HOYW-equations with estimated ACS

- ▶ Form  $\hat{\Omega}$  and  $\hat{\rho}$  by replacing  $r(k)$  with  $\hat{r}(k)$  in  $\Omega$  and  $\rho$ ,

$$\hat{\Omega}b \approx -\hat{\rho}. \quad (1)$$

- ▶ Random errors in  $\hat{r}(k)$ , so we (almost surely) get a full-rank  $\hat{\Omega}$ ,

$$\text{rank}(\hat{\Omega}) = \min(M, L) > n.$$

- ▶ **Ill-conditioned:** For reasonable large  $N$ ,  $\hat{\Omega}$  will be close to a rank- $n$  matrix. Hence, solving for  $b$  in (1) directly is numerically problematic.
- ▶ **Idea:** First approximate  $\hat{\Omega}$  with a rank- $n$  matrix  $\hat{\Omega}_n$ .

# HOYW-equations with estimated ACS

- ▶ Form  $\hat{\Omega}$  and  $\hat{\rho}$  by replacing  $r(k)$  with  $\hat{r}(k)$  in  $\Omega$  and  $\rho$ ,

$$\hat{\Omega}b \approx -\hat{\rho}. \quad (1)$$

- ▶ Random errors in  $\hat{r}(k)$ , so we (almost surely) get a full-rank  $\hat{\Omega}$ ,

$$\text{rank}(\hat{\Omega}) = \min(M, L) > n.$$

- ▶ **Ill-conditioned:** For reasonable large  $N$ ,  $\hat{\Omega}$  will be close to a rank- $n$  matrix. Hence, solving for  $b$  in (1) directly is numerically problematic.
- ▶ **Idea:** First approximate  $\hat{\Omega}$  with a rank- $n$  matrix  $\hat{\Omega}_n$ .
- ▶  $\hat{\Omega}_n$  should be closer to the true  $\Omega$  since we exploited

$$\text{rank}(\Omega) = n.$$

# HOYW-equations with estimated ACS

- ▶ Form  $\hat{\Omega}$  and  $\hat{\rho}$  by replacing  $r(k)$  with  $\hat{r}(k)$  in  $\Omega$  and  $\rho$ ,

$$\hat{\Omega}b \approx -\hat{\rho}. \quad (1)$$

- ▶ Random errors in  $\hat{r}(k)$ , so we (almost surely) get a full-rank  $\hat{\Omega}$ ,

$$\text{rank}(\hat{\Omega}) = \min(M, L) > n.$$

- ▶ **Ill-conditioned:** For reasonable large  $N$ ,  $\hat{\Omega}$  will be close to a rank- $n$  matrix. Hence, solving for  $b$  in (1) directly is numerically problematic.
- ▶ **Idea:** First approximate  $\hat{\Omega}$  with a rank- $n$  matrix  $\hat{\Omega}_n$ .
- ▶  $\hat{\Omega}_n$  should be closer to the true  $\Omega$  since we exploited

$$\text{rank}(\Omega) = n.$$

- ▶ Find  $\hat{b}$  by solving (LS)

$$\hat{\Omega}_n \hat{b} \approx -\hat{\rho} \implies \hat{b} = -\hat{\Omega}_n^\dagger \hat{\rho}.$$

# HOYW in practice

---

- Use SVD on  $\hat{\Omega}$ ,

$$\hat{\Omega} = U\Sigma V^* = \begin{bmatrix} U_1 & U_2 \end{bmatrix} \begin{bmatrix} \Sigma_1 & 0 \\ 0 & \Sigma_2 \end{bmatrix} \begin{bmatrix} V_1^* \\ V_2^* \end{bmatrix},$$

where  $\Sigma_1$  contains the  $n$  largest singular values.

# HOYW in practice

---

- ▶ Use SVD on  $\hat{\Omega}$ ,

$$\hat{\Omega} = U\Sigma V^* = \begin{bmatrix} U_1 & U_2 \end{bmatrix} \begin{bmatrix} \Sigma_1 & 0 \\ 0 & \Sigma_2 \end{bmatrix} \begin{bmatrix} V_1^* \\ V_2^* \end{bmatrix},$$

where  $\Sigma_1$  contains the  $n$  largest singular values.

- ▶ Truncate (i.e. replace  $\Sigma_2$  with 0) to get rank- $n$  matrix

$$\hat{\Omega}_n = U_1 \Sigma_1 V_1^* \approx \hat{\Omega}.$$

# HOYW in practice

- ▶ Use SVD on  $\hat{\Omega}$ ,

$$\hat{\Omega} = U\Sigma V^* = \begin{bmatrix} U_1 & U_2 \end{bmatrix} \begin{bmatrix} \Sigma_1 & 0 \\ 0 & \Sigma_2 \end{bmatrix} \begin{bmatrix} V_1^* \\ V_2^* \end{bmatrix},$$

where  $\Sigma_1$  contains the  $n$  largest singular values.

- ▶ Truncate (i.e. replace  $\Sigma_2$  with 0) to get rank- $n$  matrix

$$\hat{\Omega}_n = U_1 \Sigma_1 V_1^* \approx \hat{\Omega}.$$

- ▶ Solve for  $\hat{b}$ ,

$$\hat{b} = -\hat{\Omega}_n^\dagger \hat{\rho} =$$



# HOYW in practice

- ▶ Use SVD on  $\hat{\Omega}$ ,

$$\hat{\Omega} = U\Sigma V^* = \begin{bmatrix} U_1 & U_2 \end{bmatrix} \begin{bmatrix} \Sigma_1 & 0 \\ 0 & \Sigma_2 \end{bmatrix} \begin{bmatrix} V_1^* \\ V_2^* \end{bmatrix},$$

where  $\Sigma_1$  contains the  $n$  largest singular values.

- ▶ Truncate (i.e. replace  $\Sigma_2$  with 0) to get rank- $n$  matrix

$$\hat{\Omega}_n = U_1 \Sigma_1 V_1^* \approx \hat{\Omega}.$$

- ▶ Solve for  $\hat{b}$ ,

$$\hat{b} = -\hat{\Omega}_n^\dagger \hat{\rho} = -V_1 \Sigma_1^{-1} U_1^* \hat{\rho}.$$

# HOYW in practice

- ▶ Use SVD on  $\hat{\Omega}$ ,

$$\hat{\Omega} = U\Sigma V^* = \begin{bmatrix} U_1 & U_2 \end{bmatrix} \begin{bmatrix} \Sigma_1 & 0 \\ 0 & \Sigma_2 \end{bmatrix} \begin{bmatrix} V_1^* \\ V_2^* \end{bmatrix},$$

where  $\Sigma_1$  contains the  $n$  largest singular values.

- ▶ Truncate (i.e. replace  $\Sigma_2$  with 0) to get rank- $n$  matrix

$$\hat{\Omega}_n = U_1 \Sigma_1 V_1^* \approx \hat{\Omega}.$$

- ▶ Solve for  $\hat{b}$ ,

$$\hat{b} = -\hat{\Omega}_n^\dagger \hat{\rho} = -V_1 \Sigma_1^{-1} U_1^* \hat{\rho}.$$

- ▶  $\{\hat{\omega}_k\}_{k=1}^n$  from the  $n$  roots of  $\hat{B}(z)$  closest to the unit circle.

# HOYW in practice

- ▶ Use SVD on  $\hat{\Omega}$ ,

$$\hat{\Omega} = U\Sigma V^* = \begin{bmatrix} U_1 & U_2 \end{bmatrix} \begin{bmatrix} \Sigma_1 & 0 \\ 0 & \Sigma_2 \end{bmatrix} \begin{bmatrix} V_1^* \\ V_2^* \end{bmatrix},$$

where  $\Sigma_1$  contains the  $n$  largest singular values.

- ▶ Truncate (i.e. replace  $\Sigma_2$  with 0) to get rank- $n$  matrix

$$\hat{\Omega}_n = U_1 \Sigma_1 V_1^* \approx \hat{\Omega}.$$

- ▶ Solve for  $\hat{b}$ ,

$$\hat{b} = -\hat{\Omega}_n^\dagger \hat{\rho} = -V_1 \Sigma_1^{-1} U_1^* \hat{\rho}.$$

- ▶  $\{\hat{\omega}_k\}_{k=1}^n$  from the  $n$  roots of  $\hat{B}(z)$  closest to the unit circle.
- ▶ **User parameters:**  $M \approx L$ , and  $L + M$  a fraction of  $N$ .

# HOYW in practice

- ▶ Use SVD on  $\hat{\Omega}$ ,

$$\hat{\Omega} = U\Sigma V^* = \begin{bmatrix} U_1 & U_2 \end{bmatrix} \begin{bmatrix} \Sigma_1 & 0 \\ 0 & \Sigma_2 \end{bmatrix} \begin{bmatrix} V_1^* \\ V_2^* \end{bmatrix},$$

where  $\Sigma_1$  contains the  $n$  largest singular values.

- ▶ Truncate (i.e. replace  $\Sigma_2$  with 0) to get rank- $n$  matrix

$$\hat{\Omega}_n = U_1 \Sigma_1 V_1^* \approx \hat{\Omega}.$$

- ▶ Solve for  $\hat{b}$ ,

$$\hat{b} = -\hat{\Omega}_n^\dagger \hat{\rho} = -V_1 \Sigma_1^{-1} U_1^* \hat{\rho}.$$

- ▶  $\{\hat{\omega}_k\}_{k=1}^n$  from the  $n$  roots of  $\hat{B}(z)$  closest to the unit circle.
- ▶ **User parameters:**  $M \approx L$ , and  $L + M$  a fraction of  $N$ .
- ▶ **Note:** SVD can also give an idea about the size of  $n$ .



# MUltiple Signal Classification (MUSIC)

# The Covariance Matrix Equation

Let

$$a(\omega) = \begin{bmatrix} 1 & e^{-i\omega} & \cdots & e^{-i(m-1)\omega} \end{bmatrix}^\top \text{ for some } m > n$$

$$A = \begin{bmatrix} a(\omega_1) & \cdots & a(\omega_n) \end{bmatrix} \in \mathbb{C}^{m \times n}$$

Let  $\tilde{y}(t) = \begin{bmatrix} y(t) & \cdots & y(t-m+1) \end{bmatrix}^\top$ , then

$$\tilde{y}(t) = A\tilde{x}(t) + \tilde{e}(t)$$

$$\tilde{x}(t) = \begin{bmatrix} x_1(t) & \cdots & x_n(t) \end{bmatrix}^\top, \quad \tilde{e}(t) = \begin{bmatrix} e(t) & \cdots & e(t-m+1) \end{bmatrix}^\top$$

**Covariance matrix:**  $R = E\{\tilde{y}(t)\tilde{y}^*(t)\} = APA^* + \sigma^2 I$ , where

$$P = E\{\tilde{x}(t)\tilde{x}^*(t)\} = \begin{bmatrix} \alpha_1^2 & & 0 \\ & \ddots & \\ 0 & & \alpha_n^2 \end{bmatrix}$$

# Some rank conditions

---

- ▶  $\text{rank}(A) = n$  (if  $m \geq n$  and  $\omega_k \neq \omega_p$  for  $k \neq p$ ).

# Some rank conditions

---

- ▶  $\text{rank}(A) = n$  (if  $m \geq n$  and  $\omega_k \neq \omega_p$  for  $k \neq p$ ).
- ▶ Hence,

$$\text{rank}(APA^*) = n.$$



# Some rank conditions

---

►  $\text{rank}(A) = n$  (if  $m \geq n$  and  $\omega_k \neq \omega_p$  for  $k \neq p$ ).

► Hence,

$$\text{rank}(APA^*) = n.$$

►  $APA^*$  has  $n$  strictly positive eigenvalues, and  $m - n$  eigenvalues equal to 0,

# Some rank conditions

---

►  $\text{rank}(A) = n$  (if  $m \geq n$  and  $\omega_k \neq \omega_p$  for  $k \neq p$ ).

► Hence,

$$\text{rank}(APA^*) = n.$$

►  $APA^*$  has  $n$  strictly positive eigenvalues, and  $m - n$  eigenvalues equal to 0,

$$\tilde{\lambda}_1 \geq \tilde{\lambda}_2 \geq \cdots \geq \tilde{\lambda}_n > 0, \quad \tilde{\lambda}_{n+1} = \cdots = \tilde{\lambda}_m = 0.$$

# Some rank conditions

►  $\text{rank}(A) = n$  (if  $m \geq n$  and  $\omega_k \neq \omega_p$  for  $k \neq p$ ).

► Hence,

$$\text{rank}(APA^*) = n.$$

►  $APA^*$  has  $n$  strictly positive eigenvalues, and  $m - n$  eigenvalues equal to 0,

$$\tilde{\lambda}_1 \geq \tilde{\lambda}_2 \geq \cdots \geq \tilde{\lambda}_n > 0, \quad \tilde{\lambda}_{n+1} = \cdots = \tilde{\lambda}_m = 0.$$

► The eigenvalues  $\lambda_k$  of  $R = APA^* + \sigma^2 I$  are thus given by

$$\lambda_k =$$

# Some rank conditions

►  $\text{rank}(A) = n$  (if  $m \geq n$  and  $\omega_k \neq \omega_p$  for  $k \neq p$ ).

► Hence,

$$\text{rank}(APA^*) = n.$$

►  $APA^*$  has  $n$  strictly positive eigenvalues, and  $m - n$  eigenvalues equal to 0,

$$\tilde{\lambda}_1 \geq \tilde{\lambda}_2 \geq \cdots \geq \tilde{\lambda}_n > 0, \quad \tilde{\lambda}_{n+1} = \cdots = \tilde{\lambda}_m = 0.$$

► The eigenvalues  $\lambda_k$  of  $R = APA^* + \sigma^2 I$  are thus given by

$$\lambda_k = \tilde{\lambda}_k + \sigma^2 > 0, \quad k = 1, \dots, m,$$

# Some rank conditions

- ▶  $\text{rank}(A) = n$  (if  $m \geq n$  and  $\omega_k \neq \omega_p$  for  $k \neq p$ ).

- ▶ Hence,

$$\text{rank}(APA^*) = n.$$

- ▶  $APA^*$  has  $n$  strictly positive eigenvalues, and  $m - n$  eigenvalues equal to 0,

$$\tilde{\lambda}_1 \geq \tilde{\lambda}_2 \geq \cdots \geq \tilde{\lambda}_n > 0, \quad \tilde{\lambda}_{n+1} = \cdots = \tilde{\lambda}_m = 0.$$

- ▶ The eigenvalues  $\lambda_k$  of  $R = APA^* + \sigma^2 I$  are thus given by

$$\lambda_k = \tilde{\lambda}_k + \sigma^2 > 0, \quad k = 1, \dots, m,$$

- ▶ and thus

$$\text{rank}(R) = m.$$

# Eigendecomposition of $R$

---

$$R = APA^* + \sigma^2 I \quad (m > n)$$

► **Eigenvalues:**  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n > \lambda_{n+1} = \dots = \lambda_m = \sigma^2$

# Eigendecomposition of $R$

---

$$R = APA^* + \sigma^2 I \quad (m > n)$$

- ▶ **Eigenvalues:**  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n > \lambda_{n+1} = \dots = \lambda_m = \sigma^2$
- ▶ **Eigenvectors:**  $s_1, \dots, s_n, g_1, \dots, g_{m-n}$  orthonormal

$$S = \begin{bmatrix} s_1 & \cdots & s_n \end{bmatrix} \text{ corresponding to } \lambda_1, \dots, \lambda_n$$

$$G = \begin{bmatrix} g_1 & \cdots & g_{m-n} \end{bmatrix} \text{ corresponding to } \lambda_{n+1}, \dots, \lambda_m$$

# Eigendecomposition of $R$

$$R = APA^* + \sigma^2 I \quad (m > n)$$

- **Eigenvalues:**  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n > \lambda_{n+1} = \dots = \lambda_m = \sigma^2$
- **Eigenvectors:**  $s_1, \dots, s_n, g_1, \dots, g_{m-n}$  orthonormal

$$S = \begin{bmatrix} s_1 & \cdots & s_n \end{bmatrix} \text{ corresponding to } \lambda_1, \dots, \lambda_n$$

$$G = \begin{bmatrix} g_1 & \cdots & g_{m-n} \end{bmatrix} \text{ corresponding to } \lambda_{n+1}, \dots, \lambda_m$$

- **Eigendecomposition:**

$$R = \begin{bmatrix} S & G \end{bmatrix} \begin{bmatrix} \lambda_1 & & 0 \\ & \ddots & \\ 0 & & \lambda_m \end{bmatrix} \begin{bmatrix} S^* \\ G^* \end{bmatrix}.$$



# MUSIC, motivation

---

- ▶ Note that

$$RG =$$

# MUSIC, motivation

---

- ▶ Note that

$$RG = \sigma^2 G =$$

# MUSIC, motivation

---

- ▶ Note that

$$RG = \sigma^2 G = APA^*G + \sigma^2 G$$

# MUSIC, motivation

---

- ▶ Note that

$$RG = \sigma^2 G = APA^*G + \sigma^2 G$$

- ▶ Follows that  $APA^*G = 0$ .

# MUSIC, motivation

---

- ▶ Note that

$$RG = \sigma^2 G = APA^*G + \sigma^2 G$$

- ▶ Follows that  $APA^*G = 0$ . Since  $AP$  has full rank we get

$$0 = A^*G =$$

# MUSIC, motivation

---

- Note that

$$RG = \sigma^2 G = APA^*G + \sigma^2 G$$

- Follows that  $APA^*G = 0$ . Since  $AP$  has full rank we get

$$0 = A^*G = \begin{bmatrix} a^*(\omega_1)G \\ \vdots \\ a^*(\omega_n)G \end{bmatrix}$$

# MUSIC, motivation

- Note that

$$RG = \sigma^2 G = APA^*G + \sigma^2 G$$

- Follows that  $APA^*G = 0$ . Since  $AP$  has full rank we get

$$0 = A^*G = \begin{bmatrix} a^*(\omega_1)G \\ \vdots \\ a^*(\omega_n)G \end{bmatrix} \implies a^*(\omega)G = 0 \text{ if } \omega = \omega_k.$$

# MUSIC, motivation

- Note that

$$RG = \sigma^2 G = APA^*G + \sigma^2 G$$

- Follows that  $APA^*G = 0$ . Since  $AP$  has full rank we get

$$0 = A^*G = \begin{bmatrix} a^*(\omega_1)G \\ \vdots \\ a^*(\omega_n)G \end{bmatrix} \implies a^*(\omega)G = 0 \text{ if } \omega = \omega_k.$$

- **Result:** For  $\omega = \omega_k$  (the true frequencies)

$$a^*(\omega)GG^*a(\omega) = 0$$



# MUSIC, motivation

- ▶ Note that

$$RG = \sigma^2 G = APA^*G + \sigma^2 G$$

- ▶ Follows that  $APA^*G = 0$ . Since  $AP$  has full rank we get

$$0 = A^*G = \begin{bmatrix} a^*(\omega_1)G \\ \vdots \\ a^*(\omega_n)G \end{bmatrix} \implies a^*(\omega)G = 0 \text{ if } \omega = \omega_k.$$

- ▶ **Result:** For  $\omega = \omega_k$  (the true frequencies)

$$a^*(\omega)GG^*a(\omega) = 0$$

- ▶ It can be shown that the above holds **if and only if**  $\omega = \omega_k$  for  $k = 1, \dots, n$ .

# MUSIC, motivation

- Note that

$$RG = \sigma^2 G = APA^*G + \sigma^2 G$$

- Follows that  $APA^*G = 0$ . Since  $AP$  has full rank we get

$$0 = A^*G = \begin{bmatrix} a^*(\omega_1)G \\ \vdots \\ a^*(\omega_n)G \end{bmatrix} \implies a^*(\omega)G = 0 \text{ if } \omega = \omega_k.$$

- **Result:** For  $\omega = \omega_k$  (the true frequencies)

$$a^*(\omega)GG^*a(\omega) = 0$$

- It can be shown that the above holds **if and only if**  $\omega = \omega_k$  for  $k = 1, \dots, n$ .
- (Any  $a(\tilde{\omega})$  that satisfy the equation belongs to  $\mathcal{N}(G^*)$ , which is a subspace of dimension  $n$ . At most  $n$  vectors on the form  $a(\omega)$  can belong to this subspace, which by the above must be  $\{a(\omega_k)\}_{k=1}^n$ .)

# MUSIC

---

Compute sample covariance matrix (hermitian positive definite)

$$\hat{R} = \frac{1}{N} \sum_{t=m}^N \tilde{y}(t) \tilde{y}^*(t)$$

and its eigendecomposition (here SVD) giving matrices  $\hat{S}$  and  $\hat{G}$

# MUSIC

---

Compute sample covariance matrix (hermitian positive definite)

$$\hat{R} = \frac{1}{N} \sum_{t=m}^N \tilde{y}(t) \tilde{y}^*(t)$$

and its eigendecomposition (here SVD) giving matrices  $\hat{S}$  and  $\hat{G}$

## Spectral MUSIC

Plot pseudo spectrum and determine frequencies from the  $n$  highest peaks

$$\frac{1}{a^*(\omega) \hat{G} \hat{G}^* a(\omega)} \quad \omega \in [-\pi, \pi]$$

# MUSIC

Compute sample covariance matrix (hermitian positive definite)

$$\hat{R} = \frac{1}{N} \sum_{t=m}^N \tilde{y}(t) \tilde{y}^*(t)$$

and its eigendecomposition (here SVD) giving matrices  $\hat{S}$  and  $\hat{G}$

## Spectral MUSIC

Plot pseudo spectrum and determine frequencies from the  $n$  highest peaks

$$\frac{1}{a^*(\omega) \hat{G} \hat{G}^* a(\omega)} \quad \omega \in [-\pi, \pi]$$

## Root MUSIC

Determine frequencies from the angular positions of the  $n$  roots of

$$a^T(z^{-1}) \hat{G} \hat{G}^* a(z) = 0$$

closest to the unit circle



# Min-Norm method

# The Min-Norm method

---

- ▶ **Goal:** Reduce computational burden, and reduce risk of false frequency estimates.

# The Min-Norm method

---

- ▶ **Goal:** Reduce computational burden, and reduce risk of false frequency estimates.
- ▶ **Idea:** From our previous discussion we can see that for any vector  $g \in \mathcal{R}(G) = \mathcal{N}(A^*)$  we have

$$a^*(\omega)g = 0, \quad \text{for } \omega = \omega_k.$$



# The Min-Norm method

---

- ▶ **Goal:** Reduce computational burden, and reduce risk of false frequency estimates.
- ▶ **Idea:** From our previous discussion we can see that for any vector  $g \in \mathcal{R}(G) = \mathcal{N}(A^*)$  we have

$$a^*(\omega)g = 0, \quad \text{for } \omega = \omega_k.$$

- ▶ Computationally easier to solve this equation!

# The Min-Norm method

---

- ▶ **Goal:** Reduce computational burden, and reduce risk of false frequency estimates.
- ▶ **Idea:** From our previous discussion we can see that for any vector  $g \in \mathcal{R}(G) = \mathcal{N}(A^*)$  we have

$$a^*(\omega)g = 0, \quad \text{for } \omega = \omega_k.$$

- ▶ Computationally easier to solve this equation!
- ▶ **Which vector to use?** Are strong empirical evidence that the vector

$$g = \begin{bmatrix} 1 \\ \tilde{g} \end{bmatrix} \in \mathcal{R}(G)$$

with minimum Euclidean norm have good properties (accuracy, and reduced risk of spurious  $\omega$ -estimates).

# Min-Norm

---

Find

$$\begin{bmatrix} 1 \\ \hat{g} \end{bmatrix} \in \mathcal{R}(\hat{G})$$

with minimum Euclidean norm (see textbook).

# Min-Norm

---

Find

$$\begin{bmatrix} 1 \\ \hat{g} \end{bmatrix} \in \mathcal{R}(\hat{G})$$

with minimum Euclidean norm (see textbook).

## Spectral Min-Norm

Plot pseudo spectrum and determine frequencies from the  $n$  highest peaks

$$\frac{1}{a^*(\omega) \begin{bmatrix} 1 \\ \hat{g} \end{bmatrix}}$$

# Min-Norm

---

Find

$$\begin{bmatrix} 1 \\ \hat{g} \end{bmatrix} \in \mathcal{R}(\hat{G})$$

with minimum Euclidean norm (see textbook).

## Spectral Min-Norm

Plot pseudo spectrum and determine frequencies from the  $n$  highest peaks

$$\frac{1}{a^*(\omega) \begin{bmatrix} 1 \\ \hat{g} \end{bmatrix}}$$

## Root Min-Norm

Determine frequencies from the angular positions of the  $n$  roots of

$$a^T(z^{-1}) \begin{bmatrix} 1 \\ \hat{g} \end{bmatrix} = 0$$

closest to the unit circle



# Estimation of Signal Parameters by Rotational Invariance Techniques (ESPRIT)

**esprit (noun) – vivacious cleverness or wit**

# Theory of ESPRIT

---

- ▶ Let  $A_1$  be  $A$  with last row removed, and  $A_2$  be  $A$  with first row removed

# Theory of ESPRIT

---

- Let  $A_1$  be  $A$  with last row removed, and  $A_2$  be  $A$  with first row removed, i.e.,

$$A_1 = \begin{bmatrix} I_{m-1} & 0 \end{bmatrix} A, \quad A_2 = \begin{bmatrix} 0 & I_{m-1} \end{bmatrix} A, \quad (m-1 \times n)$$



# Theory of ESPRIT

- ▶ Let  $A_1$  be  $A$  with last row removed, and  $A_2$  be  $A$  with first row removed, i.e.,

$$A_1 = \begin{bmatrix} I_{m-1} & 0 \end{bmatrix} A, \quad A_2 = \begin{bmatrix} 0 & I_{m-1} \end{bmatrix} A, \quad (m-1 \times n)$$

- ▶ then,

$$A_2 = A_1 D, \quad D = \begin{bmatrix} e^{-i\omega_1} & & 0 \\ & \ddots & \\ 0 & & e^{-i\omega_n} \end{bmatrix}.$$

# Theory of ESPRIT

- ▶ Let  $A_1$  be  $A$  with last row removed, and  $A_2$  be  $A$  with first row removed, i.e.,

$$A_1 = \begin{bmatrix} I_{m-1} & 0 \end{bmatrix} A, \quad A_2 = \begin{bmatrix} 0 & I_{m-1} \end{bmatrix} A, \quad (m-1 \times n)$$

- ▶ then,

$$A_2 = A_1 D, \quad D = \begin{bmatrix} e^{-i\omega_1} & & 0 \\ & \ddots & \\ 0 & & e^{-i\omega_n} \end{bmatrix}.$$

- ▶ In the same way let

$$S_1 = \begin{bmatrix} I_{m-1} & 0 \end{bmatrix} S, \quad S_2 = \begin{bmatrix} 0 & I_{m-1} \end{bmatrix} S.$$

- ▶ There exist nonsingular  $C$  such that  $S = AC$ .

# Theory of ESPRIT

- ▶ Let  $A_1$  be  $A$  with last row removed, and  $A_2$  be  $A$  with first row removed, i.e.,

$$A_1 = \begin{bmatrix} I_{m-1} & 0 \end{bmatrix} A, \quad A_2 = \begin{bmatrix} 0 & I_{m-1} \end{bmatrix} A, \quad (m-1 \times n)$$

- ▶ then,

$$A_2 = A_1 D, \quad D = \begin{bmatrix} e^{-i\omega_1} & & 0 \\ & \ddots & \\ 0 & & e^{-i\omega_n} \end{bmatrix}.$$

- ▶ In the same way let

$$S_1 = \begin{bmatrix} I_{m-1} & 0 \end{bmatrix} S, \quad S_2 = \begin{bmatrix} 0 & I_{m-1} \end{bmatrix} S.$$

- ▶ There exist nonsingular  $C$  such that  $S = AC$ . Hence,

$$S_2 =$$

# Theory of ESPRIT

- ▶ Let  $A_1$  be  $A$  with last row removed, and  $A_2$  be  $A$  with first row removed, i.e.,

$$A_1 = \begin{bmatrix} I_{m-1} & 0 \end{bmatrix} A, \quad A_2 = \begin{bmatrix} 0 & I_{m-1} \end{bmatrix} A, \quad (m-1 \times n)$$

- ▶ then,

$$A_2 = A_1 D, \quad D = \begin{bmatrix} e^{-i\omega_1} & & 0 \\ & \ddots & \\ 0 & & e^{-i\omega_n} \end{bmatrix}.$$

- ▶ In the same way let

$$S_1 = \begin{bmatrix} I_{m-1} & 0 \end{bmatrix} S, \quad S_2 = \begin{bmatrix} 0 & I_{m-1} \end{bmatrix} S.$$

- ▶ There exist nonsingular  $C$  such that  $S = AC$ . Hence,

$$S_2 = A_2 C =$$

# Theory of ESPRIT

- ▶ Let  $A_1$  be  $A$  with last row removed, and  $A_2$  be  $A$  with first row removed, i.e.,

$$A_1 = \begin{bmatrix} I_{m-1} & 0 \end{bmatrix} A, \quad A_2 = \begin{bmatrix} 0 & I_{m-1} \end{bmatrix} A, \quad (m-1 \times n)$$

- ▶ then,

$$A_2 = A_1 D, \quad D = \begin{bmatrix} e^{-i\omega_1} & & 0 \\ & \ddots & \\ 0 & & e^{-i\omega_n} \end{bmatrix}.$$

- ▶ In the same way let

$$S_1 = \begin{bmatrix} I_{m-1} & 0 \end{bmatrix} S, \quad S_2 = \begin{bmatrix} 0 & I_{m-1} \end{bmatrix} S.$$

- ▶ There exist nonsingular  $C$  such that  $S = AC$ . Hence,

$$S_2 = A_2 C = A_1 D C =$$

# Theory of ESPRIT

- ▶ Let  $A_1$  be  $A$  with last row removed, and  $A_2$  be  $A$  with first row removed, i.e.,

$$A_1 = \begin{bmatrix} I_{m-1} & 0 \end{bmatrix} A, \quad A_2 = \begin{bmatrix} 0 & I_{m-1} \end{bmatrix} A, \quad (m-1 \times n)$$

- ▶ then,

$$A_2 = A_1 D, \quad D = \begin{bmatrix} e^{-i\omega_1} & & 0 \\ & \ddots & \\ 0 & & e^{-i\omega_n} \end{bmatrix}.$$

- ▶ In the same way let

$$S_1 = \begin{bmatrix} I_{m-1} & 0 \end{bmatrix} S, \quad S_2 = \begin{bmatrix} 0 & I_{m-1} \end{bmatrix} S.$$

- ▶ There exist nonsingular  $C$  such that  $S = AC$ . Hence,

$$S_2 = A_2 C = A_1 D C = S_1 \underbrace{C^{-1} D C}_{\phi}.$$

# ESPIRIT

---

► We have

$$S_2 = S_1\phi, \quad \text{where } \phi = C^{-1}DC.$$

# ESPIRIT

---

- ▶ We have

$$S_2 = S_1\phi, \quad \text{where } \phi = C^{-1}DC.$$

- ▶ Eigenvalues of  $\phi$ :  $\phi$  is similar to  $D \Rightarrow$



# ESPIRIT

---

- ▶ We have

$$S_2 = S_1\phi, \quad \text{where } \phi = C^{-1}DC.$$

- ▶ **Eigenvalues of  $\phi$ :**  $\phi$  is similar to  $D \Rightarrow$  The eigenvalues of  $\phi$  are the same as the eigenvalues of  $D$ , i.e.,  $\{e^{-i\omega_k}\}_{k=1}^n$ .

# ESPIRIT

---

- ▶ We have

$$S_2 = S_1\phi, \quad \text{where } \phi = C^{-1}DC.$$

- ▶ **Eigenvalues of  $\phi$ :**  $\phi$  is similar to  $D \Rightarrow$  The eigenvalues of  $\phi$  are the same as the eigenvalues of  $D$ , i.e.,  $\{e^{-i\omega_k}\}_{k=1}^n$ .
- ▶  $S_1$  and  $S_2$  have full rank, so  $\phi$  is uniquely given by

$$\phi = (S_1^* S_1)^{-1} S_1^* S_2.$$

# ESPIRIT

---

- ▶ We have

$$S_2 = S_1\phi, \quad \text{where } \phi = C^{-1}DC.$$

- ▶ **Eigenvalues of  $\phi$ :**  $\phi$  is similar to  $D \Rightarrow$  The eigenvalues of  $\phi$  are the same as the eigenvalues of  $D$ , i.e.,  $\{e^{-i\omega_k}\}_{k=1}^n$ .
- ▶  $S_1$  and  $S_2$  have full rank, so  $\phi$  is uniquely given by

$$\phi = (S_1^* S_1)^{-1} S_1^* S_2.$$

## ESPIRIT:

- ▶ Compute estimated  $\hat{R}$  for some  $m > n$ .

# ESPIRIT

---

- ▶ We have

$$S_2 = S_1\phi, \quad \text{where } \phi = C^{-1}DC.$$

- ▶ **Eigenvalues of  $\phi$ :**  $\phi$  is similar to  $D \Rightarrow$  The eigenvalues of  $\phi$  are the same as the eigenvalues of  $D$ , i.e.,  $\{e^{-i\omega_k}\}_{k=1}^n$ .
- ▶  $S_1$  and  $S_2$  have full rank, so  $\phi$  is uniquely given by

$$\phi = (S_1^* S_1)^{-1} S_1^* S_2.$$

## ESPIRIT:

- ▶ Compute estimated  $\hat{R}$  for some  $m > n$ .
- ▶ Use SVD to get  $\hat{S}$  and construct  $\hat{S}_1$  and  $\hat{S}_2$ .

# ESPIRIT

- ▶ We have

$$S_2 = S_1 \phi, \quad \text{where } \phi = C^{-1}DC.$$

- ▶ **Eigenvalues of  $\phi$ :**  $\phi$  is similar to  $D \Rightarrow$  The eigenvalues of  $\phi$  are the same as the eigenvalues of  $D$ , i.e.,  $\{e^{-i\omega_k}\}_{k=1}^n$ .
- ▶  $S_1$  and  $S_2$  have full rank, so  $\phi$  is uniquely given by

$$\phi = (S_1^* S_1)^{-1} S_1^* S_2.$$

## ESPIRIT:

- ▶ Compute estimated  $\hat{R}$  for some  $m > n$ .
- ▶ Use SVD to get  $\hat{S}$  and construct  $\hat{S}_1$  and  $\hat{S}_2$ .
- ▶ Let  $\hat{\phi} = (\hat{S}_1^* \hat{S}_1)^{-1} \hat{S}_1^* \hat{S}_2$  (LS-solution).

# ESPIRIT

- ▶ We have

$$S_2 = S_1 \phi, \quad \text{where } \phi = C^{-1}DC.$$

- ▶ **Eigenvalues of  $\phi$ :**  $\phi$  is similar to  $D \Rightarrow$  The eigenvalues of  $\phi$  are the same as the eigenvalues of  $D$ , i.e.,  $\{e^{-i\omega_k}\}_{k=1}^n$ .
- ▶  $S_1$  and  $S_2$  have full rank, so  $\phi$  is uniquely given by

$$\phi = (S_1^* S_1)^{-1} S_1^* S_2.$$

## ESPIRIT:

- ▶ Compute estimated  $\hat{R}$  for some  $m > n$ .
- ▶ Use SVD to get  $\hat{S}$  and construct  $\hat{S}_1$  and  $\hat{S}_2$ .
- ▶ Let  $\hat{\phi} = (\hat{S}_1^* \hat{S}_1)^{-1} \hat{S}_1^* \hat{S}_2$  (LS-solution).
- ▶ Let  $\hat{v}_k$  be the eigenvalues of  $\hat{\phi}$ . Then the estimated frequencies are

$$\hat{\omega}_k = -\arg(\hat{v}_k).$$

# ESPIRIT

- ▶ We have

$$S_2 = S_1 \phi, \quad \text{where } \phi = C^{-1}DC.$$

- ▶ **Eigenvalues of  $\phi$ :**  $\phi$  is similar to  $D \Rightarrow$  The eigenvalues of  $\phi$  are the same as the eigenvalues of  $D$ , i.e.,  $\{e^{-i\omega_k}\}_{k=1}^n$ .
- ▶  $S_1$  and  $S_2$  have full rank, so  $\phi$  is uniquely given by

$$\phi = (S_1^* S_1)^{-1} S_1^* S_2.$$

## ESPIRIT:

- ▶ Compute estimated  $\hat{R}$  for some  $m > n$ .
- ▶ Use SVD to get  $\hat{S}$  and construct  $\hat{S}_1$  and  $\hat{S}_2$ .
- ▶ Let  $\hat{\phi} = (\hat{S}_1^* \hat{S}_1)^{-1} \hat{S}_1^* \hat{S}_2$  (LS-solution).
- ▶ Let  $\hat{v}_k$  be the eigenvalues of  $\hat{\phi}$ . Then the estimated frequencies are

$$\hat{\omega}_k = -\arg(\hat{v}_k).$$

- ▶ If estimates of  $\alpha_k$  are needed, use LS with the estimated frequencies.

# Performance

- ▶ All methods can achieve super resolution (beyond the Periodogram)
- ▶ Finding the frequencies through subspace methods is tractable as opposed to NLS (typically)
- ▶ ESPRIT is typically slightly better than the others
- ▶ We still have to choose the order  $n$

Method	Computational Burden	Accuracy / Resolution	Risk for False Freq Estimates
Periodogram	small	medium-high	medium
Nonlinear LS	very high	very high	very high
Yule-Walker	medium	high	medium
Pisarenko	small	low	none
MUSIC	high	high	medium
Min-Norm	medium	high	small
ESPRIT	medium	very high	none



# MATLAB

---

Useful MATLAB functions:

- ▶ `hoyw()`
- ▶ `music()`
- ▶ `minnorm()`
- ▶ `esprit()`
- ▶ `freqaphi()`
- ▶ `lsa()`

# Summary

---

- ▶ Alternatives to NLS, based on ARMA and covariance model
- ▶ HOYW, MUSIC, Min-Norm, ESPRIT
- ▶ Complicated derivation but easy to use (and implement)

# Summary

---

- ▶ Alternatives to NLS, based on ARMA and covariance model
- ▶ HOYW, MUSIC, Min-Norm, ESPRIT
- ▶ Complicated derivation but easy to use (and implement)

## Next lecture:

- ▶ How to choose the model order?
- ▶ Information criteria
- ▶ Heuristic approaches
- ▶ Cross validation