




Nonparametric methods: The Periodogram and the Correlogram



Per Mattsson

Systems and Control
Department of Information Technology
Uppsala University

2019-09-05

Summary

The Spectral Estimation Problem

From a finite dataset $\{y(1), y(2), \dots, y(N)\}$ of a stationary signal, determine an estimate $\hat{\phi}(\omega)$ of the PSD $\phi(\omega)$ for $\omega \in [-\pi, \pi]$

- ▶ Power spectral density (PSD)

$$\phi(\omega) = \lim_{N \rightarrow \infty} E \left\{ \frac{1}{N} \left| \sum_{t=1}^N y(t) e^{-i\omega t} \right|^2 \right\}$$

- ▶ Autocovariance sequence (ACS)

$$r(k) = E\{y(t)y^*(t-k)\}$$

- ▶ DTFT of the ACS gives PSD

$$\phi(\omega) = \sum_{k=-\infty}^{\infty} r(k) e^{-i\omega k}$$



Today

- ▶ Estimation theory.
- ▶ The Periodogram and Correlogram methods
- ▶ Properties and equivalences
- ▶ Computing the periodogram in practice.
- ▶ Useful Matlab functions.



Time allocation during the course

- ▶ **Lectures:** Present key concepts and methods, Q&A
- ▶ **Exercise sessions:** You/we solve theory problems (prove/show/verify something)
- ▶ **Labs/homeworks:** Simulate things in MATLAB and try the methods
- ▶ **All over:** Discuss properties, interpretations, and intuition

Time allocation during the course

- ▶ **Lectures:** Present key concepts and methods, Q&A
- ▶ **Exercise sessions:** You/we solve theory problems (prove/show/verify something)
- ▶ **Labs/homeworks:** Simulate things in MATLAB and try the methods
- ▶ **All over:** Discuss properties, interpretations, and intuition

Beyond the lectures

- ▶ You will need to **read more in the book** to get the full picture
- ▶ The **computer labs** will let you compare theory and practice in a systematical way
- ▶ The homework will require you to **code**, try different approaches, make engineering decisions, and present your results



Estimation theory



Estimation theory



Estimation theory

- ▶ Want to estimate some quantity θ from data y .

Estimation theory

- ▶ Want to estimate some quantity θ from data y .
- ▶ Often some model connects θ and y , e.g.,

$$y = f(\theta) + e$$

Estimation theory

- ▶ Want to estimate some quantity θ from data y .
- ▶ Often some model connects θ and y , e.g.,

$$y = f(\theta) + e$$

- ▶ Uncertainty/noise in measurements: $e \implies y$ is stochastic

Estimation theory

- ▶ Want to estimate some quantity θ from data y .
- ▶ Often some model connects θ and y , e.g.,

$$y = f(\theta) + e$$

- ▶ Uncertainty/noise in measurements: $e \implies y$ is stochastic
- ▶ Fit model to data.

Estimation theory

- ▶ Want to estimate some quantity θ from data y .
- ▶ Often some model connects θ and y , e.g.,

$$y = f(\theta) + e$$

- ▶ Uncertainty/noise in measurements: $e \implies y$ is stochastic
- ▶ Fit model to data. E.g., minimize the least square error:

$$\hat{\theta} = \operatorname{argmin}_{\theta} \|y - f(\theta)\|^2$$

Estimation theory

- ▶ Want to estimate some quantity θ from data \mathbf{y} .
- ▶ Often some model connects θ and \mathbf{y} , e.g.,

$$\mathbf{y} = \mathbf{f}(\theta) + \mathbf{e}$$

- ▶ Uncertainty/noise in measurements: $\mathbf{e} \implies \mathbf{y}$ is stochastic
- ▶ Fit model to data. E.g., minimize the least square error:

$$\hat{\theta} = \underset{\theta}{\operatorname{argmin}} \|\mathbf{y} - \mathbf{f}(\theta)\|^2$$

- ▶ $\hat{\theta}$ can be seen as stochastic since \mathbf{y} is stochastic.

Estimation theory

- ▶ Want to estimate some quantity θ from data \mathbf{y} .
- ▶ Often some model connects θ and \mathbf{y} , e.g.,

$$\mathbf{y} = \mathbf{f}(\theta) + \mathbf{e}$$

- ▶ Uncertainty/noise in measurements: $\mathbf{e} \implies \mathbf{y}$ is stochastic
- ▶ Fit model to data. E.g., minimize the least square error:

$$\hat{\theta} = \underset{\theta}{\operatorname{argmin}} \|\mathbf{y} - \mathbf{f}(\theta)\|^2$$

- ▶ $\hat{\theta}$ can be seen as stochastic since \mathbf{y} is stochastic.
- ▶ For any experiment (collected data \mathbf{y}) you get one realization $\hat{\theta}$. If you redo the experiment, you will probably get a different $\hat{\theta}$.

Estimation theory

- ▶ Want to estimate some quantity θ from data y .
- ▶ Often some model connects θ and y , e.g.,

$$y = f(\theta) + e$$

- ▶ Uncertainty/noise in measurements: $e \implies y$ is stochastic
- ▶ Fit model to data. E.g., minimize the least square error:

$$\hat{\theta} = \underset{\theta}{\operatorname{argmin}} \|y - f(\theta)\|^2$$

- ▶ $\hat{\theta}$ can be seen as stochastic since y is stochastic.
- ▶ For any experiment (collected data y) you get one realization $\hat{\theta}$. If you redo the experiment, you will probably get a different $\hat{\theta}$.
- ▶ With this view, we can talk about the expected value and variance of $\hat{\theta}$.



Bias, variance, consistency

Bias, variance, consistency

- ▶ Let $\hat{\theta}$ be the estimate of θ .

Bias, variance, consistency

- ▶ Let $\hat{\theta}$ be the estimate of θ .
- ▶ **Bias:** $E\{\hat{\theta}\} - \theta$

Bias, variance, consistency

- ▶ Let $\hat{\theta}$ be the estimate of θ .
- ▶ **Bias:** $E\{\hat{\theta}\} - \theta$
- ▶ **Variance:** $E\{|\hat{\theta} - E\{\hat{\theta}\}|^2\}$

Bias, variance, consistency

- ▶ Let $\hat{\theta}$ be the estimate of θ .
- ▶ **Bias:** $E\{\hat{\theta}\} - \theta$
- ▶ **Variance:** $E\{|\hat{\theta} - E\{\hat{\theta}\}|^2\}$
- ▶ **Mean squared error:**

$$\text{MSE} = E\{|\hat{\theta} - \theta|^2\} =$$

Bias, variance, consistency

- ▶ Let $\hat{\theta}$ be the estimate of θ .
- ▶ **Bias:** $E\{\hat{\theta}\} - \theta$
- ▶ **Variance:** $E\{|\hat{\theta} - E\{\hat{\theta}\}|^2\}$
- ▶ **Mean squared error:**

$$\text{MSE} = E\{|\hat{\theta} - \theta|^2\} = \text{var}\{\hat{\theta}\} + |\text{bias}\{\hat{\theta}\}|^2$$

Bias, variance, consistency

- ▶ Let $\hat{\theta}$ be the estimate of θ .
- ▶ **Bias:** $E\{\hat{\theta}\} - \theta$
- ▶ **Variance:** $E\{|\hat{\theta} - E\{\hat{\theta}\}|^2\}$
- ▶ **Mean squared error:**

$$\text{MSE} = E\{|\hat{\theta} - \theta|^2\} = \text{var}\{\hat{\theta}\} + |\text{bias}\{\hat{\theta}\}|^2$$

- ▶ **Consistency:** The way we estimate (our estimator) is consistent if $\hat{\theta} \rightarrow \theta$ as the number of data points used for estimation goes to infinity.

Parametric vs. non-parametric

Nonparametric (Classical) Methods

No model imposing constraints, or rather, a very general model (i.e. many “parameters”)

Parametric (Modern) Methods

Model the data and estimate the model parameters.

Parametric vs. non-parametric

Nonparametric (Classical) Methods

No model imposing constraints, or rather, a very general model (i.e. many “parameters”)

Parametric (Modern) Methods

Model the data and estimate the model parameters.

Trade-offs: (Robustness, and Resolution vs. Variance)

- ▶ **Parametric methods:** better estimates **if** model is good, otherwise the results can be heavily biased.
- ▶ **Non-parametric methods:** works for any dataset, but have relatively high variance, even when $N \rightarrow \infty$.
- ▶ **Resolution** – detecting/separating spectral components.



Periodogram and correlogram

Non-parametric PSD estimation

Both

$$\phi(\omega) = \lim_{N \rightarrow \infty} E \left\{ \frac{1}{N} \left| \sum_{t=1}^N y(t) e^{-i\omega t} \right|^2 \right\}$$

and

$$\phi(\omega) = \sum_{k=-\infty}^{\infty} r(k) e^{-i\omega k} \quad \text{where} \quad r(k) = E\{y(t)y^*(t-k)\}$$

require an **infinite** number of samples ($N \rightarrow \infty$).

Non-parametric PSD estimation

Both

$$\phi(\omega) = \lim_{N \rightarrow \infty} E \left\{ \frac{1}{N} \left| \sum_{t=1}^N y(t) e^{-i\omega t} \right|^2 \right\}$$

and

$$\phi(\omega) = \sum_{k=-\infty}^{\infty} r(k) e^{-i\omega k} \quad \text{where} \quad r(k) = E\{y(t)y^*(t-k)\}$$

require an **infinite** number of samples ($N \rightarrow \infty$).

- **Problem:** Finite collection of samples $\{y(1), \dots, y(N)\}$, so cannot compute the sums and expectations.

Non-parametric PSD estimation

Both

$$\phi(\omega) = \lim_{N \rightarrow \infty} E \left\{ \frac{1}{N} \left| \sum_{t=1}^N y(t) e^{-i\omega t} \right|^2 \right\}$$

and

$$\phi(\omega) = \sum_{k=-\infty}^{\infty} r(k) e^{-i\omega k} \quad \text{where} \quad r(k) = E\{y(t)y^*(t-k)\}$$

require an **infinite** number of samples ($N \rightarrow \infty$).

- ▶ **Problem:** Finite collection of samples $\{y(1), \dots, y(N)\}$, so cannot compute the sums and expectations.
- ▶ **Idea:**

Non-parametric PSD estimation

Both

$$\phi(\omega) = \lim_{N \rightarrow \infty} E \left\{ \frac{1}{N} \left| \sum_{t=1}^N y(t) e^{-i\omega t} \right|^2 \right\}$$

and

$$\phi(\omega) = \sum_{k=-\infty}^{\infty} r(k) e^{-i\omega k} \quad \text{where} \quad r(k) = E\{y(t)y^*(t-k)\}$$

require an **infinite** number of samples ($N \rightarrow \infty$).

- ▶ **Problem:** Finite collection of samples $\{y(1), \dots, y(N)\}$, so cannot compute the sums and expectations.
- ▶ **Idea:** Skip $\lim_{N \rightarrow \infty}$ and $E\{\cdot\}$.

Non-parametric PSD estimation

Both

$$\phi(\omega) = \lim_{N \rightarrow \infty} E \left\{ \frac{1}{N} \left| \sum_{t=1}^N y(t) e^{-i\omega t} \right|^2 \right\}$$

and

$$\phi(\omega) = \sum_{k=-\infty}^{\infty} r(k) e^{-i\omega k} \quad \text{where} \quad r(k) = E\{y(t)y^*(t-k)\}$$

require an **infinite** number of samples ($N \rightarrow \infty$).

- ▶ **Problem:** Finite collection of samples $\{y(1), \dots, y(N)\}$, so cannot compute the sums and expectations.
- ▶ **Idea:** Skip $\lim_{N \rightarrow \infty}$ and $E\{\cdot\}$.
- ▶ Idea used by Schuster (~ 1900) to determine “hidden periodicities”.

Non-parametric PSD estimation

$$\phi(\omega) = \lim_{N \rightarrow \infty} E \left\{ \frac{1}{N} \left| \sum_{t=1}^N y(t) e^{-i\omega t} \right|^2 \right\} = \sum_{k=-\infty}^{\infty} r(k) e^{-i\omega k}$$

Non-parametric PSD estimation

$$\phi(\omega) = \lim_{N \rightarrow \infty} E \left\{ \frac{1}{N} \left| \sum_{t=1}^N y(t) e^{-i\omega t} \right|^2 \right\} = \sum_{k=-\infty}^{\infty} r(k) e^{-i\omega k}$$

► **Periodogram:** The first definition gives

$$\hat{\phi}_p(\omega) = \frac{1}{N} \left| \sum_{t=1}^N y(t) e^{-i\omega t} \right|^2.$$

Non-parametric PSD estimation

$$\phi(\omega) = \lim_{N \rightarrow \infty} E \left\{ \frac{1}{N} \left| \sum_{t=1}^N y(t) e^{-i\omega t} \right|^2 \right\} = \sum_{k=-\infty}^{\infty} r(k) e^{-i\omega k}$$

- **Periodogram:** The first definition gives

$$\hat{\phi}_p(\omega) = \frac{1}{N} \left| \sum_{t=1}^N y(t) e^{-i\omega t} \right|^2.$$

- **Correlogram:** The second definition gives

$$\hat{\phi}_c(\omega) = \sum_{k=-(N-1)}^{N-1} \hat{r}(k) e^{-i\omega k},$$

where $\hat{r}(k)$ is an estimate of $r(k) = E\{y(t)y^*(t-k)\}$.

Non-parametric PSD estimation

$$\phi(\omega) = \lim_{N \rightarrow \infty} E \left\{ \frac{1}{N} \left| \sum_{t=1}^N y(t) e^{-i\omega t} \right|^2 \right\} = \sum_{k=-\infty}^{\infty} r(k) e^{-i\omega k}$$

- **Periodogram:** The first definition gives

$$\hat{\phi}_p(\omega) = \frac{1}{N} \left| \sum_{t=1}^N y(t) e^{-i\omega t} \right|^2.$$

- **Correlogram:** The second definition gives

$$\hat{\phi}_c(\omega) = \sum_{k=-(N-1)}^{N-1} \hat{r}(k) e^{-i\omega k},$$

where $\hat{r}(k)$ is an estimate of $r(k) = E\{y(t)y^*(t-k)\}$.

- Finite sums \implies can be computed in practice.

Estimation of ACS

Standard unbiased estimate:

$$\hat{r}(k) = \frac{1}{N-k} \sum_{t=k+1}^N y(t)y^*(t-k), \quad 0 \leq k \leq N-1.$$

Estimation of ACS

Standard unbiased estimate:

$$\hat{r}(k) = \frac{1}{N-k} \sum_{t=k+1}^N y(t)y^*(t-k), \quad 0 \leq k \leq N-1.$$

Problems:

- For k large (close to N), $\hat{r}(k)$ has high variance (erratic).

Estimation of ACS

Standard unbiased estimate:

$$\hat{r}(k) = \frac{1}{N-k} \sum_{t=k+1}^N y(t)y^*(t-k), \quad 0 \leq k \leq N-1.$$

Problems:

- ▶ For k large (close to N), $\hat{r}(k)$ has high variance (erratic).
- ▶ May result in negative spectrum.

Estimation of ACS

Standard unbiased estimate:

$$\hat{r}(k) = \frac{1}{N-k} \sum_{t=k+1}^N y(t)y^*(t-k), \quad 0 \leq k \leq N-1.$$

Problems:

- ▶ For k large (close to N), $\hat{r}(k)$ has high variance (erratic).
- ▶ May result in negative spectrum.

Standard biased estimate:

$$\hat{r}(k) = \frac{1}{N} \sum_{t=k+1}^N y(t)y^*(t-k), \quad 0 \leq k \leq N-1.$$

Estimation of ACS

Standard unbiased estimate:

$$\hat{r}(k) = \frac{1}{N-k} \sum_{t=k+1}^N y(t)y^*(t-k), \quad 0 \leq k \leq N-1.$$

Problems:

- ▶ For k large (close to N), $\hat{r}(k)$ has high variance (erratic).
- ▶ May result in negative spectrum.

Standard biased estimate:

$$\hat{r}(k) = \frac{1}{N} \sum_{t=k+1}^N y(t)y^*(t-k), \quad 0 \leq k \leq N-1.$$

- ▶ $r(k)$ is typically small for large k , and bias is low for small k .

Estimation of ACS

Standard unbiased estimate:

$$\hat{r}(k) = \frac{1}{N-k} \sum_{t=k+1}^N y(t)y^*(t-k), \quad 0 \leq k \leq N-1.$$

Problems:

- ▶ For k large (close to N), $\hat{r}(k)$ has high variance (erratic).
- ▶ May result in negative spectrum.

Standard biased estimate:

$$\hat{r}(k) = \frac{1}{N} \sum_{t=k+1}^N y(t)y^*(t-k), \quad 0 \leq k \leq N-1.$$

- ▶ $r(k)$ is typically small for large k , and bias is low for small k .
- ▶ $\hat{r}(k)$ positive semidefinite \implies non-negative spectrum.

Estimation of ACS

Standard unbiased estimate:

$$\hat{r}(k) = \frac{1}{N-k} \sum_{t=k+1}^N y(t)y^*(t-k), \quad 0 \leq k \leq N-1.$$

Problems:

- ▶ For k large (close to N), $\hat{r}(k)$ has high variance (erratic).
- ▶ May result in negative spectrum.

Standard biased estimate:

$$\hat{r}(k) = \frac{1}{N} \sum_{t=k+1}^N y(t)y^*(t-k), \quad 0 \leq k \leq N-1.$$

- ▶ $r(k)$ is typically small for large k , and bias is low for small k .
- ▶ $\hat{r}(k)$ positive semidefinite \implies non-negative spectrum.

For both estimates: $\hat{r}(k) = \hat{r}^*(-k)$ for $k < 0$.

Equivalence

If the biased ACS estimator $\hat{r}(k)$ is used in $\hat{\phi}_c(\omega)$, then

$$\begin{aligned}\hat{\phi}_p(\omega) &= \frac{1}{N} \left| \sum_{t=1}^N y(t) e^{-i\omega t} \right|^2 \\ &= \sum_{k=-(N-1)}^{N-1} \hat{r}(k) e^{-i\omega k} = \hat{\phi}_c(\omega)\end{aligned}$$

$$\hat{\phi}_p(\omega) = \hat{\phi}_c(\omega)$$

- Can analyze the estimates simultaneously!



Analysis

Analysis of the periodogram

- ▶ Since $\phi_p(\omega) = \phi_c(\omega)$ we can use both formulations in the analysis.
- ▶ Bias-variance tradeoff:

$$\text{MSE} = \text{var}(\hat{\phi}_p(\omega)) + |\text{bias}(\hat{\phi}_p(\omega))|^2.$$

- ▶ **Wanted:** Small bias, small variance, consistent.

The bias of the periodogram

- ▶ With the biased ACS estimate we get

$$E\{\hat{\phi}_c(\omega)\} =$$

The bias of the periodogram

- With the biased ACS estimate we get

$$E\{\hat{\phi}_c(\omega)\} = \sum_{k=-(N-1)}^{N-1} E\{\hat{r}(k)\}e^{-i\omega k} = [\text{Board}]$$
$$=$$

The bias of the periodogram

- With the biased ACS estimate we get

$$\begin{aligned} E\{\hat{\phi}_c(\omega)\} &= \sum_{k=-(N-1)}^{N-1} E\{\hat{r}(k)\} e^{-i\omega k} = \text{[Board]} \\ &= \sum_{k=-(N-1)}^{N-1} \left(1 - \frac{|k|}{N}\right) r(k) e^{-i\omega k} \\ &= \end{aligned}$$

The bias of the periodogram

- With the biased ACS estimate we get

$$\begin{aligned} E\{\hat{\phi}_c(\omega)\} &= \sum_{k=-(N-1)}^{N-1} E\{\hat{r}(k)\} e^{-i\omega k} = \text{[Board]} \\ &= \sum_{k=-(N-1)}^{N-1} \left(1 - \frac{|k|}{N}\right) r(k) e^{-i\omega k} \\ &= \sum_{k=-\infty}^{\infty} w_B(k) r(k) e^{-i\omega k}, \end{aligned}$$

where the **Bartlett (triangular) window** is given by

$$w_B(k) = \begin{cases} 1 - \frac{|k|}{N}, & |k| \leq N-1 \\ 0, & |k| \geq N \end{cases}.$$

The bias of the periodogram

- ▶ With the biased ACS estimate we get

$$\begin{aligned} E\{\hat{\phi}_c(\omega)\} &= \sum_{k=-(N-1)}^{N-1} E\{\hat{r}(k)\} e^{-i\omega k} = \text{[Board]} \\ &= \sum_{k=-(N-1)}^{N-1} \left(1 - \frac{|k|}{N}\right) r(k) e^{-i\omega k} \\ &= \sum_{k=-\infty}^{\infty} w_B(k) r(k) e^{-i\omega k}, \end{aligned}$$

where the **Bartlett (triangular) window** is given by

$$w_B(k) = \begin{cases} 1 - \frac{|k|}{N}, & |k| \leq N-1 \\ 0, & |k| \geq N \end{cases}.$$

- ▶ Large bias for small N since $w_B(k) \neq 1$.

The bias of the periodogram

- $E\{\hat{\phi}_c(\omega)\}$ is the DTFT of $w_B(k)r(k)$.

The bias of the periodogram

- ▶ $E\{\hat{\phi}_c(\omega)\}$ is the DTFT of $w_B(k)r(k)$.
- ▶ Time-domain multiplication \iff Frequency-domain convolution

The bias of the periodogram

- ▶ $E\{\hat{\phi}_c(\omega)\}$ is the DTFT of $w_B(k)r(k)$.
- ▶ Time-domain multiplication \iff Frequency-domain convolution:

$$E\{\hat{\phi}(\omega)\} = \frac{1}{2\pi} \int_{-\pi}^{\pi} \phi(\zeta) W_B(\omega - \zeta) d\zeta$$

The bias of the periodogram

- ▶ $E\{\hat{\phi}_c(\omega)\}$ is the DTFT of $w_B(k)r(k)$.
- ▶ Time-domain multiplication \iff Frequency-domain convolution:

$$E\{\hat{\phi}(\omega)\} = \frac{1}{2\pi} \int_{-\pi}^{\pi} \phi(\zeta) W_B(\omega - \zeta) d\zeta,$$

where the DTFT of w_B is given by

$$W_B(\omega) = \frac{1}{N} \left(\frac{\sin(\omega N/2)}{\sin(\omega/2)} \right)^2.$$

The bias of the periodogram

- ▶ $E\{\hat{\phi}_c(\omega)\}$ is the DTFT of $w_B(k)r(k)$.
- ▶ Time-domain multiplication \iff Frequency-domain convolution:

$$E\{\hat{\phi}(\omega)\} = \frac{1}{2\pi} \int_{-\pi}^{\pi} \phi(\zeta) W_B(\omega - \zeta) d\zeta,$$

where the DTFT of w_B is given by

$$W_B(\omega) = \frac{1}{N} \left(\frac{\sin(\omega N/2)}{\sin(\omega/2)} \right)^2.$$

- ▶ Ideally $W_B(\omega)$ should approximate a Dirac impulse $\delta(\omega)$.

The Bartlett window




Figure 2.1. $W_B(\omega)/W_B(0)$, for $N = 25$.

The Bartlett window




Figure 2.1. $W_B(\omega)/W_B(0)$, for $N = 25$.

- Main lobe 3dB width $\approx 2\pi/N$ radians (or $1/N$ in frequency).

The Bartlett window




Figure 2.1. $W_B(\omega)/W_B(0)$, for $N = 25$.

- ▶ Main lobe 3dB width $\approx 2\pi/N$ radians (or $1/N$ in frequency).
- ▶ For small N , quite different from $\delta(\omega)$ (severe bias).

The Bartlett window




Figure 2.1. $W_B(\omega)/W_B(0)$, for $N = 25$.

- ▶ Main lobe 3dB width $\approx 2\pi/N$ radians (or $1/N$ in frequency).
- ▶ For small N , quite different from $\delta(\omega)$ (severe bias).
- ▶ Non-zero main lobe \implies Smearing (smoothing).

The Bartlett window




Figure 2.1. $W_B(\omega)/W_B(0)$, for $N = 25$.

- ▶ Main lobe 3dB width $\approx 2\pi/N$ radians (or $1/N$ in frequency).
- ▶ For small N , quite different from $\delta(\omega)$ (severe bias).
- ▶ Non-zero main lobe \implies Smearing (smoothing).
- ▶ Sidelobes \implies Leakage.

Summary of the periodogram bias properties

- ▶ For small N , severe bias.

Summary of the periodogram bias properties

- ▶ For small N , severe bias.
- ▶ However,

$$\lim_{N \rightarrow \infty} W_B(\omega) = 2\pi\delta(\omega),$$

Summary of the periodogram bias properties

- ▶ For small N , severe bias.
- ▶ However,

$$\lim_{N \rightarrow \infty} W_B(\omega) = 2\pi\delta(\omega),$$

so,

$$\lim_{N \rightarrow \infty} E\{\hat{\phi}_p(\omega)\} = \int_{-\pi}^{\pi} \phi(\zeta)\delta(\omega - \zeta)d\zeta =$$

Summary of the periodogram bias properties

- ▶ For small N , severe bias.
- ▶ However,

$$\lim_{N \rightarrow \infty} W_B(\omega) = 2\pi\delta(\omega),$$

so,

$$\lim_{N \rightarrow \infty} E\{\hat{\phi}_p(\omega)\} = \int_{-\pi}^{\pi} \phi(\zeta)\delta(\omega - \zeta)d\zeta = \phi(\omega).$$

Summary of the periodogram bias properties

- ▶ For small N , severe bias.
- ▶ However,

$$\lim_{N \rightarrow \infty} W_B(\omega) = 2\pi\delta(\omega),$$

so,

$$\lim_{N \rightarrow \infty} E\{\hat{\phi}_p(\omega)\} = \int_{-\pi}^{\pi} \phi(\zeta)\delta(\omega - \zeta)d\zeta = \phi(\omega).$$

- ▶ That is, the periodogram is **asymptotically unbiased**.

Summary of the periodogram bias properties

- ▶ For small N , severe bias.
- ▶ However,

$$\lim_{N \rightarrow \infty} W_B(\omega) = 2\pi\delta(\omega),$$

so,

$$\lim_{N \rightarrow \infty} E\{\hat{\phi}_p(\omega)\} = \int_{-\pi}^{\pi} \phi(\zeta)\delta(\omega - \zeta)d\zeta = \phi(\omega).$$

- ▶ That is, the periodogram is **asymptotically unbiased**.
- ▶ If bias was the only problem with periodogram, we could solve it by increasing the number of data points used. However....

The asymptotic variance/covariance

► As $N \rightarrow \infty$,

$$E \left\{ \left[\hat{\phi}_p(\omega_1) - \phi(\omega_1) \right] \left[\hat{\phi}_p(\omega_2) - \phi(\omega_2) \right] \right\}$$

=

The asymptotic variance/covariance

► As $N \rightarrow \infty$,

$$\begin{aligned} E \left\{ \left[\hat{\phi}_p(\omega_1) - \phi(\omega_1) \right] \left[\hat{\phi}_p(\omega_2) - \phi(\omega_2) \right] \right\} \\ = \begin{cases} \phi^2(\omega_1), & \omega_1 = \omega_2 \\ 0, & \omega_1 \neq \omega_2 \end{cases} \end{aligned}$$

The asymptotic variance/covariance

- As $N \rightarrow \infty$,

$$\begin{aligned} E \left\{ \left[\hat{\phi}_p(\omega_1) - \phi(\omega_1) \right] \left[\hat{\phi}_p(\omega_2) - \phi(\omega_2) \right] \right\} \\ = \begin{cases} \phi^2(\omega_1), & \omega_1 = \omega_2 \\ 0, & \omega_1 \neq \omega_2 \end{cases} \end{aligned}$$

- **Inconsistent estimate.** In general $\lim_{N \rightarrow \infty} \hat{\phi}_p(\omega) \neq \phi(\omega)$.

The asymptotic variance/covariance

- As $N \rightarrow \infty$,

$$\begin{aligned} E \left\{ \left[\hat{\phi}_p(\omega_1) - \phi(\omega_1) \right] \left[\hat{\phi}_p(\omega_2) - \phi(\omega_2) \right] \right\} \\ = \begin{cases} \phi^2(\omega_1), & \omega_1 = \omega_2 \\ 0, & \omega_1 \neq \omega_2 \end{cases} \end{aligned}$$


- **Inconsistent estimate.** In general $\lim_{N \rightarrow \infty} \hat{\phi}_p(\omega) \neq \phi(\omega)$.
- **Erratic behavior.** Estimate at ω_1 uncorrelated (for large N) from estimate at $\omega_2 \neq \omega_1$.

The asymptotic variance/covariance

- As $N \rightarrow \infty$,

$$E \left\{ \left[\hat{\phi}_p(\omega_1) - \phi(\omega_1) \right] \left[\hat{\phi}_p(\omega_2) - \phi(\omega_2) \right] \right\} = \begin{cases} \phi^2(\omega_1), & \omega_1 = \omega_2 \\ 0, & \omega_1 \neq \omega_2 \end{cases}$$

- **Inconsistent estimate.** In general $\lim_{N \rightarrow \infty} \hat{\phi}_p(\omega) \neq \phi(\omega)$.
- **Erratic behavior.** Estimate at ω_1 uncorrelated (for large N) from estimate at $\omega_2 \neq \omega_1$.



Summary of periodogram properties

$$\hat{\phi}_p = \frac{1}{N} \left| \sum_{t=1}^N y(t) e^{-i\omega t} \right|^2 = \sum_{k=-(N-1)}^{N-1} \hat{r}(k) e^{-i\omega k} = \phi_c(\omega)$$

- ▶ Asymptotically unbiased:

$$\lim_{N \rightarrow \infty} E\{\hat{\phi}_p(\omega)\} = \phi(\omega).$$

- ▶ Large variance, even as $N \rightarrow \infty$.
- ▶ Thus, $\hat{\phi}(\omega)$ has bad performance.

Summary of periodogram properties

$$\hat{\phi}_p = \frac{1}{N} \left| \sum_{t=1}^N y(t) e^{-i\omega t} \right|^2 = \sum_{k=-(N-1)}^{N-1} \hat{r}(k) e^{-i\omega k} = \phi_c(\omega)$$

- ▶ Asymptotically unbiased:

$$\lim_{N \rightarrow \infty} E\{\hat{\phi}_p(\omega)\} = \phi(\omega).$$

- ▶ Large variance, even as $N \rightarrow \infty$.
- ▶ Thus, $\hat{\phi}(\omega)$ has bad performance.

Intuitive explanation:

- ▶ $\hat{r}(k) - r(k)$ may be large for large $|k|$.
- ▶ Even if this is not case, the sum of many small errors can become large.



Computing the periodogram

Computing the periodogram

$$\phi_p(\omega) = \frac{1}{N} |Y_N(\omega)|^2, \quad Y_N(\omega) = \sum_{t=1}^N y(t) e^{-i\omega t}.$$

Computing the periodogram

$$\phi_p(\omega) = \frac{1}{N} |Y_N(\omega)|^2, \quad Y_N(\omega) = \sum_{t=1}^N y(t) e^{-i\omega t}.$$

- ▶ Can only compute for a finite number of ω .

Computing the periodogram

$$\phi_p(\omega) = \frac{1}{N} |Y_N(\omega)|^2, \quad Y_N(\omega) = \sum_{t=1}^N y(t) e^{-i\omega t}.$$

- ▶ Can only compute for a finite number of ω .
- ▶ For large N , heavy computations (sum of N complex multiplications for each frequency).

Computing the periodogram

$$\phi_p(\omega) = \frac{1}{N} |Y_N(\omega)|^2, \quad Y_N(\omega) = \sum_{t=1}^N y(t) e^{-i\omega t}.$$

- ▶ Can only compute for a finite number of ω .
- ▶ For large N , heavy computations (sum of N complex multiplications for each frequency).

Common solution:

- ▶ Compute for $\omega = \frac{2\pi}{N}k$, $k = 0, 1, \dots, N-1$.

Computing the periodogram

$$\phi_p(\omega) = \frac{1}{N} |Y_N(\omega)|^2, \quad Y_N(\omega) = \sum_{t=1}^N y(t) e^{-i\omega t}.$$

- ▶ Can only compute for a finite number of ω .
- ▶ For large N , heavy computations (sum of N complex multiplications for each frequency).

Common solution:

- ▶ Compute for $\omega = \frac{2\pi}{N}k$, $k = 0, 1, \dots, N-1$.
- ▶ Then we obtain the DFT,

$$Y(k) = \sum_{t=1}^N y(t) e^{-i2\pi kt/N}.$$

Computing the periodogram

$$\phi_p(\omega) = \frac{1}{N} |Y_N(\omega)|^2, \quad Y_N(\omega) = \sum_{t=1}^N y(t) e^{-i\omega t}.$$

- ▶ Can only compute for a finite number of ω .
- ▶ For large N , heavy computations (sum of N complex multiplications for each frequency).

Common solution:

- ▶ Compute for $\omega = \frac{2\pi}{N}k$, $k = 0, 1, \dots, N-1$.
- ▶ Then we obtain the DFT,

$$Y(k) = \sum_{t=1}^N y(t) e^{-i2\pi kt/N}.$$

- ▶ Can be computed using Fast Fourier Transform (FFT) (computations in the order of $N \log_2 N$ vs N^2 for direct approach). Often assumes $N = 2^m$.

Zeropadding

- ▶ Append the data vector given to FFT with zeros,

$$\underbrace{\{y(1), \dots, y(N), 0, \dots, 0\}}_L$$

Zeropadding

- ▶ Append the data vector given to FFT with zeros,

$$\underbrace{\{y(1), \dots, y(N), 0, \dots, 0\}}_L$$

- ▶ Does not change $\hat{\phi}_p(\omega)$,

$$\hat{\phi}_p(\omega) = \frac{1}{N} \left| \sum_{t=1}^L y(t) e^{-i\omega t} \right|^2, \quad \text{with } y(t) = 0 \text{ for } t > N.$$

Zeropadding

- ▶ Append the data vector given to FFT with zeros,

$$\underbrace{\{y(1), \dots, y(N), 0, \dots, 0\}}_L$$

- ▶ Does not change $\hat{\phi}_p(\omega)$,

$$\hat{\phi}_p(\omega) = \frac{1}{N} \left| \sum_{t=1}^L y(t) e^{-i\omega t} \right|^2, \quad \text{with } y(t) = 0 \text{ for } t > N.$$

- ▶ Can ensure that $L = 2^m$ (not necessary for `fft` in Matlab).

Zeropadding

- ▶ Append the data vector given to FFT with zeros,

$$\underbrace{\{y(1), \dots, y(N), 0, \dots, 0\}}_L$$

- ▶ Does not change $\hat{\phi}_p(\omega)$,

$$\hat{\phi}_p(\omega) = \frac{1}{N} \left| \sum_{t=1}^L y(t) e^{-i\omega t} \right|^2, \quad \text{with } y(t) = 0 \text{ for } t > N.$$

- ▶ Can ensure that $L = 2^m$ (not necessary for `fft` in Matlab).
- ▶ Increases the number of frequencies computed by FFT,

$$\omega = \frac{2\pi}{N} k, \quad k = 0, 1, \dots, L - 1.$$

Zeropadding

- ▶ Append the data vector given to FFT with zeros,

$$\underbrace{\{y(1), \dots, y(N), 0, \dots, 0\}}_L$$

- ▶ Does not change $\hat{\phi}_p(\omega)$,


$$\hat{\phi}_p(\omega) = \frac{1}{N} \left| \sum_{t=1}^L y(t) e^{-i\omega t} \right|^2, \quad \text{with } y(t) = 0 \text{ for } t > N.$$

- ▶ Can ensure that $L = 2^m$ (not necessary for `fft` in Matlab).
- ▶ Increases the number of frequencies computed by FFT,

$$\omega = \frac{2\pi}{N} k, \quad k = 0, 1, \dots, L - 1.$$


- ▶ There are FFT algorithms that eliminate operations on zeros.

Example: Zeropadding



6 samples (no zeropadding)

Example: Zeropadding



60 samples ($L = 10N$ zeropadding)

Summary

- ▶ The periodogram and correlogram.
- ▶ Smearing and leakage.
- ▶ Asymptotically unbiased.
- ▶ “High” variance even when $N \rightarrow \infty$.
- ▶ Using the FFT for computing the periodogram.

Summary

- ▶ The periodogram and correlogram.
- ▶ Smearing and leakage.
- ▶ Asymptotically unbiased.
- ▶ “High” variance even when $N \rightarrow \infty$.
- ▶ Using the FFT for computing the periodogram.

Try the following:

- ▶ Simulate a sinusoid/complex exponential in noise and plot the Periodogram (compute the periodogram manually and with FFT!)
- ▶ Add another periodic component, study the effect of smearing.
- ▶ Exercise 1.5, 1.7, 1.10, 2.3, 2.6

MATLAB

Useful functions:

- ▶ `randn(N,1)` – Generates Gaussian white noise.
- ▶ `fft(y,L)` – Performs FFT padded with zeros.
- ▶ `fftshift(phi)` – Shifts frequencies from $[0, 2\pi]$ to $[-\pi, \pi]$.
- ▶ `periodogramse(y,v,L)` – (Windowed) periodogram.
- ▶ `filter(B,A,e)` – Generate $y(t) = \frac{B(q)}{A(q)}e(t)$. A and B are vectors with the coefficients of the polynomials $A(q), B(q)$.

MATLAB

Useful functions:

- ▶ `randn(N,1)` – Generates Gaussian white noise.
- ▶ `fft(y,L)` – Performs FFT padded with zeros.
- ▶ `fftshift(phi)` – Shifts frequencies from $[0, 2\pi]$ to $[-\pi, \pi]$.
- ▶ `periodogramse(y,v,L)` – (Windowed) periodogram.
- ▶ `filter(B,A,e)` – Generate $y(t) = \frac{B(q)}{A(q)}e(t)$. A and B are vectors with the coefficients of the polynomials $A(q), B(q)$.

With `fft`, the corresponding normalized frequencies are

- » `omega = 0:2*pi/L:2*pi(L-1)/L; %or`
- » `omega = (0:L-1)*2*pi/L;`

MATLAB

Useful functions:

- ▶ `randn(N,1)` – Generates Gaussian white noise.
- ▶ `fft(y,L)` – Performs FFT padded with zeros.
- ▶ `fftshift(phi)` – Shifts frequencies from $[0, 2\pi]$ to $[-\pi, \pi]$.
- ▶ `periodogramse(y,v,L)` – (Windowed) periodogram.
- ▶ `filter(B,A,e)` – Generate $y(t) = \frac{B(q)}{A(q)}e(t)$. A and B are vectors with the coefficients of the polynomials $A(q), B(q)$.

With `fft`, the corresponding normalized frequencies are

- » `omega = 0:2*pi/L:2*pi(L-1)/L; %or`
- » `omega = (0:L-1)*2*pi/L;`

If you have also applied `fftshift`, the normalized frequencies are

- » `omega = (0:L-1)*2*pi/L - pi;`