




# Parametric methods for Rational Spectra



Per Mattsson

Systems and Control  
Department of Information Technology  
Uppsala University

2019-09-13

# Summary from last lecture

## Refined non-parametric methods

$$\text{Blackman-Tukey: } \hat{\phi}_{\text{BT}}(\omega) = \sum_{k=-(M-1)}^{M-1} w(k) \hat{r}(k) e^{-i\omega k}$$

$$\text{Welch: } \hat{\phi}_{\text{W}}(\omega) = \frac{1}{S} \sum_{j=1}^S \frac{1}{MP} \left| \sum_{t=1}^M v(t) y_j(t) e^{-i\omega t} \right|^2$$

- ▶ Windowed Correlogram/Periodogram (smoothing)
- ▶ Window functions (Bartlett, Hamming, Chebyshev, Kaiser)
- ▶ Bias/Variance or Resolution(main)/Leakage(side) trade-off
- ▶ Estimating continuous/line spectra

**Today:** Parametric methods for rational (continuous) spectra

# Parametric methods

---

- ▶ Model the data/PSD using fewer parameters (cf. Periodogram has one for each  $\omega$ , i.e.  $N$  parameters)
- ▶ If the model is poor, the result will be biased.
- ▶ Some prior knowledge needed.

# Parametric methods

- ▶ Model the data/PSD using fewer parameters (cf. Periodogram has one for each  $\omega$ , i.e.  $N$  parameters)
- ▶ If the model is poor, the result will be biased.
- ▶ Some prior knowledge needed.

## Occam's razor or Principle of Parsimony

*If two theories explain the data equally well, then the simpler theory is preferable.*

# Parametric methods

- ▶ Model the data/PSD using fewer parameters (cf. Periodogram has one for each  $\omega$ , i.e.  $N$  parameters)
- ▶ If the model is poor, the result will be biased.
- ▶ Some prior knowledge needed.

## Occam's razor or Principle of Parsimony

*If two theories explain the data equally well, then the simpler theory is preferable.*

- ▶ “Keep it as simple as possible, but no simpler” – Albert Einstein.

# Parametric methods

- ▶ Model the data/PSD using fewer parameters (cf. Periodogram has one for each  $\omega$ , i.e.  $N$  parameters)
- ▶ If the model is poor, the result will be biased.
- ▶ Some prior knowledge needed.

## Occam's razor or Principle of Parsimony

*If two theories explain the data equally well, then the simpler theory is preferable.*

- ▶ “Keep it as simple as possible, but no simpler” – Albert Einstein.
- ▶ “With four parameters I can fit an elephant  
” – John von Neumann.

# Parametric methods

- ▶ Model the data/PSD using fewer parameters (cf. Periodogram has one for each  $\omega$ , i.e.  $N$  parameters)
- ▶ If the model is poor, the result will be biased.
- ▶ Some prior knowledge needed.

## Occam's razor or Principle of Parsimony

*If two theories explain the data equally well, then the simpler theory is preferable.*

- ▶ “Keep it as simple as possible, but no simpler” – Albert Einstein.
- ▶ “With four parameters I can fit an elephant, and with five I can make him wiggle his trunk.” – John von Neumann.

# Parametric methods

- ▶ Model the data/PSD using fewer parameters (cf. Periodogram has one for each  $\omega$ , i.e.  $N$  parameters)
- ▶ If the model is poor, the result will be biased.
- ▶ Some prior knowledge needed.

## Occam's razor or Principle of Parsimony

*If two theories explain the data equally well, then the simpler theory is preferable.*

- ▶ “Keep it as simple as possible, but no simpler” – Albert Einstein.
- ▶ “With four parameters I can fit an elephant, and with five I can make him wiggle his trunk.” – John von Neumann.
- ▶ Bias/variance trade-off



# Parametric methods

- ▶ Model the data/PSD using fewer parameters (cf. Periodogram has one for each  $\omega$ , i.e.  $N$  parameters)
- ▶ If the model is poor, the result will be biased.
- ▶ Some prior knowledge needed.

## Occam's razor or Principle of Parsimony

*If two theories explain the data equally well, then the simpler theory is preferable.*

- ▶ “Keep it as simple as possible, but no simpler” – Albert Einstein.
- ▶ “With four parameters I can fit an elephant, and with five I can make him wiggle his trunk.” – John von Neumann.
- ▶ Bias/variance trade-off
- ▶ Overfitting.

# Parametric methods

- ▶ Model the data/PSD using fewer parameters (cf. Periodogram has one for each  $\omega$ , i.e.  $N$  parameters)
- ▶ If the model is poor, the result will be biased.
- ▶ Some prior knowledge needed.

## Occam's razor or Principle of Parsimony

*If two theories explain the data equally well, then the simpler theory is preferable.*

- ▶ “Keep it as simple as possible, but no simpler” – Albert Einstein.
- ▶ “With four parameters I can fit an elephant, and with five I can make him wiggle his trunk.” – John von Neumann.
- ▶ Bias/variance trade-off
- ▶ Overfitting.
- ▶ Model order selection is important, but difficult (more in L7)



# Signals with rational spectra

# Rational spectra

## Rational PSD

$$\phi(\omega) = \frac{\sum_{k=-m}^m \gamma_k e^{-i\omega k}}{\sum_{k=-n}^n \rho_k e^{-i\omega k}}$$

where  $\gamma_{-k} = \gamma_k^*$  and  $\rho_{-k} = \rho_k^*$ , can approximate any **continuous** PSD for **sufficiently large**  $m, n$

# Rational spectra

## Rational PSD

$$\phi(\omega) = \frac{\sum_{k=-m}^m \gamma_k e^{-i\omega k}}{\sum_{k=-n}^n \rho_k e^{-i\omega k}}$$

where  $\gamma_{-k} = \gamma_k^*$  and  $\rho_{-k} = \rho_k^*$ , can approximate any **continuous** PSD for **sufficiently large**  $m, n$

Spectral factorization theorem says we can write a rational  $\phi(\omega)$  as

$$\phi(\omega) = \left| \frac{B(\omega)}{A(\omega)} \right|^2 \sigma^2$$

for some positive scalar  $\sigma^2$ , where


$$A(z) = 1 + a_1 z^{-1} + \dots + a_n z^{-n}$$

$$B(z) = 1 + b_1 z^{-1} + \dots + b_m z^{-m}$$

and  $A(\omega) = A(z)|_{z=e^{i\omega}}$ .

# Signal modeling interpretation


White noise  $e(t)$  with variance  $\sigma^2$  filtered through a linear filter:



►  $y(t) = \frac{B(z)}{A(z)} e(t).$

# Signal modeling interpretation


White noise  $e(t)$  with variance  $\sigma^2$  filtered through a linear filter:



- ▶  $y(t) = \frac{B(z)}{A(z)}e(t).$
- ▶ **Input spectrum:**  $\phi_e(\omega) =$

# Signal modeling interpretation

White noise  $e(t)$  with variance  $\sigma^2$  filtered through a linear filter:




- ▶  $y(t) = \frac{B(z)}{A(z)}e(t).$
- ▶ **Input spectrum:**  $\phi_e(\omega) = \sigma^2.$



# Signal modeling interpretation

White noise  $e(t)$  with variance  $\sigma^2$  filtered through a linear filter:




- ▶  $y(t) = \frac{B(z)}{A(z)}e(t).$
- ▶ **Input spectrum:**  $\phi_e(\omega) = \sigma^2.$
- ▶ **Output spectrum:**

$$\phi_y(\omega) =$$

# Signal modeling interpretation

White noise  $e(t)$  with variance  $\sigma^2$  filtered through a linear filter:




- ▶  $y(t) = \frac{B(z)}{A(z)}e(t).$
- ▶ **Input spectrum:**  $\phi_e(\omega) = \sigma^2.$
- ▶ **Output spectrum:**

$$\phi_y(\omega) = \left| \frac{B(\omega)}{A(\omega)} \right|^2 \sigma^2$$

# Signal modeling interpretation

White noise  $e(t)$  with variance  $\sigma^2$  filtered through a linear filter:



- ▶  $y(t) = \frac{B(z)}{A(z)}e(t)$ .
- ▶ **Input spectrum:**  $\phi_e(\omega) = \sigma^2$ .
- ▶ **Output spectrum:**

$$\phi_y(\omega) = \left| \frac{B(\omega)}{A(\omega)} \right|^2 \sigma^2$$

- ▶ **Conclusion:** A rational spectrum can be modelled as the spectrum of white noise filtered through a linear filter.

# Poles, zeros and the spectrum

---

**Linear filter:**

$$y(t) = \frac{B(z)}{A(z)}e(t) = \sum_{k=0}^{\infty} h_k e(t-k), \quad h_0 = 1 \text{ in these slides}$$

# Poles, zeros and the spectrum

---

**Linear filter:**

$$y(t) = \frac{B(z)}{A(z)}e(t) = \sum_{k=0}^{\infty} h_k e(t-k), \quad h_0 = 1 \text{ in these slides}$$

► **Poles:** Zeros of  $A(z)$ . **Zeros:** Zeros of  $B(z)$ .

# Poles, zeros and the spectrum

---

## Linear filter:

$$y(t) = \frac{B(z)}{A(z)}e(t) = \sum_{k=0}^{\infty} h_k e(t-k), \quad h_0 = 1 \text{ in these slides}$$

- ▶ **Poles:** Zeros of  $A(z)$ . **Zeros:** Zeros of  $B(z)$ .
- ▶ **Stable** if all poles lies inside the unit circle.

# Poles, zeros and the spectrum

## Linear filter:

$$y(t) = \frac{B(z)}{A(z)}e(t) = \sum_{k=0}^{\infty} h_k e(t-k), \quad h_0 = 1 \text{ in these slides}$$

- **Poles:** Zeros of  $A(z)$ . **Zeros:** Zeros of  $B(z)$ .
- **Stable** if all poles lies inside the unit circle.

## Spectrum:

$$\phi_y(z) = \frac{B(z)B^*(1/z^*)}{A(z)A^*(1/z^*)}\sigma^2 \xrightarrow{z=e^{i\omega}} \phi_y(\omega) = \left| \frac{B(\omega)}{A(\omega)} \right|^2 \sigma^2$$

# Poles, zeros and the spectrum

## Linear filter:

$$y(t) = \frac{B(z)}{A(z)}e(t) = \sum_{k=0}^{\infty} h_k e(t-k), \quad h_0 = 1 \text{ in these slides}$$

- **Poles:** Zeros of  $A(z)$ . **Zeros:** Zeros of  $B(z)$ .
- **Stable** if all poles lies inside the unit circle.

## Spectrum:

$$\phi_y(z) = \frac{B(z)B^*(1/z^*)}{A(z)A^*(1/z^*)}\sigma^2 \xrightarrow{z=e^{i\omega}} \phi_y(\omega) = \left| \frac{B(\omega)}{A(\omega)} \right|^2 \sigma^2$$

- $\phi(\omega)$  is finite for all  $\omega \iff$



# Poles, zeros and the spectrum

## Linear filter:

$$y(t) = \frac{B(z)}{A(z)}e(t) = \sum_{k=0}^{\infty} h_k e(t-k), \quad h_0 = 1 \text{ in these slides}$$

- **Poles:** Zeros of  $A(z)$ . **Zeros:** Zeros of  $B(z)$ .
- **Stable** if all poles lies inside the unit circle.

## Spectrum:

$$\phi_y(z) = \frac{B(z)B^*(1/z^*)}{A(z)A^*(1/z^*)}\sigma^2 \xrightarrow{z=e^{i\omega}} \phi_y(\omega) = \left| \frac{B(\omega)}{A(\omega)} \right|^2 \sigma^2$$

- $\phi(\omega)$  is finite for all  $\omega \iff$  No poles on the unit circle.

# Poles, zeros and the spectrum

## Linear filter:

$$y(t) = \frac{B(z)}{A(z)} e(t) = \sum_{k=0}^{\infty} h_k e(t-k), \quad h_0 = 1 \text{ in these slides}$$

- ▶ **Poles:** Zeros of  $A(z)$ . **Zeros:** Zeros of  $B(z)$ .
- ▶ **Stable** if all poles lies inside the unit circle.

## Spectrum:

$$\phi_y(z) = \frac{B(z)B^*(1/z^*)}{A(z)A^*(1/z^*)} \sigma^2 \xrightarrow{z=e^{i\omega}} \phi_y(\omega) = \left| \frac{B(\omega)}{A(\omega)} \right|^2 \sigma^2$$

- ▶  $\phi(\omega)$  is finite for all  $\omega \iff$  No poles on the unit circle.
- ▶ Can always choose  $A(z)$  and  $B(z)$  so that the poles and zeros lie inside the unit circle. (If  $\phi(\omega)$  is finite)

# Poles, zeros and the spectrum

## Linear filter:

$$y(t) = \frac{B(z)}{A(z)} e(t) = \sum_{k=0}^{\infty} h_k e(t-k), \quad h_0 = 1 \text{ in these slides}$$

- ▶ **Poles:** Zeros of  $A(z)$ . **Zeros:** Zeros of  $B(z)$ .
- ▶ **Stable** if all poles lies inside the unit circle.

## Spectrum:

$$\phi_y(z) = \frac{B(z)B^*(1/z^*)}{A(z)A^*(1/z^*)} \sigma^2 \xrightarrow{z=e^{i\omega}} \phi_y(\omega) = \left| \frac{B(\omega)}{A(\omega)} \right|^2 \sigma^2$$

- ▶  $\phi(\omega)$  is finite for all  $\omega \iff$  No poles on the unit circle.
- ▶ Can always choose  $A(z)$  and  $B(z)$  so that the poles and zeros lie inside the unit circle. (If  $\phi(\omega)$  is finite)
- ▶ Pole at  $z_1 \approx e^{i\omega_1}$  (near unit circle)

# Poles, zeros and the spectrum

## Linear filter:

$$y(t) = \frac{B(z)}{A(z)} e(t) = \sum_{k=0}^{\infty} h_k e(t-k), \quad h_0 = 1 \text{ in these slides}$$

- ▶ **Poles:** Zeros of  $A(z)$ . **Zeros:** Zeros of  $B(z)$ .
- ▶ **Stable** if all poles lies inside the unit circle.

## Spectrum:

$$\phi_y(z) = \frac{B(z)B^*(1/z^*)}{A(z)A^*(1/z^*)} \sigma^2 \xrightarrow{z=e^{i\omega}} \phi_y(\omega) = \left| \frac{B(\omega)}{A(\omega)} \right|^2 \sigma^2$$

- ▶  $\phi(\omega)$  is finite for all  $\omega \iff$  No poles on the unit circle.
- ▶ Can always choose  $A(z)$  and  $B(z)$  so that the poles and zeros lie inside the unit circle. (If  $\phi(\omega)$  is finite)
- ▶ Pole at  $z_1 \approx e^{i\omega_1}$  (near unit circle)  $\implies \phi(\omega_1)$  large.

# Poles, zeros and the spectrum

## Linear filter:

$$y(t) = \frac{B(z)}{A(z)} e(t) = \sum_{k=0}^{\infty} h_k e(t-k), \quad h_0 = 1 \text{ in these slides}$$



- **Poles:** Zeros of  $A(z)$ . **Zeros:** Zeros of  $B(z)$ .
- **Stable** if all poles lies inside the unit circle.

## Spectrum:

$$\phi_y(z) = \frac{B(z)B^*(1/z^*)}{A(z)A^*(1/z^*)} \sigma^2 \xrightarrow{z=e^{i\omega}} \phi_y(\omega) = \left| \frac{B(\omega)}{A(\omega)} \right|^2 \sigma^2$$

- $\phi(\omega)$  is finite for all  $\omega \iff$  No poles on the unit circle.
- Can always choose  $A(z)$  and  $B(z)$  so that the poles and zeros lie inside the unit circle. (If  $\phi(\omega)$  is finite)
- Pole at  $z_1 \approx e^{i\omega_1}$  (near unit circle)  $\implies \phi(\omega_1)$  large.
- Zero at  $z_1 \approx e^{i\omega_1}$  (near unit circle)  $\implies \phi(\omega_1)$  small.

# Example



# AR(MA) (1)

## Autoregressive moving average (pole-zero model)

$$\text{ARMA: } A(z)y(t) = B(z)e(t)$$

Special cases

$$\text{AR: } A(z)y(t) = e(t)$$

$$\text{MA: } y(t) = B(z)e(t)$$

Estimate parameters:  $\{a_i\}_{i=1}^n, \{b_i\}_{i=1}^m, \sigma^2 \implies$  PSD estimate

# AR(MA) (1)

## Autoregressive moving average (pole-zero model)

$$\text{ARMA: } A(z)y(t) = B(z)e(t)$$



Special cases

$$\text{AR: } A(z)y(t) = e(t)$$

$$\text{MA: } y(t) = B(z)e(t)$$

Estimate parameters:  $\{a_i\}_{i=1}^n, \{b_i\}_{i=1}^m, \sigma^2 \implies$  **PSD estimate**

**AR Ex.:** `N=256; e=randn(N,1); y=filter(1,[1 -0.95],e);`






# AR(MA) (2)

```
N=256; e=randn(N,1); y=filter(1,[1 -0.95],e);
```

Single real-valued pole ( $A(z) = 0$ ) at  $z = 0.95$ , i.e. close to unit circle at  $\omega = 0$



# Covariance structure

---

ARMA can be written out as (with  $b_0 = 1$ )

$$y(t) + \sum_{i=1}^n a_i y(t-i) = \sum_{j=0}^m b_j e(t-j)$$

# Covariance structure

---

ARMA can be written out as (with  $b_0 = 1$ )

$$y(t) + \sum_{i=1}^n a_i y(t-i) = \sum_{j=0}^m b_j e(t-j)$$

Multiplying by  $y^*(t-k)$  and taking the expectation gives

# Covariance structure

---

ARMA can be written out as (with  $b_0 = 1$ )

$$y(t) + \sum_{i=1}^n a_i y(t-i) = \sum_{j=0}^m b_j e(t-j)$$

Multiplying by  $y^*(t-k)$  and taking the expectation gives

$$r(k) + \sum_{i=1}^n a_i r(k-i) = \sigma^2 \sum_{j=0}^m b_j h_{j-k}^*$$

# Covariance structure

---

ARMA can be written out as (with  $b_0 = 1$ )

$$y(t) + \sum_{i=1}^n a_i y(t-i) = \sum_{j=0}^m b_j e(t-j)$$

Multiplying by  $y^*(t-k)$  and taking the expectation gives

$$r(k) + \sum_{i=1}^n a_i r(k-i) = \sigma^2 \sum_{j=0}^m b_j h_{j-k}^*$$

But  $h_s = 0$  for  $s < 0$  (causality), which gives

# Covariance structure

ARMA can be written out as (with  $b_0 = 1$ )

$$y(t) + \sum_{i=1}^n a_i y(t-i) = \sum_{j=0}^m b_j e(t-j)$$

Multiplying by  $y^*(t-k)$  and taking the expectation gives

$$r(k) + \sum_{i=1}^n a_i r(k-i) = \sigma^2 \sum_{j=0}^m b_j h_{j-k}^*$$

But  $h_s = 0$  for  $s < 0$  (causality), which gives

## Covariance structure

$$r(k) + \sum_{i=1}^n a_i r(k-i) = 0, \quad \text{for } k > m$$



# Methods for AR-models

# AR: Yule-Walker equations

---

**AR-model:**  $A(z)y(t) = e(t)$ . Here  $m = 0$ , and  $B(z) = b_0 = 1$ , so

$$r(0) + \sum_{i=1}^n a_i r(-i) =$$



# AR: Yule-Walker equations

---

**AR-model:**  $A(z)y(t) = e(t)$ . Here  $m = 0$ , and  $B(z) = b_0 = 1$ , so

$$r(0) + \sum_{i=1}^n a_i r(-i) = \sigma^2 b_0 h_0 =$$

# AR: Yule-Walker equations

---

**AR-model:**  $A(z)y(t) = e(t)$ . Here  $m = 0$ , and  $B(z) = b_0 = 1$ , so

$$r(0) + \sum_{i=1}^n a_i r(-i) = \sigma^2 b_0 h_0 = \sigma^2,$$

# AR: Yule-Walker equations

---

**AR-model:**  $A(z)y(t) = e(t)$ . Here  $m = 0$ , and  $B(z) = b_0 = 1$ , so

$$r(0) + \sum_{i=1}^n a_i r(-i) = \sigma^2 b_0 h_0 = \sigma^2,$$

$$r(k) + \sum_{i=1}^n a_i r(k-i) = 0, \quad k = 1, \dots, n$$

These are called the **Yule-Walker equations**.

# AR: Yule-Walker equations

**AR-model:**  $A(z)y(t) = e(t)$ . Here  $m = 0$ , and  $B(z) = b_0 = 1$ , so

$$r(0) + \sum_{i=1}^n a_i r(-i) = \sigma^2 b_0 h_0 = \sigma^2,$$

$$r(k) + \sum_{i=1}^n a_i r(k-i) = 0, \quad k = 1, \dots, n$$

These are called the **Yule-Walker equations**.

On matrix form:

$$\begin{bmatrix} r(0) & r(-1) & \cdots & r(-n) \\ r(1) & r(0) & \cdots & r(-n+1) \\ \vdots & & \ddots & \vdots \\ r(n) & r(n-1) & \cdots & r(0) \end{bmatrix} \begin{bmatrix} 1 \\ a_1 \\ \vdots \\ a_n \end{bmatrix} = \begin{bmatrix} \sigma^2 \\ 0 \\ \vdots \\ 0 \end{bmatrix}$$

# Yule-Walker method

---

- ▶ Want to find an estimate of  $\theta = \begin{bmatrix} a_1 & \cdots & a_n \end{bmatrix}^\top$ .

# Yule-Walker method

---

- ▶ Want to find an estimate of  $\theta = \begin{bmatrix} a_1 & \cdots & a_n \end{bmatrix}^\top$ .
- ▶ **Idea:** Replace all  $r(k)$  with **standard biased estimate**  $\hat{r}(k)$ .

# Yule-Walker method

- ▶ Want to find an estimate of  $\theta = [a_1 \ \cdots \ a_n]^\top$ .
- ▶ **Idea:** Replace all  $r(k)$  with **standard biased estimate**  $\hat{r}(k)$ .
- ▶ If we remove the first row we get

$$\underbrace{\begin{bmatrix} \hat{r}(1) \\ \vdots \\ \hat{r}(n) \end{bmatrix}}_{\hat{r}_n} + \underbrace{\begin{bmatrix} \hat{r}(0) & \cdots & \hat{r}(-n+1) \\ \vdots & \ddots & \vdots \\ \hat{r}(n-1) & \cdots & \hat{r}(0) \end{bmatrix}}_{\hat{R}_n} \underbrace{\begin{bmatrix} \hat{a}_1 \\ \vdots \\ \hat{a}_n \end{bmatrix}}_{\hat{\theta}} = \begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix}$$

# Yule-Walker method

- ▶ Want to find an estimate of  $\theta = [a_1 \ \cdots \ a_n]^\top$ .
- ▶ **Idea:** Replace all  $r(k)$  with **standard biased estimate**  $\hat{r}(k)$ .
- ▶ If we remove the first row we get

$$\underbrace{\begin{bmatrix} \hat{r}(1) \\ \vdots \\ \hat{r}(n) \end{bmatrix}}_{\hat{r}_n} + \underbrace{\begin{bmatrix} \hat{r}(0) & \cdots & \hat{r}(-n+1) \\ \vdots & \ddots & \vdots \\ \hat{r}(n-1) & \cdots & \hat{r}(0) \end{bmatrix}}_{\hat{R}_n} \underbrace{\begin{bmatrix} \hat{a}_1 \\ \vdots \\ \hat{a}_n \end{bmatrix}}_{\hat{\theta}} = \begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix}$$

- ▶ **Solution:**  $\hat{\theta} =$



# Yule-Walker method

- ▶ Want to find an estimate of  $\theta = [a_1 \ \cdots \ a_n]^\top$ .
- ▶ **Idea:** Replace all  $r(k)$  with **standard biased estimate**  $\hat{r}(k)$ .
- ▶ If we remove the first row we get

$$\underbrace{\begin{bmatrix} \hat{r}(1) \\ \vdots \\ \hat{r}(n) \end{bmatrix}}_{\hat{r}_n} + \underbrace{\begin{bmatrix} \hat{r}(0) & \cdots & \hat{r}(-n+1) \\ \vdots & \ddots & \vdots \\ \hat{r}(n-1) & \cdots & \hat{r}(0) \end{bmatrix}}_{\hat{R}_n} \underbrace{\begin{bmatrix} \hat{a}_1 \\ \vdots \\ \hat{a}_n \end{bmatrix}}_{\hat{\theta}} = \begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix}$$

- ▶ **Solution:**  $\hat{\theta} = -\hat{R}_n^{-1}\hat{r}_n$ .

# Yule-Walker method

- ▶ Want to find an estimate of  $\theta = [a_1 \ \cdots \ a_n]^\top$ .
- ▶ **Idea:** Replace all  $r(k)$  with **standard biased estimate**  $\hat{r}(k)$ .
- ▶ If we remove the first row we get

$$\underbrace{\begin{bmatrix} \hat{r}(1) \\ \vdots \\ \hat{r}(n) \end{bmatrix}}_{\hat{r}_n} + \underbrace{\begin{bmatrix} \hat{r}(0) & \cdots & \hat{r}(-n+1) \\ \vdots & \ddots & \vdots \\ \hat{r}(n-1) & \cdots & \hat{r}(0) \end{bmatrix}}_{\hat{R}_n} \underbrace{\begin{bmatrix} \hat{a}_1 \\ \vdots \\ \hat{a}_n \end{bmatrix}}_{\hat{\theta}} = \begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix}$$

- ▶ **Solution:**  $\hat{\theta} = -\hat{R}_n^{-1}\hat{r}_n$ .
- ▶ Using  $\hat{\theta}$ , the first row now gives

$$\hat{\sigma}^2 =$$

# Yule-Walker method

- ▶ Want to find an estimate of  $\theta = [a_1 \ \cdots \ a_n]^\top$ .
- ▶ **Idea:** Replace all  $r(k)$  with **standard biased estimate**  $\hat{r}(k)$ .
- ▶ If we remove the first row we get

$$\underbrace{\begin{bmatrix} \hat{r}(1) \\ \vdots \\ \hat{r}(n) \end{bmatrix}}_{\hat{r}_n} + \underbrace{\begin{bmatrix} \hat{r}(0) & \cdots & \hat{r}(-n+1) \\ \vdots & \ddots & \vdots \\ \hat{r}(n-1) & \cdots & \hat{r}(0) \end{bmatrix}}_{\hat{R}_n} \underbrace{\begin{bmatrix} \hat{a}_1 \\ \vdots \\ \hat{a}_n \end{bmatrix}}_{\hat{\theta}} = \begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix}$$

- ▶ **Solution:**  $\hat{\theta} = -\hat{R}_n^{-1}\hat{r}_n$ .
- ▶ Using  $\hat{\theta}$ , the first row now gives

$$\hat{\sigma}^2 = \hat{r}(0) + [\hat{r}(-1) \ \cdots \ \hat{r}(-n)] \hat{\theta}.$$

# AR as linear regression

---

The AR-model  $A(z)y(t) = e(t)$  can be written as

$$e(t) = y(t) + \sum_{i=1}^n a_i y(t-i) =$$

# AR as linear regression

---

The AR-model  $A(z)y(t) = e(t)$  can be written as

$$e(t) = y(t) + \sum_{i=1}^n a_i y(t-i) = y(t) - \varphi^\top(t)\theta,$$

where

$$\varphi(t) = \begin{bmatrix} -y(t-1) \\ \vdots \\ -y(t-n) \end{bmatrix}, \quad \theta = \begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix}.$$

# AR as linear regression

---

The AR-model  $A(z)y(t) = e(t)$  can be written as

$$e(t) = y(t) + \sum_{i=1}^n a_i y(t-i) = y(t) - \varphi^\top(t)\theta,$$

where

$$\varphi(t) = \begin{bmatrix} -y(t-1) \\ \vdots \\ -y(t-n) \end{bmatrix}, \quad \theta = \begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix}.$$

- **Prediction error:** If we predict  $y(t)$  using  $\hat{y}(t) = \varphi^\top(t)\theta$ , then  $e(t) = y(t) - \hat{y}(t)$  is the prediction error.

# AR as linear regression

The AR-model  $A(z)y(t) = e(t)$  can be written as

$$e(t) = y(t) + \sum_{i=1}^n a_i y(t-i) = y(t) - \varphi^\top(t)\theta,$$

where

$$\varphi(t) = \begin{bmatrix} -y(t-1) \\ \vdots \\ -y(t-n) \end{bmatrix}, \quad \theta = \begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix}.$$

- **Prediction error:** If we predict  $y(t)$  using  $\hat{y}(t) = \varphi^\top(t)\theta$ , then  $e(t) = y(t) - \hat{y}(t)$  is the prediction error.
- **Note:** In the book  $\varphi^\top(t)$  is defined without the minus sign, so we have  $e(t) = y(t) + \varphi^\top(t)\theta$ .

# AR: The least squares method

---

- **Idea:** Estimate  $\theta$  by minimizing the prediction errors

$$\hat{\theta} = \arg \min_{\theta} \sum_{t=1}^N |e(t)|^2 =$$



# AR: The least squares method

---

- **Idea:** Estimate  $\theta$  by minimizing the prediction errors

$$\begin{aligned}\hat{\theta} &= \arg \min_{\theta} \sum_{t=1}^N |e(t)|^2 = \arg \min_{\theta} \sum_{t=1}^N |y - \varphi^{\top}(t)\theta|^2 \\ &= \end{aligned}$$

# AR: The least squares method

- **Idea:** Estimate  $\theta$  by minimizing the prediction errors

$$\begin{aligned}\hat{\theta} &= \arg \min_{\theta} \sum_{t=1}^N |e(t)|^2 = \arg \min_{\theta} \sum_{t=1}^N |y - \varphi^{\top}(t)\theta|^2 \\ &= \arg \min_{\theta} \|y - \Phi\theta\|_2^2,\end{aligned}$$

where

$$y = \begin{bmatrix} y(1) \\ \vdots \\ y(N) \end{bmatrix}, \quad \Phi = \begin{bmatrix} \varphi^{\top}(1) \\ \vdots \\ \varphi^{\top}(N) \end{bmatrix}.$$

# AR: The least squares method

- **Idea:** Estimate  $\theta$  by minimizing the prediction errors

$$\begin{aligned}\hat{\theta} &= \arg \min_{\theta} \sum_{t=1}^N |e(t)|^2 = \arg \min_{\theta} \sum_{t=1}^N |y - \varphi^{\top}(t)\theta|^2 \\ &= \arg \min_{\theta} \|y - \Phi\theta\|_2^2,\end{aligned}$$

where

$$y = \begin{bmatrix} y(1) \\ \vdots \\ y(N) \end{bmatrix}, \quad \Phi = \begin{bmatrix} \varphi^{\top}(1) \\ \vdots \\ \varphi^{\top}(N) \end{bmatrix}.$$

- **Solution:**

$$\hat{\theta} =$$

# AR: The least squares method

- **Idea:** Estimate  $\theta$  by minimizing the prediction errors

$$\begin{aligned}\hat{\theta} &= \arg \min_{\theta} \sum_{t=1}^N |e(t)|^2 = \arg \min_{\theta} \sum_{t=1}^N |y - \varphi^{\top}(t)\theta|^2 \\ &= \arg \min_{\theta} \|y - \Phi\theta\|_2^2,\end{aligned}$$

where

$$y = \begin{bmatrix} y(1) \\ \vdots \\ y(N) \end{bmatrix}, \quad \Phi = \begin{bmatrix} \varphi^{\top}(1) \\ \vdots \\ \varphi^{\top}(N) \end{bmatrix}.$$

- **Solution:**

$$\hat{\theta} = (\Phi^* \Phi)^{-1} \Phi^* y.$$

# AR: The least squares method

- **Idea:** Estimate  $\theta$  by minimizing the prediction errors

$$\begin{aligned}\hat{\theta} &= \arg \min_{\theta} \sum_{t=1}^N |e(t)|^2 = \arg \min_{\theta} \sum_{t=1}^N |y - \varphi^{\top}(t)\theta|^2 \\ &= \arg \min_{\theta} \|y - \Phi\theta\|_2^2,\end{aligned}$$

where

$$y = \begin{bmatrix} y(1) \\ \vdots \\ y(N) \end{bmatrix}, \quad \Phi = \begin{bmatrix} \varphi^{\top}(1) \\ \vdots \\ \varphi^{\top}(N) \end{bmatrix}.$$

- **Solution:**

$$\hat{\theta} = (\Phi^* \Phi)^{-1} \Phi^* y.$$

- Again, the book uses  $Y = -\Phi$ , so  $\hat{\theta} = -(Y^* Y)^{-1} Y^* y$ .



# What about negative $t$ ?

---

# What about negative $t$ ?

---

- If, for example,  $n = 3$  then

$$\varphi^{\top}(1) = \begin{bmatrix} y(0) & y(-1) & y(-2) \end{bmatrix}$$

$$\varphi^{\top}(2) = \begin{bmatrix} y(1) & y(0) & y(-1) \end{bmatrix}$$

$$\varphi^{\top}(3) = \begin{bmatrix} y(2) & y(1) & y(0) \end{bmatrix}$$

# What about negative $t$ ?

---

- ▶ If, for example,  $n = 3$  then

$$\varphi^\top(1) = \begin{bmatrix} y(0) & y(-1) & y(-2) \end{bmatrix}$$

$$\varphi^\top(2) = \begin{bmatrix} y(1) & y(0) & y(-1) \end{bmatrix}$$

$$\varphi^\top(3) = \begin{bmatrix} y(2) & y(1) & y(0) \end{bmatrix}$$

- ▶ We have only measured  $y(t)$  for  $t \geq 1$ . How to handle this?



# What about negative $t$ ?

---

- ▶ If, for example,  $n = 3$  then

$$\varphi^\top(1) = \begin{bmatrix} y(0) & y(-1) & y(-2) \end{bmatrix}$$

$$\varphi^\top(2) = \begin{bmatrix} y(1) & y(0) & y(-1) \end{bmatrix}$$

$$\varphi^\top(3) = \begin{bmatrix} y(2) & y(1) & y(0) \end{bmatrix}$$

- ▶ We have only measured  $y(t)$  for  $t \geq 1$ . How to handle this?
- ▶ **Option 1:** Let  $y(t) = 0$  for  $t \leq 0$ .

# What about negative $t$ ?

---

- ▶ If, for example,  $n = 3$  then

$$\varphi^\top(1) = \begin{bmatrix} y(0) & y(-1) & y(-2) \end{bmatrix}$$

$$\varphi^\top(2) = \begin{bmatrix} y(1) & y(0) & y(-1) \end{bmatrix}$$

$$\varphi^\top(3) = \begin{bmatrix} y(2) & y(1) & y(0) \end{bmatrix}$$

- ▶ We have only measured  $y(t)$  for  $t \geq 1$ . How to handle this?
- ▶ **Option 1:** Let  $y(t) = 0$  for  $t \leq 0$ .
- ▶ **Option 2:** Skip the first  $n$  rows in  $Y$  and  $\Phi$ .

# What about negative $t$ ?

---

- ▶ If, for example,  $n = 3$  then

$$\varphi^\top(1) = \begin{bmatrix} y(0) & y(-1) & y(-2) \end{bmatrix}$$

$$\varphi^\top(2) = \begin{bmatrix} y(1) & y(0) & y(-1) \end{bmatrix}$$

$$\varphi^\top(3) = \begin{bmatrix} y(2) & y(1) & y(0) \end{bmatrix}$$

- ▶ We have only measured  $y(t)$  for  $t \geq 1$ . How to handle this?
- ▶ **Option 1:** Let  $y(t) = 0$  for  $t \leq 0$ .
- ▶ **Option 2:** Skip the first  $n$  rows in  $Y$  and  $\Phi$ .
- ▶ Could also add rows corresponding to e.g.  
 $t = N + 1, \dots, N + n$  and assume that  $y(t) = 0$  at these rows.

# AR: The least squares method

$$y = \begin{bmatrix} y(1) \\ y(2) \\ \vdots \\ y(n+1) \\ y(n+2) \\ \vdots \\ y(N) \\ 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \quad Y = \begin{bmatrix} 0 & 0 & \dots & 0 \\ y(1) & 0 & & \vdots \\ \vdots & \ddots & \ddots & 0 \\ y(n) & y(n-1) & \dots & y(1) \\ y(n+1) & y(n) & \dots & y(2) \\ \vdots & \vdots & & \vdots \\ y(N-1) & y(N-2) & \dots & y(N-n) \\ y(N) & y(N-1) & \dots & y(N-n+1) \\ 0 & y(N) & & \\ \vdots & \ddots & \ddots & \vdots \\ 0 & \dots & 0 & y(N) \end{bmatrix}$$

**Note:** With notation used in these slides  $\Phi = -Y$ .

# AR: The least squares method

$$y = \begin{bmatrix} y(1) \\ y(2) \\ \vdots \\ y(n+1) \\ y(n+2) \\ \vdots \\ y(N) \\ 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \quad Y = \begin{bmatrix} 0 & 0 & \dots & 0 \\ y(1) & 0 & & \vdots \\ \vdots & \ddots & \ddots & 0 \\ y(n) & y(n-1) & \dots & y(1) \\ y(n+1) & y(n) & \dots & y(2) \\ \vdots & \vdots & & \vdots \\ y(N-1) & y(N-2) & \dots & y(N-n) \\ y(N) & y(N-1) & \dots & y(N-n+1) \\ 0 & y(N) & & \\ \vdots & \ddots & \ddots & \vdots \\ 0 & \dots & 0 & y(N) \end{bmatrix}$$

**Note:** With notation used in these slides  $\Phi = -Y$ .

► **Autocorrelation method:** Use full matrices. Equiv. to YW.

# AR: The least squares method

$$y = \begin{bmatrix} y(1) \\ y(2) \\ \vdots \\ y(n+1) \\ y(n+2) \\ \vdots \\ y(N) \\ 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \quad Y = \begin{bmatrix} 0 & 0 & \dots & 0 \\ y(1) & 0 & & \vdots \\ \vdots & \ddots & \ddots & 0 \\ y(n) & y(n-1) & \dots & y(1) \\ y(n+1) & y(n) & \dots & y(2) \\ \vdots & \vdots & & \vdots \\ y(N-1) & y(N-2) & \dots & y(N-n) \\ y(N) & y(N-1) & \dots & y(N-n+1) \\ 0 & y(N) & & \\ \vdots & \ddots & \ddots & \vdots \\ 0 & \dots & 0 & y(N) \end{bmatrix}$$

**Note:** With notation used in these slides  $\Phi = -Y$ .

- **Autocorrelation method:** Use full matrices. Equiv. to YW.
- **LS method:** Only keep middle part. Approx YW.



# Methods for ARMA-models

# ARMA-models

---

- ▶ Sharp peaks captured by AR-part, and deep valleys by MA-part:

$$A(q)y(t) = B(q)e(t).$$



# ARMA-models

---

- ▶ Sharp peaks captured by AR-part, and deep valleys by MA-part:

$$A(q)y(t) = B(q)e(t).$$

- ▶ Much harder to estimate an ARMA-model!

# ARMA-models

---

- ▶ Sharp peaks captured by AR-part, and deep valleys by MA-part:

$$A(q)y(t) = B(q)e(t).$$

- ▶ Much harder to estimate an ARMA-model!
- ▶ Here two methods that often works will be discussed.

# ARMA-models

---

- ▶ Sharp peaks captured by AR-part, and deep valleys by MA-part:

$$A(q)y(t) = B(q)e(t).$$

- ▶ Much harder to estimate an ARMA-model!
- ▶ Here two methods that often works will be discussed.
- ▶ Many more estimators in the literature.

# The two-stage LS method

---

We can write the ARMA-model as

$$e(t) = y(t) - \varphi^{\top}(t)\theta$$

with

$$\varphi(t) =$$

$$\theta =$$

# The two-stage LS method

---

We can write the ARMA-model as

$$e(t) = y(t) - \varphi^\top(t)\theta$$

with

$$\varphi(t) = \begin{bmatrix} -y(t-1) \\ \vdots \\ -y(t-n) \\ -e(t-1) \\ \vdots \\ -e(t-m) \end{bmatrix}, \quad \theta = \begin{bmatrix} a_1 \\ \vdots \\ a_n \\ b_1 \\ \vdots \\ b_m \end{bmatrix}.$$

# The two-stage LS method

We can write the ARMA-model as

$$e(t) = y(t) - \varphi^\top(t)\theta$$

with

$$\varphi(t) = \begin{bmatrix} -y(t-1) \\ \vdots \\ -y(t-n) \\ -e(t-1) \\ \vdots \\ -e(t-m) \end{bmatrix}, \quad \theta = \begin{bmatrix} a_1 \\ \vdots \\ a_n \\ b_1 \\ \vdots \\ b_m \end{bmatrix}.$$

► **Idea:** Minimize  $\sum_{t=1}^N |e(t)|^2 = \|y - \Phi\theta\|_2^2$ .

# The two-stage LS method

We can write the ARMA-model as

$$e(t) = y(t) - \varphi^\top(t)\theta$$

with

$$\varphi(t) = \begin{bmatrix} -y(t-1) \\ \vdots \\ -y(t-n) \\ -e(t-1) \\ \vdots \\ -e(t-m) \end{bmatrix}, \quad \theta = \begin{bmatrix} a_1 \\ \vdots \\ a_n \\ b_1 \\ \vdots \\ b_m \end{bmatrix}.$$

- ▶ **Idea:** Minimize  $\sum_{t=1}^N |e(t)|^2 = \|y - \Phi\theta\|_2^2$ .
- ▶ **Problem:** Can't create  $\Phi$  since we do not have  $e(t)$ .

# The two-stage LS method

We can write the ARMA-model as

$$e(t) = y(t) - \varphi^\top(t)\theta$$

with

$$\varphi(t) = \begin{bmatrix} -y(t-1) \\ \vdots \\ -y(t-n) \\ -e(t-1) \\ \vdots \\ -e(t-m) \end{bmatrix}, \quad \theta = \begin{bmatrix} a_1 \\ \vdots \\ a_n \\ b_1 \\ \vdots \\ b_m \end{bmatrix}.$$

- ▶ **Idea:** Minimize  $\sum_{t=1}^N |e(t)|^2 = \|y - \Phi\theta\|_2^2$ .
- ▶ **Problem:** Can't create  $\Phi$  since we do not have  $e(t)$ .
- ▶ **Potential solution:** First estimate  $e(t)$ , then use LS to estimate  $\theta$ .



# The two-stage LS method

---

- ▶ We can assume that the ARMA-model is minimum phase.
- ▶ Then the ARMA-model can be written as an infinite dimensional AR-model

$$e(t) = \frac{A(q)}{B(q)}y(t) = y(t) + \alpha_1 y(t-1) + \alpha_2 y(t-2) + \dots,$$

where  $\alpha_k \rightarrow 0$  as  $k \rightarrow \infty$ .

# The two-stage LS method

---

- ▶ We can assume that the ARMA-model is minimum phase.
- ▶ Then the ARMA-model can be written as an infinite dimensional AR-model

$$e(t) = \frac{A(q)}{B(q)}y(t) = y(t) + \alpha_1 y(t-1) + \alpha_2 y(t-2) + \dots,$$

where  $\alpha_k \rightarrow 0$  as  $k \rightarrow \infty$ .

- ▶ For large enough  $K$  we should be able to approximate the ARMA-model with an AR-model of order  $K$ .

# The two-stage LS method

---

- ▶ We can assume that the ARMA-model is minimum phase.
- ▶ Then the ARMA-model can be written as an infinite dimensional AR-model

$$e(t) = \frac{A(q)}{B(q)}y(t) = y(t) + \alpha_1 y(t-1) + \alpha_2 y(t-2) + \dots,$$

where  $\alpha_k \rightarrow 0$  as  $k \rightarrow \infty$ .

- ▶ For large enough  $K$  we should be able to approximate the ARMA-model with an AR-model of order  $K$ .
- ▶ **Potential problems:**
  - ▶  $K$  should not be too large with respect to  $N$  (overfitting).

# The two-stage LS method

---

- ▶ We can assume that the ARMA-model is minimum phase.
- ▶ Then the ARMA-model can be written as an infinite dimensional AR-model

$$e(t) = \frac{A(q)}{B(q)}y(t) = y(t) + \alpha_1 y(t-1) + \alpha_2 y(t-2) + \dots,$$

where  $\alpha_k \rightarrow 0$  as  $k \rightarrow \infty$ .

- ▶ For large enough  $K$  we should be able to approximate the ARMA-model with an AR-model of order  $K$ .
- ▶ **Potential problems:**
  - ▶  $K$  should not be too large with respect to  $N$  (overfitting).
  - ▶ With **zeros close to the unit circle** (deep valleys in the PSD),  $K$  must be very large to keep the bias errors low.

# The two-stage LS method

---

- **Step 1:** Choose a sufficiently large  $K$ , and estimate an AR-model of order  $K$ . Let

$$\hat{e}(t) = y(t) + \sum_{k=1}^K \hat{\alpha}_k y(t - k)$$

# The two-stage LS method

---

- ▶ **Step 1:** Choose a sufficiently large  $K$ , and estimate an AR-model of order  $K$ . Let

$$\hat{e}(t) = y(t) + \sum_{k=1}^K \hat{\alpha}_k y(t-k)$$

- ▶ **Step 2:** Estimate the ARMA-model by minimizing

$$\|y - \hat{\Phi}\theta\|_2^2$$

where we get  $\hat{\Phi}$  by replacing  $e(t)$  with  $\hat{e}(t)$ .

# The two-stage LS method

- ▶ **Step 1:** Choose a sufficiently large  $K$ , and estimate an AR-model of order  $K$ . Let

$$\hat{e}(t) = y(t) + \sum_{k=1}^K \hat{\alpha}_k y(t-k)$$

- ▶ **Step 2:** Estimate the ARMA-model by minimizing

$$\|y - \hat{\Phi}\theta\|_2^2$$

where we get  $\hat{\Phi}$  by replacing  $e(t)$  with  $\hat{e}(t)$ . Can use either YW-method or LS method for this.

# The two-stage LS method

- ▶ **Step 1:** Choose a sufficiently large  $K$ , and estimate an AR-model of order  $K$ . Let

$$\hat{e}(t) = y(t) + \sum_{k=1}^K \hat{\alpha}_k y(t-k)$$

- ▶ **Step 2:** Estimate the ARMA-model by minimizing

$$\|y - \hat{\Phi}\theta\|_2^2$$

where we get  $\hat{\Phi}$  by replacing  $e(t)$  with  $\hat{e}(t)$ . Can use either YW-method or LS method for this.

- ▶ The variance  $\sigma^2$  is estimated by the sample variance of  $\hat{e}(t)$ .



# The two-stage LS method

- ▶ **Step 1:** Choose a sufficiently large  $K$ , and estimate an AR-model of order  $K$ . Let

$$\hat{e}(t) = y(t) + \sum_{k=1}^K \hat{\alpha}_k y(t-k)$$

- ▶ **Step 2:** Estimate the ARMA-model by minimizing

$$\|y - \hat{\Phi}\theta\|_2^2$$

where we get  $\hat{\Phi}$  by replacing  $e(t)$  with  $\hat{e}(t)$ . Can use either YW-method or LS method for this.

- ▶ The variance  $\sigma^2$  is estimated by the sample variance of  $\hat{e}(t)$ .
- ▶ Finally, the estimate of the spectrum is

$$\hat{\phi}(\omega) = \left| \frac{\hat{B}(\omega)}{\hat{A}(\omega)} \right|^2 \hat{\sigma}^2.$$

# Modified YW

---

- We want to estimate  $A(q)$  and  $B(q)$  in

$$A(q)y(k) = B(q)e(k)$$

# Modified YW

---

- ▶ We want to estimate  $A(q)$  and  $B(q)$  in

$$A(q)y(k) = B(q)e(k)$$

- ▶ From earlier in this lecture

$$r(k) + \sum_{i=1}^n a_i r(k-i) = 0, \quad k > m.$$

# Modified YW

---

- ▶ We want to estimate  $A(q)$  and  $B(q)$  in

$$A(q)y(k) = B(q)e(k)$$

- ▶ From earlier in this lecture

$$r(k) + \sum_{i=1}^n a_i r(k-i) = 0, \quad k > m.$$

- ▶ **Idea:** Use these equations to estimate  $A(q)$ .

# Modified YW

---

- ▶ We want to estimate  $A(q)$  and  $B(q)$  in

$$A(q)y(k) = B(q)e(k)$$

- ▶ From earlier in this lecture

$$r(k) + \sum_{i=1}^n a_i r(k-i) = 0, \quad k > m.$$

- ▶ **Idea:** Use these equations to estimate  $A(q)$ . We can then use  $\hat{A}(q)$  to estimate  $B(q)$ .

# Estimating the AR-part using modified YW

---

- Replace  $r(k)$  with estimates  $\hat{r}(k)$  to get,

$$\hat{r}(k) + \sum_{i=1}^n a_i \hat{r}(k-i) = 0, \quad k > m.$$

# Estimating the AR-part using modified YW

- ▶ Replace  $r(k)$  with estimates  $\hat{r}(k)$  to get,

$$\hat{r}(k) + \sum_{i=1}^n a_i \hat{r}(k-i) = 0, \quad k > m.$$

- ▶ or on matrix form for  $k = m+1, \dots, m+M$ ,

$$\begin{bmatrix} \hat{r}(m+1) \\ \vdots \\ \hat{r}(m+M) \end{bmatrix} + \begin{bmatrix} \hat{r}(m) & \cdots & \hat{r}(m-n+1) \\ \vdots & \ddots & \vdots \\ \hat{r}(m+M-1) & \cdots & \hat{r}(m-n+M) \end{bmatrix} \begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix} = \begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix}$$

# Estimating the AR-part using modified YW

- ▶ Replace  $r(k)$  with estimates  $\hat{r}(k)$  to get,

$$\hat{r}(k) + \sum_{i=1}^n a_i \hat{r}(k-i) = 0, \quad k > m.$$

- ▶ or on matrix form for  $k = m+1, \dots, m+M$ ,

$$\begin{bmatrix} \hat{r}(m+1) \\ \vdots \\ \hat{r}(m+M) \end{bmatrix} + \begin{bmatrix} \hat{r}(m) & \cdots & \hat{r}(m-n+1) \\ \vdots & \ddots & \vdots \\ \hat{r}(m+M-1) & \cdots & \hat{r}(m-n+M) \end{bmatrix} \begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix} = \begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix}$$

- ▶ Can estimate  $a_1, \dots, a_n$  with least squares.



# Estimating the AR-part using modified YW

- ▶ Replace  $r(k)$  with estimates  $\hat{r}(k)$  to get,

$$\hat{r}(k) + \sum_{i=1}^n a_i \hat{r}(k-i) = 0, \quad k > m.$$

- ▶ or on matrix form for  $k = m+1, \dots, m+M$ ,

$$\begin{bmatrix} \hat{r}(m+1) \\ \vdots \\ \hat{r}(m+M) \end{bmatrix} + \begin{bmatrix} \hat{r}(m) & \cdots & \hat{r}(m-n+1) \\ \vdots & \ddots & \vdots \\ \hat{r}(m+M-1) & \cdots & \hat{r}(m-n+M) \end{bmatrix} \begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix} = \begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix}$$

- ▶ Can estimate  $a_1, \dots, a_n$  with least squares.
- ▶ **User choice:** How large should  $M$  be?

# Estimating the AR-part using modified YW

- ▶ Replace  $r(k)$  with estimates  $\hat{r}(k)$  to get,

$$\hat{r}(k) + \sum_{i=1}^n a_i \hat{r}(k-i) = 0, \quad k > m.$$

- ▶ or on matrix form for  $k = m+1, \dots, m+M$ ,

$$\begin{bmatrix} \hat{r}(m+1) \\ \vdots \\ \hat{r}(m+M) \end{bmatrix} + \begin{bmatrix} \hat{r}(m) & \cdots & \hat{r}(m-n+1) \\ \vdots & \ddots & \vdots \\ \hat{r}(m+M-1) & \cdots & \hat{r}(m-n+M) \end{bmatrix} \begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix} = \begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix}$$

- ▶ Can estimate  $a_1, \dots, a_n$  with least squares.
- ▶ **User choice:** How large should  $M$  be?
  - ▶  $M = n$  gives square matrix.  $M > n$  overdetermined system.

# Estimating the AR-part using modified YW

- ▶ Replace  $r(k)$  with estimates  $\hat{r}(k)$  to get,

$$\hat{r}(k) + \sum_{i=1}^n a_i \hat{r}(k-i) = 0, \quad k > m.$$

- ▶ or on matrix form for  $k = m+1, \dots, m+M$ ,

$$\begin{bmatrix} \hat{r}(m+1) \\ \vdots \\ \hat{r}(m+M) \end{bmatrix} + \begin{bmatrix} \hat{r}(m) & \cdots & \hat{r}(m-n+1) \\ \vdots & \ddots & \vdots \\ \hat{r}(m+M-1) & \cdots & \hat{r}(m-n+M) \end{bmatrix} \begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix} = \begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix}$$

- ▶ Can estimate  $a_1, \dots, a_n$  with least squares.
- ▶ **User choice:** How large should  $M$  be?
  - ▶  $M = n$  gives square matrix.  $M > n$  overdetermined system.
  - ▶ Larger  $M$  needed when poles and zeros are closely spaced near the unit circle (narrowband spectrum).

# Estimating the AR-part using modified YW

- ▶ Replace  $r(k)$  with estimates  $\hat{r}(k)$  to get,

$$\hat{r}(k) + \sum_{i=1}^n a_i \hat{r}(k-i) = 0, \quad k > m.$$

- ▶ or on matrix form for  $k = m+1, \dots, m+M$ ,

$$\begin{bmatrix} \hat{r}(m+1) \\ \vdots \\ \hat{r}(m+M) \end{bmatrix} + \begin{bmatrix} \hat{r}(m) & \cdots & \hat{r}(m-n+1) \\ \vdots & \ddots & \vdots \\ \hat{r}(m+M-1) & \cdots & \hat{r}(m-n+M) \end{bmatrix} \begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix} = \begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix}$$

- ▶ Can estimate  $a_1, \dots, a_n$  with least squares.
- ▶ **User choice:** How large should  $M$  be?
  - ▶  $M = n$  gives square matrix.  $M > n$  overdetermined system.
  - ▶ Larger  $M$  needed when poles and zeros are closely spaced near the unit circle (narrowband spectrum).
  - ▶ When poles and zeros are both well inside the unit circle,  $M = n$  typically works well.

# The MA-part

---

- ▶ ARMA-model:  $A(q)y(k) = B(q)e(k)$ . Let  $x(k) = B(q)e(k)$ .

# The MA-part

---

- ▶ ARMA-model:  $A(q)y(k) = B(q)e(k)$ . Let  $x(k) = B(q)e(k)$ .
- ▶ Note that

$$\phi_y(\omega) = \frac{1}{|A(\omega)|^2} \phi_x(\omega).$$

# The MA-part

---

- ▶ ARMA-model:  $A(q)y(k) = B(q)e(k)$ . Let  $x(k) = B(q)e(k)$ .
- ▶ Note that

$$\phi_y(\omega) = \frac{1}{|A(\omega)|^2} \phi_x(\omega).$$

- ▶ Let  $\gamma_k$  be the ACS for  $x(k)$ , then  $\gamma_k = 0$  for  $|k| > m$ , and

$$\phi_x(\omega) =$$

# The MA-part

---

- ▶ ARMA-model:  $A(q)y(k) = B(q)e(k)$ . Let  $x(k) = B(q)e(k)$ .
- ▶ Note that

$$\phi_y(\omega) = \frac{1}{|A(\omega)|^2} \phi_x(\omega).$$

- ▶ Let  $\gamma_k$  be the ACS for  $x(k)$ , then  $\gamma_k = 0$  for  $|k| > m$ , and

$$\phi_x(\omega) = |B(\omega)|^2 \sigma^2 =$$



# The MA-part

- ▶ ARMA-model:  $A(q)y(k) = B(q)e(k)$ . Let  $x(k) = B(q)e(k)$ .
- ▶ Note that

$$\phi_y(\omega) = \frac{1}{|A(\omega)|^2} \phi_x(\omega).$$

- ▶ Let  $\gamma_k$  be the ACS for  $x(k)$ , then  $\gamma_k = 0$  for  $|k| > m$ , and

$$\phi_x(\omega) = |B(\omega)|^2 \sigma^2 = \sum_{k=-m}^m \gamma_k e^{-i\omega k}.$$

# The MA-part

- ▶ ARMA-model:  $A(q)y(k) = B(q)e(k)$ . Let  $x(k) = B(q)e(k)$ .
- ▶ Note that

$$\phi_y(\omega) = \frac{1}{|A(\omega)|^2} \phi_x(\omega).$$

- ▶ Let  $\gamma_k$  be the ACS for  $x(k)$ , then  $\gamma_k = 0$  for  $|k| > m$ , and

$$\phi_x(\omega) = |B(\omega)|^2 \sigma^2 = \sum_{k=-m}^m \gamma_k e^{-i\omega k}.$$

- ▶ Finally,

$$\gamma_k =$$

# The MA-part

- ▶ ARMA-model:  $A(q)y(k) = B(q)e(k)$ . Let  $x(k) = B(q)e(k)$ .
- ▶ Note that

$$\phi_y(\omega) = \frac{1}{|A(\omega)|^2} \phi_x(\omega).$$

- ▶ Let  $\gamma_k$  be the ACS for  $x(k)$ , then  $\gamma_k = 0$  for  $|k| > m$ , and

$$\phi_x(\omega) = |B(\omega)|^2 \sigma^2 = \sum_{k=-m}^m \gamma_k e^{-i\omega k}.$$

- ▶ Finally,

$$\gamma_k = E\{x(t)x^*(t-k)\} =$$

# The MA-part

- ▶ ARMA-model:  $A(q)y(k) = B(q)e(k)$ . Let  $x(k) = B(q)e(k)$ .
- ▶ Note that

$$\phi_y(\omega) = \frac{1}{|A(\omega)|^2} \phi_x(\omega).$$

- ▶ Let  $\gamma_k$  be the ACS for  $x(k)$ , then  $\gamma_k = 0$  for  $|k| > m$ , and

$$\phi_x(\omega) = |B(\omega)|^2 \sigma^2 = \sum_{k=-m}^m \gamma_k e^{-i\omega k}.$$

- ▶ Finally,

$$\begin{aligned} \gamma_k &= E\{x(t)x^*(t-k)\} = E\{[B(q)e(t)][B(q)e(t-k)]^*\} \\ &= \end{aligned}$$

# The MA-part

- ▶ ARMA-model:  $A(q)y(k) = B(q)e(k)$ . Let  $x(k) = B(q)e(k)$ .
- ▶ Note that

$$\phi_y(\omega) = \frac{1}{|A(\omega)|^2} \phi_x(\omega).$$

- ▶ Let  $\gamma_k$  be the ACS for  $x(k)$ , then  $\gamma_k = 0$  for  $|k| > m$ , and

$$\phi_x(\omega) = |B(\omega)|^2 \sigma^2 = \sum_{k=-m}^m \gamma_k e^{-i\omega k}.$$

- ▶ Finally,

$$\begin{aligned} \gamma_k &= E\{x(t)x^*(t-k)\} = E\{[B(q)e(t)][B(q)e(t-k)]^*\} \\ &= E\{[A(q)y(t)][A(q)y(t-k)]^*\} \\ &= \end{aligned}$$

# The MA-part

- ▶ ARMA-model:  $A(q)y(k) = B(q)e(k)$ . Let  $x(k) = B(q)e(k)$ .
- ▶ Note that

$$\phi_y(\omega) = \frac{1}{|A(\omega)|^2} \phi_x(\omega).$$

- ▶ Let  $\gamma_k$  be the ACS for  $x(k)$ , then  $\gamma_k = 0$  for  $|k| > m$ , and

$$\phi_x(\omega) = |B(\omega)|^2 \sigma^2 = \sum_{k=-m}^m \gamma_k e^{-i\omega k}.$$

- ▶ Finally,

$$\begin{aligned} \gamma_k &= E\{x(t)x^*(t-k)\} = E\{[B(q)e(t)][B(q)e(t-k)]^*\} \\ &= E\{[A(q)y(t)][A(q)y(t-k)]^*\} \\ &= \sum_{j=0}^n \sum_{p=0}^n a_j a_p^* r(k+p-j), \quad a_0 = 1. \end{aligned}$$

# Summary of modified YW

---

$$A(q)y(t) = B(q)e(t)$$

# Summary of modified YW

---

$$A(q)y(t) = B(q)e(t)$$

- ▶ Compute estimate  $\hat{r}(k)$  of  $r(k)$ .



# Summary of modified YW

---

$$A(q)y(t) = B(q)e(t)$$

- ▶ Compute estimate  $\hat{r}(k)$  of  $r(k)$ .
- ▶ Find an estimate  $\hat{A}(q)$  of  $A(q)$  from

$$\begin{bmatrix} \hat{r}(m+1) \\ \vdots \\ \hat{r}(m+M) \end{bmatrix} + \begin{bmatrix} \hat{r}(m) & \cdots & \hat{r}(m-n+1) \\ \vdots & \ddots & \vdots \\ \hat{r}(m+M-1) & \cdots & \hat{r}(m-n+M) \end{bmatrix} \begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix} = \begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix}$$

# Summary of modified YW

$$A(q)y(t) = B(q)e(t)$$

- ▶ Compute estimate  $\hat{r}(k)$  of  $r(k)$ .
- ▶ Find an estimate  $\hat{A}(q)$  of  $A(q)$  from

$$\begin{bmatrix} \hat{r}(m+1) \\ \vdots \\ \hat{r}(m+M) \end{bmatrix} + \begin{bmatrix} \hat{r}(m) & \cdots & \hat{r}(m-n+1) \\ \vdots & \ddots & \vdots \\ \hat{r}(m+M-1) & \cdots & \hat{r}(m-n+M) \end{bmatrix} \begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix} = \begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix}$$

- ▶ Estimate the ACS of  $B(q)e(t)$  with

$$\hat{\gamma}_k = \sum_{j=0}^n \sum_{p=0}^n \hat{a}_j \hat{a}_p^* \hat{r}(k+p-j).$$

# Summary of modified YW

$$A(q)y(t) = B(q)e(t)$$

- ▶ Compute estimate  $\hat{r}(k)$  of  $r(k)$ .
- ▶ Find an estimate  $\hat{A}(q)$  of  $A(q)$  from

$$\begin{bmatrix} \hat{r}(m+1) \\ \vdots \\ \hat{r}(m+M) \end{bmatrix} + \begin{bmatrix} \hat{r}(m) & \cdots & \hat{r}(m-n+1) \\ \vdots & \ddots & \vdots \\ \hat{r}(m+M-1) & \cdots & \hat{r}(m-n+M) \end{bmatrix} \begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix} = \begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix}$$

- ▶ Estimate the ACS of  $B(q)e(t)$  with

$$\hat{\gamma}_k = \sum_{j=0}^n \sum_{p=0}^n \hat{a}_j \hat{a}_p^* \hat{r}(k+p-j).$$

- ▶ The estimate of the ARMA spectrum is then

$$\hat{\phi}(\omega) =$$

# Summary of modified YW

$$A(q)y(t) = B(q)e(t)$$

- ▶ Compute estimate  $\hat{r}(k)$  of  $r(k)$ .
- ▶ Find an estimate  $\hat{A}(q)$  of  $A(q)$  from

$$\begin{bmatrix} \hat{r}(m+1) \\ \vdots \\ \hat{r}(m+M) \end{bmatrix} + \begin{bmatrix} \hat{r}(m) & \cdots & \hat{r}(m-n+1) \\ \vdots & \ddots & \vdots \\ \hat{r}(m+M-1) & \cdots & \hat{r}(m-n+M) \end{bmatrix} \begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix} = \begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix}$$

- ▶ Estimate the ACS of  $B(q)e(t)$  with

$$\hat{\gamma}_k = \sum_{j=0}^n \sum_{p=0}^n \hat{a}_j \hat{a}_p^* \hat{r}(k+p-j).$$

- ▶ The estimate of the ARMA spectrum is then

$$\hat{\phi}(\omega) = \frac{\sum_{k=-m}^m \hat{\gamma}_k e^{-i\omega k}}{|\hat{A}(\omega)|^2}.$$

# Summing up

- ▶ Parametric approach
- ▶ Rational spectra – AR(MA) processes
- ▶ Pole/zero placement (intuition)
- ▶ Covariance structure for AR(MA)
- ▶ Yule-Walker for AR(MA)
- ▶ Least squares for AR(MA)

Method	Computational Burden	Accuracy	Guarantee $\hat{\phi}(\omega) \geq 0$ ?	Use for
AR: YW or LS	low	medium	Yes	Spectra with (narrow) peaks but no valley
MA: BT	low	low-medium	No	Broadband spectra possibly with valleys but no peaks
ARMA: MYW	low-medium	medium	No	Spectra with both peaks and (not too deep) valleys
ARMA: 2-Stage LS	medium-high	medium-high	Yes	As above

# MATLAB

---

Useful MATLAB functions:

- ▶ `lsar(y,n)`
- ▶ `lsarma(y,n,m,K)`
- ▶ `yulewalker(y,n)`
- ▶ `mywarma(y,n,m,M)`
- ▶ `armase(b,a,sig2,L)`
- ▶ `argamse(gamma,a,L)`
- ▶ `zplane(b,a)`

**Note:** These methods assume that the mean value of  $y$  is 0. If this is not the case for your data set, start by removing the mean!