




# Line spectra, Covariance structure, and NLS



Per Mattsson

Systems and Control  
Department of Information Technology  
Uppsala University

2019-09-17

# Summary from last lecture

---

- ▶ Parametric methods  $\implies$  fewer parameters to estimate (bias/variance)
- ▶ AR and ARMA models for rational (continuous) spectra
- ▶ AR – only peaks
- ▶ ARMA – peaks and valleys
- ▶ 4 approaches:
  - ▶ YW (AR)
  - ▶ LSAR (AR)
  - ▶ MYW (ARMA, two-stage)
  - ▶ LSARMA (ARMA, two-stage)
- ▶ ARMA is nonlinear, methods have user parameters

**Today:** Estimating line spectra.



# Sinusoidal model

# Sinusoidal model

---

- ▶ Many applications (e.g. communications, radar/sonar etc) have signals with (near) sinusoidal components.

# Sinusoidal model

---

- ▶ Many applications (e.g. communications, radar/sonar etc) have signals with (near) sinusoidal components.
- ▶ Stable ARMA-models,  $y(t) = \frac{B(z)}{A(z)}e(t)$ , are poor approximations.

# Sinusoidal model

---

- ▶ Many applications (e.g. communications, radar/sonar etc) have signals with (near) sinusoidal components.
- ▶ Stable ARMA-models,  $y(t) = \frac{B(z)}{A(z)}e(t)$ , are poor approximations.
- ▶ **Complex sinusoid:**

$$x_k(t) = \alpha_k e^{i(\omega_k t + \varphi_k)}$$

# Sinusoidal model

---

- ▶ Many applications (e.g. communications, radar/sonar etc) have signals with (near) sinusoidal components.
- ▶ Stable ARMA-models,  $y(t) = \frac{B(z)}{A(z)}e(t)$ , are poor approximations.
- ▶ **Complex sinusoid:**

$$x_k(t) = \alpha_k e^{i(\omega_k t + \varphi_k)}$$

- ▶ Frequency:  $\omega_k$ . Amplitude:  $\alpha_k$ . Phase:  $\varphi_k$ .

# Sinusoidal model

- ▶ Many applications (e.g. communications, radar/sonar etc) have signals with (near) sinusoidal components.
- ▶ Stable ARMA-models,  $y(t) = \frac{B(z)}{A(z)}e(t)$ , are poor approximations.
- ▶ **Complex sinusoid:**

$$x_k(t) = \alpha_k e^{i(\omega_k t + \varphi_k)}$$

- ▶ Frequency:  $\omega_k$ . Amplitude:  $\alpha_k$ . Phase:  $\varphi_k$ .
- ▶ Model with  $n$  components and noise,

$$x(t) = \sum_{k=1}^n \alpha_k e^{i(\omega_k t + \varphi_k)} = \sum_{k=1}^n x_k(t),$$

$$y(t) = x(t) + e(t).$$



# Sinusoidal model, assumptions

---

$$y(t) = \underbrace{\sum_{k=1}^n \underbrace{\alpha_k e^{i(\omega_k t + \varphi_k)}}_{x_k(t)}}_{x(t)} + e(t)$$

# Sinusoidal model, assumptions

---

$$y(t) = \underbrace{\sum_{k=1}^n \underbrace{\alpha_k e^{i(\omega_k t + \varphi_k)}}_{x_k(t)}}_{x(t)} + e(t)$$

- **A1:**  $\alpha_k > 0$  and  $\omega_k \in [-\pi, \pi]$ . Prevents model ambiguities.

# Sinusoidal model, assumptions

---

$$y(t) = \underbrace{\sum_{k=1}^n \underbrace{\alpha_k e^{i(\omega_k t + \varphi_k)}}_{x_k(t)}}_{x(t)} + e(t)$$

- ▶ **A1:**  $\alpha_k > 0$  and  $\omega_k \in [-\pi, \pi]$ . Prevents model ambiguities.
- ▶ **A2:**  $\varphi_k$  are independent random variables uniformly distributed on  $[-\pi, \pi]$ .

# Sinusoidal model, assumptions

---

$$y(t) = \underbrace{\sum_{k=1}^n \underbrace{\alpha_k e^{i(\omega_k t + \varphi_k)}}_{x_k(t)}}_{x(t)} + e(t)$$

- ▶ **A1:**  $\alpha_k > 0$  and  $\omega_k \in [-\pi, \pi]$ . Prevents model ambiguities.
- ▶ **A2:**  $\varphi_k$  are independent random variables uniformly distributed on  $[-\pi, \pi]$ .
  - ▶ Mathematically convenient.

# Sinusoidal model, assumptions

---

$$y(t) = \underbrace{\sum_{k=1}^n \underbrace{\alpha_k e^{i(\omega_k t + \varphi_k)}}_{x_k(t)}}_{x(t)} + e(t)$$

- ▶ **A1:**  $\alpha_k > 0$  and  $\omega_k \in [-\pi, \pi]$ . Prevents model ambiguities.
- ▶ **A2:**  $\varphi_k$  are independent random variables uniformly distributed on  $[-\pi, \pi]$ .
  - ▶ Mathematically convenient.
  - ▶ Nuisance parameters (we are not interested in their values).

# Sinusoidal model, assumptions

---

$$y(t) = \underbrace{\sum_{k=1}^n \underbrace{\alpha_k e^{i(\omega_k t + \varphi_k)}}_{x_k(t)}}_{x(t)} + e(t)$$

- ▶ **A1:**  $\alpha_k > 0$  and  $\omega_k \in [-\pi, \pi]$ . Prevents model ambiguities.
- ▶ **A2:**  $\varphi_k$  are independent random variables uniformly distributed on  $[-\pi, \pi]$ .
  - ▶ Mathematically convenient.
  - ▶ Nuisance parameters (we are not interested in their values).
  - ▶ Realistic (no reason to assume any particular values).

# Sinusoidal model, assumptions

$$y(t) = \underbrace{\sum_{k=1}^n \underbrace{\alpha_k e^{i(\omega_k t + \varphi_k)}}_{x_k(t)}}_{x(t)} + e(t)$$

- ▶ **A1:**  $\alpha_k > 0$  and  $\omega_k \in [-\pi, \pi]$ . Prevents model ambiguities.
- ▶ **A2:**  $\varphi_k$  are independent random variables uniformly distributed on  $[-\pi, \pi]$ .
  - ▶ Mathematically convenient.
  - ▶ Nuisance parameters (we are not interested in their values).
  - ▶ Realistic (no reason to assume any particular values).
- ▶ **A3:**  $e(t)$  is circular white noise with variance  $\sigma^2$ .

# Sinusoidal model, assumptions

$$y(t) = \underbrace{\sum_{k=1}^n \underbrace{\alpha_k e^{i(\omega_k t + \varphi_k)}}_{x_k(t)}}_{x(t)} + e(t)$$

- ▶ **A1:**  $\alpha_k > 0$  and  $\omega_k \in [-\pi, \pi]$ . Prevents model ambiguities.
- ▶ **A2:**  $\varphi_k$  are independent random variables uniformly distributed on  $[-\pi, \pi]$ .
  - ▶ Mathematically convenient.
  - ▶ Nuisance parameters (we are not interested in their values).
  - ▶ Realistic (no reason to assume any particular values).
- ▶ **A3:**  $e(t)$  is circular white noise with variance  $\sigma^2$ . That is,

$$E\{e(t)e^*(s)\} = \sigma^2 \delta_{t,s}, \quad E\{e(t)e(s)\} = 0.$$



# Sinusoidal model, assumptions

$$y(t) = \underbrace{\sum_{k=1}^n \underbrace{\alpha_k e^{i(\omega_k t + \varphi_k)}}_{x_k(t)}}_{x(t)} + e(t)$$

- ▶ **A1:**  $\alpha_k > 0$  and  $\omega_k \in [-\pi, \pi]$ . Prevents model ambiguities.
- ▶ **A2:**  $\varphi_k$  are independent random variables uniformly distributed on  $[-\pi, \pi]$ .
  - ▶ Mathematically convenient.
  - ▶ Nuisance parameters (we are not interested in their values).
  - ▶ Realistic (no reason to assume any particular values).
- ▶ **A3:**  $e(t)$  is circular white noise with variance  $\sigma^2$ . That is,

$$E\{e(t)e^*(s)\} = \sigma^2 \delta_{t,s}, \quad E\{e(t)e(s)\} = 0.$$

- ▶ May be achieved by slow sampling or filtering.

# Sinusoidal model, assumptions

$$y(t) = \underbrace{\sum_{k=1}^n \underbrace{\alpha_k e^{i(\omega_k t + \varphi_k)}}_{x_k(t)}}_{x(t)} + e(t)$$

- ▶ **A1:**  $\alpha_k > 0$  and  $\omega_k \in [-\pi, \pi]$ . Prevents model ambiguities.
- ▶ **A2:**  $\varphi_k$  are independent random variables uniformly distributed on  $[-\pi, \pi]$ .
  - ▶ Mathematically convenient.
  - ▶ Nuisance parameters (we are not interested in their values).
  - ▶ Realistic (no reason to assume any particular values).
- ▶ **A3:**  $e(t)$  is circular white noise with variance  $\sigma^2$ . That is,

$$E\{e(t)e^*(s)\} = \sigma^2 \delta_{t,s}, \quad E\{e(t)e(s)\} = 0.$$

- ▶ May be achieved by slow sampling or filtering.
- ▶ Some methods are poor for colored noise!

# The spectrum

---

- ▶ **PSD**:  $\phi(\omega) = \sum_{k=-\infty}^{\infty} r(k)e^{-i\omega k}$ ,  $r(k) = E\{y(t)y^*(t-k)\}$ .
- ▶ With the sinusoidal model

$$E\{y(t)y^*(t-k)\} = \text{[Board]} =$$

# The spectrum

---

- ▶ **PSD**:  $\phi(\omega) = \sum_{k=-\infty}^{\infty} r(k)e^{-i\omega k}$ ,  $r(k) = E\{y(t)y^*(t-k)\}$ .
- ▶ With the sinusoidal model

$$E\{y(t)y^*(t-k)\} = \text{[Board]} = E\{x(t)x^*(t-k)\} + \sigma^2\delta_{k,0}.$$

# The spectrum

- ▶ **PSD:**  $\phi(\omega) = \sum_{k=-\infty}^{\infty} r(k)e^{-i\omega k}$ ,  $r(k) = E\{y(t)y^*(t-k)\}$ .
- ▶ With the sinusoidal model

$$E\{y(t)y^*(t-k)\} = \text{[Board]} = E\{x(t)x^*(t-k)\} + \sigma^2\delta_{k,0}.$$

- ▶ Note that

$$E\{x_p(t)x_j^*(t-k)\} =$$

# The spectrum

- ▶ **PSD:**  $\phi(\omega) = \sum_{k=-\infty}^{\infty} r(k)e^{-i\omega k}$ ,  $r(k) = E\{y(t)y^*(t-k)\}$ .
- ▶ With the sinusoidal model

$$E\{y(t)y^*(t-k)\} = [\text{Board}] = E\{x(t)x^*(t-k)\} + \sigma^2\delta_{k,0}.$$

- ▶ Note that

$$\begin{aligned} E\{x_p(t)x_j^*(t-k)\} &= \alpha_p\alpha_j e^{i(\omega_p-\omega_j)t} e^{i\omega_j k} E\{e^{i\varphi_p} e^{-i\varphi_j}\} \\ &= [\text{Board}] = \end{aligned}$$

# The spectrum

- ▶ **PSD:**  $\phi(\omega) = \sum_{k=-\infty}^{\infty} r(k)e^{-i\omega k}$ ,  $r(k) = E\{y(t)y^*(t-k)\}$ .
- ▶ With the sinusoidal model

$$E\{y(t)y^*(t-k)\} = [\text{Board}] = E\{x(t)x^*(t-k)\} + \sigma^2\delta_{k,0}.$$

- ▶ Note that

$$\begin{aligned} E\{x_p(t)x_j^*(t-k)\} &= \alpha_p\alpha_j e^{i(\omega_p-\omega_j)t} e^{i\omega_j k} E\{e^{i\varphi_p} e^{-i\varphi_j}\} \\ &= [\text{Board}] = \alpha_p^2 e^{i\omega_p k} \delta_{p,j}. \end{aligned}$$

# The spectrum

- ▶ **PSD:**  $\phi(\omega) = \sum_{k=-\infty}^{\infty} r(k)e^{-i\omega k}$ ,  $r(k) = E\{y(t)y^*(t-k)\}$ .
- ▶ With the sinusoidal model

$$E\{y(t)y^*(t-k)\} = [\text{Board}] = E\{x(t)x^*(t-k)\} + \sigma^2\delta_{k,0}.$$

- ▶ Note that

$$\begin{aligned} E\{x_p(t)x_j^*(t-k)\} &= \alpha_p\alpha_j e^{i(\omega_p-\omega_j)t} e^{i\omega_j k} E\{e^{i\varphi_p} e^{-i\varphi_j}\} \\ &= [\text{Board}] = \alpha_p^2 e^{i\omega_p k} \delta_{p,j}. \end{aligned}$$

- ▶ Hence

$$r(k) = [\text{Board}] =$$



# The spectrum

- ▶ **PSD:**  $\phi(\omega) = \sum_{k=-\infty}^{\infty} r(k)e^{-i\omega k}$ ,  $r(k) = E\{y(t)y^*(t-k)\}$ .
- ▶ With the sinusoidal model

$$E\{y(t)y^*(t-k)\} = [\text{Board}] = E\{x(t)x^*(t-k)\} + \sigma^2\delta_{k,0}.$$

- ▶ Note that

$$\begin{aligned} E\{x_p(t)x_j^*(t-k)\} &= \alpha_p\alpha_j e^{i(\omega_p-\omega_j)t} e^{i\omega_j k} E\{e^{i\varphi_p} e^{-i\varphi_j}\} \\ &= [\text{Board}] = \alpha_p^2 e^{i\omega_p k} \delta_{p,j}. \end{aligned}$$

- ▶ Hence

$$r(k) = [\text{Board}] = \sum_{p=1}^n \alpha_p^2 e^{i\omega_p k} + \sigma^2\delta_{k,0}.$$

# The spectrum

- ▶ **PSD:**  $\phi(\omega) = \sum_{k=-\infty}^{\infty} r(k)e^{-i\omega k}$ ,  $r(k) = E\{y(t)y^*(t-k)\}$ .
- ▶ With the sinusoidal model

$$E\{y(t)y^*(t-k)\} = [\text{Board}] = E\{x(t)x^*(t-k)\} + \sigma^2\delta_{k,0}.$$

- ▶ Note that

$$\begin{aligned} E\{x_p(t)x_j^*(t-k)\} &= \alpha_p\alpha_j e^{i(\omega_p-\omega_j)t} e^{i\omega_j k} E\{e^{i\varphi_p} e^{-i\varphi_j}\} \\ &= [\text{Board}] = \alpha_p^2 e^{i\omega_p k} \delta_{p,j}. \end{aligned}$$

- ▶ Hence

$$r(k) = [\text{Board}] = \sum_{p=1}^n \alpha_p^2 e^{i\omega_p k} + \sigma^2\delta_{k,0}.$$

- ▶ so,

$$\phi(\omega) = [\text{Board}] =$$

# The spectrum

- ▶ **PSD:**  $\phi(\omega) = \sum_{k=-\infty}^{\infty} r(k)e^{-i\omega k}$ ,  $r(k) = E\{y(t)y^*(t-k)\}$ .
- ▶ With the sinusoidal model

$$E\{y(t)y^*(t-k)\} = [\text{Board}] = E\{x(t)x^*(t-k)\} + \sigma^2\delta_{k,0}.$$

- ▶ Note that

$$\begin{aligned} E\{x_p(t)x_j^*(t-k)\} &= \alpha_p\alpha_j e^{i(\omega_p-\omega_j)t} e^{i\omega_j k} E\{e^{i\varphi_p} e^{-i\varphi_j}\} \\ &= [\text{Board}] = \alpha_p^2 e^{i\omega_p k} \delta_{p,j}. \end{aligned}$$

- ▶ Hence


$$r(k) = [\text{Board}] = \sum_{p=1}^n \alpha_p^2 e^{i\omega_p k} + \sigma^2\delta_{k,0}.$$

- ▶ so,

$$\phi(\omega) = [\text{Board}] = 2\pi \sum_{p=1}^n \alpha_p^2 \delta(\omega - \omega_p) + \sigma^2.$$

# The theoretical spectrum

$$\phi(\omega) = 2\pi \sum_{p=1}^n \alpha_p^2 \delta(\omega - \omega_p) + \sigma^2$$



Need to estimate  $\{\omega_k\}$ ,  $\{\alpha_k\}$  and  $\sigma^2$ .



# Parameter estimation

# Parameter estimation

---

- If  $\omega_k$  is given, then  $\alpha_k$  (and  $\varphi_k$ ) can be estimated from

$$y(t) = \sum_{k=1}^n \beta_k e^{i\omega_k t} + e(t),$$

with  $\beta_k = \alpha_k e^{i\varphi_k} \implies \alpha_k = |\beta_k|$ ,  $\varphi_k = \arg(\beta_k)$ .

# Parameter estimation

- If  $\omega_k$  is given, then  $\alpha_k$  (and  $\varphi_k$ ) can be estimated from

$$y(t) = \sum_{k=1}^n \beta_k e^{i\omega_k t} + e(t),$$

with  $\beta_k = \alpha_k e^{i\varphi_k} \implies \alpha_k = |\beta_k|$ ,  $\varphi_k = \arg(\beta_k)$ .

- Using least squares, we get

$$\min_{\beta} \|y - B\beta\|_2^2,$$

$$y = \begin{bmatrix} y(1) \\ \vdots \\ y(N) \end{bmatrix}, \quad B = \begin{bmatrix} e^{i\omega_1} & \dots & e^{i\omega_n} \\ \vdots & & \vdots \\ e^{iN\omega_1} & \dots & e^{iN\omega_n} \end{bmatrix}, \quad \beta = \begin{bmatrix} \beta_1 \\ \vdots \\ \beta_n \end{bmatrix}$$

# Parameter estimation

- If  $\omega_k$  is given, then  $\alpha_k$  (and  $\varphi_k$ ) can be estimated from

$$y(t) = \sum_{k=1}^n \beta_k e^{i\omega_k t} + e(t),$$

with  $\beta_k = \alpha_k e^{i\varphi_k} \implies \alpha_k = |\beta_k|$ ,  $\varphi_k = \arg(\beta_k)$ .

- Using least squares, we get

$$\min_{\beta} \|y - B\beta\|_2^2,$$

$$y = \begin{bmatrix} y(1) \\ \vdots \\ y(N) \end{bmatrix}, \quad B = \begin{bmatrix} e^{i\omega_1} & \dots & e^{i\omega_n} \\ \vdots & & \vdots \\ e^{iN\omega_1} & \dots & e^{iN\omega_n} \end{bmatrix}, \quad \beta = \begin{bmatrix} \beta_1 \\ \vdots \\ \beta_n \end{bmatrix}$$

- However, in general  $\omega_k$  are not given!



# Nonlinear least squares

---

- What if we optimize over  $\omega_k$  too? That is, minimize

$$f(\omega, \alpha, \varphi) = \sum_{k=1}^N \left| y(t) - \sum_{k=1}^n \alpha_k e^{i(\omega_k t + \varphi_k)} \right|^2 = \|y - B\beta\|_2^2.$$

# Nonlinear least squares

- ▶ What if we optimize over  $\omega_k$  too? That is, minimize

$$f(\omega, \alpha, \varphi) = \sum_{k=1}^N \left| y(t) - \sum_{k=1}^n \alpha_k e^{i(\omega_k t + \varphi_k)} \right|^2 = \|y - B\beta\|_2^2.$$

- ▶ (+) Maximum likelihood estimator if  $e(t)$  is Gaussian.

# Nonlinear least squares

- ▶ What if we optimize over  $\omega_k$  too? That is, minimize

$$f(\omega, \alpha, \varphi) = \sum_{k=1}^N \left| y(t) - \sum_{k=1}^n \alpha_k e^{i(\omega_k t + \varphi_k)} \right|^2 = \|y - B\beta\|_2^2.$$

- ▶ (+) Maximum likelihood estimator if  $e(t)$  is Gaussian.
- ▶ (-) Nonlinear in  $\omega_k$ , hard to solve (local minima).

# Nonlinear least squares

- ▶ What if we optimize over  $\omega_k$  too? That is, minimize

$$f(\omega, \alpha, \varphi) = \sum_{k=1}^N \left| y(t) - \sum_{k=1}^n \alpha_k e^{i(\omega_k t + \varphi_k)} \right|^2 = \|y - B\beta\|_2^2.$$

- ▶ (+) Maximum likelihood estimator if  $e(t)$  is Gaussian.
- ▶ (-) Nonlinear in  $\omega_k$ , hard to solve (local minima).

**Concentrate out  $\beta$ :**

- ▶ The  $\beta$  that minimizes  $f$  is given by

$$\hat{\beta} =$$

# Nonlinear least squares

- ▶ What if we optimize over  $\omega_k$  too? That is, minimize

$$f(\omega, \alpha, \varphi) = \sum_{k=1}^N \left| y(t) - \sum_{k=1}^n \alpha_k e^{i(\omega_k t + \varphi_k)} \right|^2 = \|y - B\beta\|_2^2.$$

- ▶ (+) Maximum likelihood estimator if  $e(t)$  is Gaussian.
- ▶ (-) Nonlinear in  $\omega_k$ , hard to solve (local minima).

**Concentrate out  $\beta$ :**

- ▶ The  $\beta$  that minimizes  $f$  is given by

$$\hat{\beta} = (B^* B)^{-1} B^* y.$$

# Nonlinear least squares

- ▶ What if we optimize over  $\omega_k$  too? That is, minimize

$$f(\omega, \alpha, \varphi) = \sum_{k=1}^N \left| y(t) - \sum_{k=1}^n \alpha_k e^{i(\omega_k t + \varphi_k)} \right|^2 = \|y - B\beta\|_2^2.$$

- ▶ (+) Maximum likelihood estimator if  $e(t)$  is Gaussian.
- ▶ (-) Nonlinear in  $\omega_k$ , hard to solve (local minima).

**Concentrate out  $\beta$ :**

- ▶ The  $\beta$  that minimizes  $f$  is given by

$$\hat{\beta} = (B^* B)^{-1} B^* y.$$

- ▶ Inserting this back into  $f$ , we can see that  $\hat{\omega}$  can be found by

$$\min_{\omega} \|y - B\hat{\beta}\|_2^2 =$$

# Nonlinear least squares

- ▶ What if we optimize over  $\omega_k$  too? That is, minimize

$$f(\omega, \alpha, \varphi) = \sum_{k=1}^N \left| y(t) - \sum_{k=1}^n \alpha_k e^{i(\omega_k t + \varphi_k)} \right|^2 = \|y - B\beta\|_2^2.$$

- ▶ (+) Maximum likelihood estimator if  $e(t)$  is Gaussian.
- ▶ (-) Nonlinear in  $\omega_k$ , hard to solve (local minima).

**Concentrate out  $\beta$ :**

- ▶ The  $\beta$  that minimizes  $f$  is given by

$$\hat{\beta} = (B^* B)^{-1} B^* y.$$

- ▶ Inserting this back into  $f$ , we can see that  $\hat{\omega}$  can be found by

$$\min_{\omega} \|y - B\hat{\beta}\|_2^2 = \min_{\omega} \|y - B(B^* B)^{-1} B^* y\|_2^2$$

# Nonlinear least squares, properties

---


$$\begin{aligned}\hat{\omega} &= \underset{\omega}{\operatorname{argmin}} \|y - B(B^*B)^{-1}B^*y\|_2^2 \\ \hat{\beta} &= (B^*B)^{-1}B^*y \Big|_{\omega=\hat{\omega}}\end{aligned}$$

- ▶ Gives a consistent estimate.
- ▶ For  $N \gg 1$ ,  $\operatorname{var}(\hat{\omega}_k) = \frac{6\sigma^2}{N^3\alpha_k^2}$ .
- ▶ Best (minimum variance) unbiased estimator of  $\omega_k$ .
- ▶ **Problem:** No (general) good method available to carry out the minimization.



# Nonlinear least squares, example

The cost function when  $y(t) = e^{it} + e(t)$ , and  $N = 100$ :



No general way to find the global minimum!

# Periodogram vs NLS

---

- ▶ For  $n = 1$  (one complex sinusoid) it can be shown that the NLS estimate is given by

$$\hat{\omega}_1 =$$

# Periodogram vs NLS

---

- ▶ For  $n = 1$  (one complex sinusoid) it can be shown that the NLS estimate is given by

$$\hat{\omega}_1 = \underset{\omega}{\operatorname{argmax}} \hat{\phi}_p(\omega).$$

# Periodogram vs NLS

---

- ▶ For  $n = 1$  (one complex sinusoid) it can be shown that the NLS estimate is given by

$$\hat{\omega}_1 = \operatorname{argmax}_{\omega} \hat{\phi}_p(\omega).$$

- ▶ For  $n > 1$  and

$$|\omega_p - \omega_k| > 2\pi/N, \quad (\text{resolution limit})$$

the NLS estimates is approximately the location of the  $n$  largest peaks in  $\hat{\phi}_p(\omega)$ .

# Periodogram vs NLS

---

- ▶ For  $n = 1$  (one complex sinusoid) it can be shown that the NLS estimate is given by

$$\hat{\omega}_1 = \operatorname{argmax}_{\omega} \hat{\phi}_p(\omega).$$

- ▶ For  $n > 1$  and

$$|\omega_p - \omega_k| > 2\pi/N, \quad (\text{resolution limit})$$

the NLS estimates is approximately the location of the  $n$  largest peaks in  $\hat{\phi}_p(\omega)$ .

- ▶ For tightly spaced peaks, super resolution are needed!



# Models of sinusoidal signals in noise

# ARMA-model

---

$$y(t) = \sum_{k=1}^n \underbrace{\alpha_k e^{i(\omega_k t + \varphi_k)}}_{x_k(t)} + e(t).$$

# ARMA-model

---

$$y(t) = \sum_{k=1}^n \underbrace{\alpha_k e^{i(\omega_k t + \varphi_k)}}_{x_k(t)} + e(t).$$

► **Annihilating filter:**  $(1 - e^{i\omega_k} z^{-1})x_k(t) = 0$ .



# ARMA-model

---

$$y(t) = \sum_{k=1}^n \underbrace{\alpha_k e^{i(\omega_k t + \varphi_k)}}_{x_k(t)} + e(t).$$

► **Annihilating filter:**  $(1 - e^{i\omega_k} z^{-1})x_k(t) = 0$ .

► Let

$$A(z) = \prod_{k=1}^n (1 - e^{i\omega_k} z^{-1}),$$

then

$$A(z)y(t) =$$

# ARMA-model

---

$$y(t) = \sum_{k=1}^n \underbrace{\alpha_k e^{i(\omega_k t + \varphi_k)}}_{x_k(t)} + e(t).$$

► **Annihilating filter:**  $(1 - e^{i\omega_k} z^{-1})x_k(t) = 0$ .

► Let

$$A(z) = \prod_{k=1}^n (1 - e^{i\omega_k} z^{-1}),$$

then

$$A(z)y(t) = A(z)e(t)$$

# ARMA-model

---

$$y(t) = \sum_{k=1}^n \underbrace{\alpha_k e^{i(\omega_k t + \varphi_k)}}_{x_k(t)} + e(t).$$

► **Annihilating filter:**  $(1 - e^{i\omega_k} z^{-1})x_k(t) = 0$ .

► Let

$$A(z) = \prod_{k=1}^n (1 - e^{i\omega_k} z^{-1}),$$

then

$$A(z)y(t) = A(z)e(t)$$

► Can estimate  $\omega_k$  by estimating  $A(z)$

# ARMA-model

$$y(t) = \sum_{k=1}^n \underbrace{\alpha_k e^{i(\omega_k t + \varphi_k)}}_{x_k(t)} + e(t).$$

► **Annihilating filter:**  $(1 - e^{i\omega_k} z^{-1})x_k(t) = 0$ .

► Let

$$A(z) = \prod_{k=1}^n (1 - e^{i\omega_k} z^{-1}),$$

then

$$A(z)y(t) = A(z)e(t)$$

► Can estimate  $\omega_k$  by estimating  $A(z)$

► **Note:** Can't divide away  $A(z)$  due to zeros on the unit circle!

# ARMA-model

$$y(t) = \sum_{k=1}^n \underbrace{\alpha_k e^{i(\omega_k t + \varphi_k)}}_{x_k(t)} + e(t).$$

► **Annihilating filter:**  $(1 - e^{i\omega_k} z^{-1})x_k(t) = 0$ .

► Let

$$A(z) = \prod_{k=1}^n (1 - e^{i\omega_k} z^{-1}),$$

then

$$A(z)y(t) = A(z)e(t)$$

- Can estimate  $\omega_k$  by estimating  $A(z)$
- **Note:** Can't divide away  $A(z)$  due to zeros on the unit circle!
- Use Modified Yule-Walker?

# ARMA-model

$$y(t) = \sum_{k=1}^n \underbrace{\alpha_k e^{i(\omega_k t + \varphi_k)}}_{x_k(t)} + e(t).$$

► **Annihilating filter:**  $(1 - e^{i\omega_k} z^{-1})x_k(t) = 0$ .

► Let

$$A(z) = \prod_{k=1}^n (1 - e^{i\omega_k} z^{-1}),$$

then

$$A(z)y(t) = A(z)e(t)$$

- Can estimate  $\omega_k$  by estimating  $A(z)$
- **Note:** Can't divide away  $A(z)$  due to zeros on the unit circle!
- Use Modified Yule-Walker?
- Can be improved by instead estimating  $B(z) = A(z)\tilde{A}(z)$  in

$$B(z)y(t) = B(z)e(t).$$

# Covariance model (1)

---

Notation:

$$a(\omega) \triangleq [1, e^{-i\omega}, \dots, e^{-i(m-1)\omega}]^T \quad (m \times 1)$$

$$A_m = [a(\omega_1), \dots, a(\omega_n)] \quad (m \times n)$$

For some  $m \geq n$

$\text{rank}(A_m) = n \quad \text{if} \quad \omega_k \neq \omega_p \text{ for } k \neq p$

# Covariance model (1)

Notation:

$$a(\omega) \triangleq [1, e^{-i\omega}, \dots, e^{-i(m-1)\omega}]^T \quad (m \times 1)$$

$$A_m = [a(\omega_1), \dots, a(\omega_n)] \quad (m \times n)$$

For some  $m \geq n$

$$\boxed{\text{rank}(A_m) = n \quad \text{if} \quad \omega_k \neq \omega_p \text{ for } k \neq p}$$

We can write

$$\tilde{y}_m(t) \triangleq \begin{bmatrix} y(t) \\ y(t-1) \\ \vdots \\ y(t-m+1) \end{bmatrix} = A_m \tilde{x}_m(t) + \tilde{e}_m(t)$$

where

$$\tilde{x}_m(t) = [x_1(t), \dots, x_n(t)]^T, \quad \tilde{e}_m(t) = [e(t), \dots, e(t-m+1)]^T.$$



## Covariance model (2)

---

This gives the covariance matrix model

$$R \triangleq E\{\tilde{y}_m(t)\tilde{y}_m^*(t)\} = A_m P A_m^* + \sigma^2 I, \quad P = \begin{bmatrix} \alpha_1^2 & & 0 \\ & \ddots & \\ 0 & & \alpha_n^2 \end{bmatrix}$$

The eigenstructure of  $R$  contains complete frequency information.

# Covariance model (2)

This gives the covariance matrix model

$$R \triangleq E\{\tilde{y}_m(t)\tilde{y}_m^*(t)\} = A_m P A_m^* + \sigma^2 I, \quad P = \begin{bmatrix} \alpha_1^2 & & 0 \\ & \ddots & \\ 0 & & \alpha_n^2 \end{bmatrix}$$

The eigenstructure of  $R$  contains complete frequency information. Furthermore, using non-overlapping parts of  $\{y(t)\}$  (uncorrelated)

$$\Gamma \triangleq E\{\tilde{y}_M(t-L-1)\tilde{y}_{L+1}^*(t)\} = A_M P_{L+1} A_{L+1}^*$$

where

$$P_K = \begin{bmatrix} \alpha_1^2 e^{-i\omega_1 K} & & 0 \\ & \ddots & \\ 0 & & \alpha_n^2 e^{-i\omega_n K} \end{bmatrix}$$

Null space of  $\Gamma$  ( $L, M \geq n$ ) contains complete frequency info.

# Singular value decomposition (SVD)

## Definition

Factor a matrix  $A \in \mathbb{C}^{m \times n}$  as

$$A = U \Sigma V^*$$

where  $U \in \mathbb{C}^{m \times m}$  and  $V \in \mathbb{C}^{n \times n}$  are unitary matrices, and  $\Sigma \in \mathbb{R}^{m \times n}$  is diagonal with non-negative diagonal values  $\sigma_i$  (singular values) ordered in decreasing order

(See A.4 in the book for more details)

- ▶ Low-rank approximations
- ▶ Useful to find the best fit subspaces of  $A$
- ▶ Projection onto subspace – perpendicular distance
- ▶ LS fit – vertical distance

## SVD (2)

$A \in \mathbb{C}^{m \times n}$  is of rank  $r < \min(m, n)$  then

$$A = \begin{bmatrix} U_1 & U_2 \end{bmatrix} \begin{bmatrix} \Sigma_1 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} V_1^* \\ V_2^* \end{bmatrix} = U_1 \Sigma_1 V_1^*$$

where  $\Sigma_1 \in \mathbb{R}^{r \times r}$ , then

- ▶  $U_1$  is an orthonormal basis in  $\mathcal{R}(A)$
- ▶  $U_2$  is an orthonormal basis in  $\mathcal{N}(A^*)$
- ▶  $V_1$  is an orthonormal basis in  $\mathcal{R}(A^*)$
- ▶  $V_2$  is an orthonormal basis in  $\mathcal{N}(A)$

Moreover,  $\mathcal{R}(A)$  and  $\mathcal{N}(A^*)$  are orthogonal to each other and they span  $\mathbb{C}^m$

( $\mathcal{N}(A^*)$  is the orthogonal complement to  $\mathcal{R}(A)$  and vice versa)

# Summary

---

- ▶ Signal model for line spectra
- ▶ Estimating the spectrum  $\implies$  frequency estimation
- ▶ Models of the covariance matrix and ARMA-type model
- ▶ Nonlinear least squares

# Summary

---

- ▶ Signal model for line spectra
- ▶ Estimating the spectrum  $\implies$  frequency estimation
- ▶ Models of the covariance matrix and ARMA-type model
- ▶ Nonlinear least squares

## Next lecture:

- ▶ Methods for estimating line spectra (super-resolution):  
HOYW, MUSIC, Min-norm, ESPRIT
- ▶ All exploit eigen-decomposition of the covariance matrix and  
corresponding subspaces

Interesting complements: 4.9.5 Minimization methods, Appendix B  
Cramér-Rao bound