




## STABILITY OF IMAGE RECONSTRUCTION ALGORITHMS

Pol del Aguila Pila, Ph.D.

Research Staff Scientist, CIBM SP EPFL Mathematical Imaging Section  
Postdoctoral researcher, Biomedical Imaging Group, EPFL  
[go.epfl.ch/Pol](mailto:go.epfl.ch/Pol), [pol.delaguila.pila@epfl.ch](mailto:pol.delaguila.pila@epfl.ch)  
CIBM Breakfast and Science, March 29, 2022




Joint work with Dr. S. Neumayer and Prof. M. Unser




## IMAGING AS AN INVERSE PROBLEM

### BIOMEDICAL IMAGING AS AN INVERSE PROBLEM




### BIOMEDICAL IMAGING AS AN INVERSE PROBLEM



- $\mathbf{n}$ : (approximately) additive statistical variability

## BIOMEDICAL IMAGING AS AN INVERSE PROBLEM




- $\mathbf{n}$ : (approximately) additive statistical variability
- Ill-posed inverse problem: Many  $\mathbf{s}$  could lead to the same  $y$




## BIOMEDICAL IMAGING AS AN INVERSE PROBLEM

Example: PET



## BIOMEDICAL IMAGING AS AN INVERSE PROBLEM

Example: PET



## FORWARD IMAGING MODEL

Unknown molecular/anatomical map:  $s(\mathbf{r}), \mathbf{r} = (x, y, z, t) \in \mathbb{R}^d$   
*defined over a continuum in space-time*

## FORWARD IMAGING MODEL

Unknown molecular/anatomical map:  $s(\mathbf{r}), \mathbf{r} = (x, y, z, t) \in \mathbb{R}^d$

*defined over a continuum in space-time*

$$s \in L_2(\mathbb{R}^d) \quad (\text{space of finite-energy functions})$$

## FORWARD IMAGING MODEL

Unknown molecular/anatomical map:  $s(\mathbf{r}), \mathbf{r} = (x, y, z, t) \in \mathbb{R}^d$

*defined over a continuum in space-time*

$$s \in L_2(\mathbb{R}^d) \quad (\text{space of finite-energy functions})$$

Imaging operator  $\mathbf{H} : s \mapsto \mathbf{y} = (y_1, \dots, y_M) = \mathbf{H}\{s\}$

*from continuum to discrete (finite dimensional)*

$$\mathbf{H} : L_2(\mathbb{R}^d) \rightarrow \mathbb{R}^M$$

5

5

## FORWARD IMAGING MODEL

Unknown molecular/anatomical map:  $s(\mathbf{r}), \mathbf{r} = (x, y, z, t) \in \mathbb{R}^d$

*defined over a continuum in space-time*

$$s \in L_2(\mathbb{R}^d) \quad (\text{space of finite-energy functions})$$

Imaging operator  $\mathbf{H} : s \mapsto \mathbf{y} = (y_1, \dots, y_M) = \mathbf{H}\{s\}$

*from continuum to discrete (finite dimensional)*

$$\mathbf{H} : L_2(\mathbb{R}^d) \rightarrow \mathbb{R}^M$$

Linearity assumption: for all  $s_1, s_2 \in L_2(\mathbb{R}^d), \alpha_1, \alpha_2 \in \mathbb{R}$

$$\mathbf{H}\{\alpha_1 s_1 + \alpha_2 s_2\} = \alpha_1 \mathbf{H}\{s_1\} + \alpha_2 \mathbf{H}\{s_2\}$$

## FORWARD IMAGING MODEL

Unknown molecular/anatomical map:  $s(\mathbf{r}), \mathbf{r} = (x, y, z, t) \in \mathbb{R}^d$

*defined over a continuum in space-time*

$$s \in L_2(\mathbb{R}^d) \quad (\text{space of finite-energy functions})$$

Imaging operator  $\mathbf{H} : s \mapsto \mathbf{y} = (y_1, \dots, y_M) = \mathbf{H}\{s\}$

*from continuum to discrete (finite dimensional)*

$$\mathbf{H} : L_2(\mathbb{R}^d) \rightarrow \mathbb{R}^M$$

Linearity assumption: for all  $s_1, s_2 \in L_2(\mathbb{R}^d), \alpha_1, \alpha_2 \in \mathbb{R}$

$$\mathbf{H}\{\alpha_1 s_1 + \alpha_2 s_2\} = \alpha_1 \mathbf{H}\{s_1\} + \alpha_2 \mathbf{H}\{s_2\}$$

$$\Rightarrow [\mathbf{y}]_m = y_m = \langle \eta_m, s \rangle = \int_{\mathbb{R}^d} \eta_m(\mathbf{r}) s(\mathbf{r}) d\mathbf{r}$$

↑ impulse response of the  $m$ th detector  
(by the Riesz representation theorem)

5

5

## FORWARD IMAGING MODEL

Unknown molecular/anatomical map:  $s(\mathbf{r}), \mathbf{r} = (x, y, z, t) \in \mathbb{R}^d$

*defined over a continuum in space-time*

$$s \in L_2(\mathbb{R}^d) \quad (\text{space of finite-energy functions})$$

Imaging operator  $H : s \mapsto \mathbf{y} = (y_1, \dots, y_M) = H\{s\}$

*from continuum to discrete (finite dimensional)*

$$H : L_2(\mathbb{R}^d) \rightarrow \mathbb{R}^M$$

Linearity assumption: for all  $s_1, s_2 \in L_2(\mathbb{R}^d), \alpha_1, \alpha_2 \in \mathbb{R}$

$$H\{\alpha_1 s_1 + \alpha_2 s_2\} = \alpha_1 H\{s_1\} + \alpha_2 H\{s_2\}$$

$$\Rightarrow [y]_m = y_m = \langle \eta_m, s \rangle = \int_{\mathbb{R}^d} \eta_m(\mathbf{r}) s(\mathbf{r}) d\mathbf{r}$$

(by the Riesz representation theorem)

$\eta_m(\cdot)$ : analysis function

5

## FORWARD IMAGING MODEL

Unknown molecular/anatomical map:  $s(\mathbf{r}), \mathbf{r} = (x, y, z, t) \in \mathbb{R}^d$

*defined over a continuum in space-time*

$$s \in L_2(\mathbb{R}^d) \quad (\text{space of finite-energy functions})$$

Imaging operator  $H : s \mapsto \mathbf{y} = (y_1, \dots, y_M) = H\{s\}$

*from continuum to discrete (finite dimensional)*

$$H : L_2(\mathbb{R}^d) \rightarrow \mathbb{R}^M$$


Linearity assumption: for all  $s_1, s_2 \in L_2(\mathbb{R}^d), \alpha_1, \alpha_2 \in \mathbb{R}$

$$H\{\alpha_1 s_1 + \alpha_2 s_2\} = \alpha_1 H\{s_1\} + \alpha_2 H\{s_2\}$$

$$\Rightarrow [y]_m = y_m = \langle \eta_m, s \rangle = \int_{\mathbb{R}^d} \eta_m(\mathbf{r}) s(\mathbf{r}) d\mathbf{r}$$

(by the Riesz representation theorem)

$\eta_m(\cdot)$ : analysis function



5

## FORWARD IMAGING MODEL

Unknown molecular/anatomical map:  $s(\mathbf{r}), \mathbf{r} = (x, y, z, t) \in \mathbb{R}^d$

*defined over a continuum in space-time*

$$s \in L_2(\mathbb{R}^d) \quad (\text{space of finite-energy functions})$$

Imaging operator  $H : s \mapsto \mathbf{y} = (y_1, \dots, y_M) = H\{s\}$

*from continuum to discrete (finite dimensional)*

$$H : L_2(\mathbb{R}^d) \rightarrow \mathbb{R}^M$$

Linearity assumption: for all  $s_1, s_2 \in L_2(\mathbb{R}^d), \alpha_1, \alpha_2 \in \mathbb{R}$


$$H\{\alpha_1 s_1 + \alpha_2 s_2\} = \alpha_1 H\{s_1\} + \alpha_2 H\{s_2\}$$

$$\Rightarrow [y]_m = y_m = \langle \eta_m, s \rangle = \int_{\mathbb{R}^d} \eta_m(\mathbf{r}) s(\mathbf{r}) d\mathbf{r}$$

(by the Riesz representation theorem)

$\eta_m(\cdot)$ : analysis function

5



## FORWARD IMAGING MODEL

Unknown molecular/anatomical map:  $s(\mathbf{r}), \mathbf{r} = (x, y, z, t) \in \mathbb{R}^d$

*defined over a continuum in space-time*

$$s \in L_2(\mathbb{R}^d) \quad (\text{space of finite-energy functions})$$

Imaging operator  $H : s \mapsto \mathbf{y} = (y_1, \dots, y_M) = H\{s\}$

*from continuum to discrete (finite dimensional)*

$$H : L_2(\mathbb{R}^d) \rightarrow \mathbb{R}^M$$


Linearity assumption: for all  $s_1, s_2 \in L_2(\mathbb{R}^d), \alpha_1, \alpha_2 \in \mathbb{R}$

$$H\{\alpha_1 s_1 + \alpha_2 s_2\} = \alpha_1 H\{s_1\} + \alpha_2 H\{s_2\}$$

$$\Rightarrow [y]_m = y_m = \langle \eta_m, s \rangle = \int_{\mathbb{R}^d} \eta_m(\mathbf{r}) s(\mathbf{r}) d\mathbf{r}$$

(by the Riesz representation theorem)

$\eta_m(\cdot)$ : analysis function



5

## FORWARD IMAGING MODEL

Unknown molecular/anatomical map:  $s(\mathbf{r}), \mathbf{r} = (x, y, z, t) \in \mathbb{R}^d$

*defined over a continuum in space-time*

$$s \in L_2(\mathbb{R}^d) \quad (\text{space of finite-energy functions})$$

Imaging operator  $H : s \mapsto \mathbf{y} = (y_1, \dots, y_M) = H\{s\}$

*from continuum to discrete (finite dimensional)*

$$H : L_2(\mathbb{R}^d) \rightarrow \mathbb{R}^M$$

Linearity assumption: for all  $s_1, s_2 \in L_2(\mathbb{R}^d), \alpha_1, \alpha_2 \in \mathbb{R}$

$$H\{\alpha_1 s_1 + \alpha_2 s_2\} = \alpha_1 H\{s_1\} + \alpha_2 H\{s_2\}$$

$$\Rightarrow [\mathbf{y}]_m = y_m = \langle \eta_m, s \rangle = \int_{\mathbb{R}^d} \eta_m(\mathbf{r}) s(\mathbf{r}) d\mathbf{r}$$

(by the Riesz representation theorem)

$\eta_m(\cdot)$ : analysis function




5

## FORWARD IMAGING MODEL

Example: MRI

■ Magnetic resonance:  $\omega_0 = \gamma B_0$

Frequency encode:




## FORWARD IMAGING MODEL

Example: MRI

■ Magnetic resonance:  $\omega_0 = \gamma B_0$

Frequency encode:



■ Linear forward model for MRI

$$\hat{s}(\omega_m) = \int_{\mathbb{R}^3} s(\mathbf{r}) e^{-j\langle \omega_m, \mathbf{r} \rangle} d\mathbf{r}$$

(sampling of Fourier transform)

Equivalent analysis functions:  $\eta_m(\mathbf{r}) = e^{j\langle \omega_m, \mathbf{r} \rangle}$  (complex sinusoids)



6


6

## FORWARD IMAGING MODEL

Example: MRI

- Magnetic resonance:  $\omega_0 = \gamma B_0$

Frequency encode:



- Linear forward model for MRI

$$\mathbf{r} = (x, y, z)$$

$$\hat{s}(\omega_m) = \int_{\mathbb{R}^3} s(\mathbf{r}) e^{-j\langle \omega_m, \mathbf{r} \rangle} d\mathbf{r} \quad (\text{sampling of Fourier transform})$$

$$\text{Equivalent analysis functions: } \eta_m(\mathbf{r}) = e^{j\langle \omega_m, \mathbf{r} \rangle} \quad (\text{complex sinusoids})$$

- Extended forward model with coil sensitivity

$$\hat{s}_w(\omega_m) = \int_{\mathbb{R}^3} w(\mathbf{r}) s(\mathbf{r}) e^{-j\langle \omega_m, \mathbf{r} \rangle} d\mathbf{r}$$

$$\eta_m(\mathbf{r}) = w(\mathbf{r}) e^{j\langle \omega_m, \mathbf{r} \rangle}$$

6



## FORWARD IMAGING MODEL

In summary

$$[\mathbf{H}\mathbf{s}]_m = y_m = \langle \eta_m, \mathbf{s} \rangle \\ = \int_{\mathbb{R}^d} \eta_m(\mathbf{r}) s(\mathbf{r}) d\mathbf{r}$$

Modality	Radiation	Analysis function $\eta_m$
2D or 3D tomography	coherent x-ray	$\delta(t_m - \langle \mathbf{r}, \theta_m \rangle)$ $\theta_m$ : mth sampled direction

## FORWARD IMAGING MODEL

In summary

$$[\mathbf{H}\mathbf{s}]_m = y_m = \langle \eta_m, \mathbf{s} \rangle \\ = \int_{\mathbb{R}^d} \eta_m(\mathbf{r}) s(\mathbf{r}) d\mathbf{r}$$

Modality	Radiation	Analysis function $\eta_m$
2D or 3D tomography	coherent x-ray	$\delta(t_m - \langle \mathbf{r}, \theta_m \rangle)$ $\theta_m$ : mth sampled direction
3D deconvolution microscopy	fluorescence	$h(\mathbf{r}_m - \mathbf{r})$ $h$ : PSF of microscope
Structured illumination microscopy (SIM)	fluorescence	$h(\mathbf{r}_m - \mathbf{r}) w_m(\mathbf{r})$ $w_m$ : mth illumination pattern
Positron emission tomography (PET)	gamma rays	$\delta(t_m - \langle \mathbf{r}, \theta_m \rangle)$ $(\theta_m, t_m)$ : mth LoR
Magnetic resonance imaging (MRI)	radio frequency	$e^{j\langle \omega_m, \mathbf{r} \rangle}$
Cardiac MRI (parallel, non-uniform)	radio frequency	$w_m(\mathbf{r}) e^{j\langle \omega_m, \mathbf{r} \rangle}$ $w_m$ : sensitivity of the mth coil

## FORWARD IMAGING MODEL

In summary

$$[\mathbf{H}\mathbf{s}]_m = y_m = \langle \eta_m, \mathbf{s} \rangle \\ = \int_{\mathbb{R}^d} \eta_m(\mathbf{r}) s(\mathbf{r}) d\mathbf{r}$$


Modality	Radiation	Analysis function $\eta_m$
2D or 3D tomography	coherent x-ray	$\delta(t_m - \langle \mathbf{r}, \theta_m \rangle)$ $\theta_m$ : mth sampled direction
3D deconvolution microscopy	fluorescence	$h(\mathbf{r}_m - \mathbf{r})$ $h$ : PSF of microscope
Structured illumination microscopy (SIM)	fluorescence	$h(\mathbf{r}_m - \mathbf{r}) w_m(\mathbf{r})$ $w_m$ : mth illumination pattern
Positron emission tomography (PET)	gamma rays	$\delta(t_m - \langle \mathbf{r}, \theta_m \rangle)$ $(\theta_m, t_m)$ : mth LoR
Magnetic resonance imaging (MRI)	radio frequency	$e^{j\langle \omega_m, \mathbf{r} \rangle}$
Cardiac MRI (parallel, non-uniform)	radio frequency	$w_m(\mathbf{r}) e^{j\langle \omega_m, \mathbf{r} \rangle}$ $w_m$ : sensitivity of the mth coil

How do we represent  $s \in L_2(\mathbb{R}^d)$  as  $\mathbf{s} \in \mathbb{R}^K$ ?

## DISCRETIZATION: FINITE DIMENSIONAL SIGNAL

Represent the continuous signal on a finite combination of synthesis functions  $\beta_k(\mathbf{r})$

$$s(\mathbf{r}) = \sum_{\mathbf{k} \in \Omega} s[\mathbf{k}] \beta_k(\mathbf{r}) \quad \text{with the signal vector (of coefficients): } \mathbf{s} = [s[\mathbf{k}]]_{\mathbf{k} \in \Omega} \text{ of dimension } K = |\Omega|.$$



8

## DISCRETIZATION: FINITE DIMENSIONAL SIGNAL


Represent the continuous signal on a finite combination of synthesis functions  $\beta_k(\mathbf{r})$

$$s(\mathbf{r}) = \sum_{\mathbf{k} \in \Omega} s[\mathbf{k}] \beta_k(\mathbf{r}) \quad \text{with the signal vector (of coefficients): } \mathbf{s} = [s[\mathbf{k}]]_{\mathbf{k} \in \Omega} \text{ of dimension } K = |\Omega|.$$

- Applying the forward imaging model to such a signal

$$\begin{aligned} y_m &= \int_{\mathbb{R}^d} s(\mathbf{r}) \eta_m(\mathbf{r}) d\mathbf{r} + n[m] = \langle s, \eta_m \rangle + n[m], \quad (m = 1, \dots, M) \\ &= \sum_{\mathbf{k} \in \Omega} \langle \beta_k, \eta_m \rangle s[\mathbf{k}] + n[m] \end{aligned}$$

$\eta_m$ : analysis function,  $\beta_k$ : synthesis function  
 $n[\cdot]$ : additive noise



8

## DISCRETIZATION: FINITE DIMENSIONAL SIGNAL


Represent the continuous signal on a finite combination of synthesis functions  $\beta_k(\mathbf{r})$

$$s(\mathbf{r}) = \sum_{\mathbf{k} \in \Omega} s[\mathbf{k}] \beta_k(\mathbf{r}) \quad \text{with the signal vector (of coefficients): } \mathbf{s} = [s[\mathbf{k}]]_{\mathbf{k} \in \Omega} \text{ of dimension } K = |\Omega|.$$

- Applying the forward imaging model to such a signal

$$\begin{aligned} y_m &= \int_{\mathbb{R}^d} s(\mathbf{r}) \eta_m(\mathbf{r}) d\mathbf{r} + n[m] = \langle s, \eta_m \rangle + n[m], \quad (m = 1, \dots, M) \\ &= \sum_{\mathbf{k} \in \Omega} \langle \beta_k, \eta_m \rangle s[\mathbf{k}] + n[m] \end{aligned}$$

$\eta_m$ : analysis function,  $\beta_k$ : synthesis function  
 $n[\cdot]$ : additive noise



8

## DISCRETIZATION: FINITE DIMENSIONAL SIGNAL

### Example basis functions

Shift-invariant representation:  $\beta_k(\mathbf{r}) = \beta(\mathbf{r} - \mathbf{k})$

$$\Rightarrow \mathbf{y} = \mathbf{H}\mathbf{s} + \mathbf{n} \quad (M \times K) \text{ system matrix : } [\mathbf{H}]_{m,k} = \langle \eta_m, \beta_k \rangle = \int_{\mathbb{R}^d} \eta_m(\mathbf{r}) \beta_k(\mathbf{r}) d\mathbf{r}$$

9

## DISCRETIZATION: FINITE DIMENSIONAL SIGNAL

### Example basis functions

Shift-invariant representation:  $\beta_k(\mathbf{r}) = \beta(\mathbf{r} - \mathbf{k})$

Separable generator:  $\beta(\mathbf{r}) = \prod_{n=1}^d \beta(\mathbf{r}[n])$

**Pixelated model**

## DISCRETIZATION: FINITE DIMENSIONAL SIGNAL

### Example basis functions

Shift-invariant representation:  $\beta_k(\mathbf{r}) = \beta(\mathbf{r} - \mathbf{k})$

Separable generator:  $\beta(\mathbf{r}) = \prod_{n=1}^d \beta(\mathbf{r}[n])$

**Pixelated model**

$$\beta(x) = \text{rect}(x)$$

9

9

## DISCRETIZATION: FINITE DIMENSIONAL SIGNAL

### Example basis functions

Shift-invariant representation:  $\beta_k(\mathbf{r}) = \beta(\mathbf{r} - \mathbf{k})$


Separable generator:  $\beta(\mathbf{r}) = \prod_{n=1}^d \beta(\mathbf{r}[n])$

**Pixelated model**

$$\beta(x) = \text{rect}(x)$$

**Bilinear model**

$$\beta(x) = (\text{rect} * \text{rect})(x) = \text{tri}(x)$$



## DISCRETIZATION: FINITE DIMENSIONAL SIGNAL

### Example basis functions

Shift-invariant representation:  $\beta_k(\mathbf{r}) = \beta(\mathbf{r} - \mathbf{k})$

Separable generator:  $\beta(\mathbf{r}) = \prod_{n=1}^d \beta(\mathbf{r}[n])$

**Pixelated model**



$$\beta(x) = \text{rect}(x)$$

**Bilinear model**

$$\beta(x) = (\text{rect} * \text{rect})(x) = \text{tri}(x)$$


**Bandlimited representation**

$$\beta(x) = \text{sinc}(x)$$



9

9



## A CONDENSED HISTORY OF IMAGE RECONSTRUCTION

See my 40 min tutorial at: [go.epfl.ch/ImRecTutorial](http://go.epfl.ch/ImRecTutorial)

## HISTORY OF IMAGE RECONSTRUCTION



© CIBM | Center for Biomedical Imaging

11

## HISTORY OF IMAGE RECONSTRUCTION

Classical image reconstruction  
(1st gen., 20th century)



Sparcity-based image reconstruction  
(2nd gen., 21st century)



The learning revolution  
(3rd gen.)



## HISTORY OF IMAGE RECONSTRUCTION



CIBM.CH

© CIBM | Center for Biomedical Imaging

11

© CIBM | Center for Biomedical Imaging

12

## HISTORY OF IMAGE RECONSTRUCTION



$$\text{Least-squares: } \min_{\tilde{s} \in \mathbb{R}_+^M} \{\|y - H\tilde{s}\|^2\}$$

## HISTORY OF IMAGE RECONSTRUCTION



$$\text{Least-squares: } \min_{\tilde{s} \in \mathbb{R}_+^M} \{\|y - H\tilde{s}\|^2\}$$

$$\min_{\tilde{s} \in \mathbb{R}_+^K} \underbrace{\|y - H\tilde{s}\|_2^2}_{\text{data consistency}} + \underbrace{\lambda \|L\tilde{s}\|_2^2}_{\text{regularization or Gaussian prior}}$$

## HISTORY OF IMAGE RECONSTRUCTION



$$\text{Least-squares: } \min_{\tilde{s} \in \mathbb{R}_+^M} \{\|y - H\tilde{s}\|^2\}$$

$$\min_{\tilde{s} \in \mathbb{R}_+^K} \underbrace{\|y - H\tilde{s}\|_2^2}_{\text{data consistency}} + \underbrace{\lambda \|L\tilde{s}\|_2^2}_{\text{regularization or Gaussian prior}}$$

"Filtered" backprojection

$$\Rightarrow s = (H^T H + \lambda L^T L)^{-1} H^T y$$

## HISTORY OF IMAGE RECONSTRUCTION



$$\text{Least-squares: } \min_{s \in \mathbb{R}_+^M} \{\|y - Hs\|^2\}$$

$$\min_{s \in \mathbb{R}_+^K} \underbrace{\|y - Hs\|_2^2}_{\text{data consistency}} + \underbrace{\lambda \|Ls\|_2^2}_{\text{regularization or Gaussian prior}}$$

"Filtered" backprojection

$$\Rightarrow s = (H^T H + \lambda L^T L)^{-1} H^T y$$

## HISTORY OF IMAGE RECONSTRUCTION



Tikhonov  
Wiener

Least-squares:  $\min_{\tilde{s} \in \mathbb{R}_+^M} \{\|y - H\tilde{s}\|_2^2\}$

$$\min_{\tilde{s} \in \mathbb{R}_+^K} \underbrace{\|y - H\tilde{s}\|_2^2}_{\text{data consistency}} + \underbrace{\lambda \|L\tilde{s}\|_2^2}_{\text{regularization or Gaussian prior}}$$

"Filtered" backprojection

$$\Rightarrow s = (H^T H + \lambda L^T L)^{-1} H^T y$$



$$\min_{s \in \mathbb{R}_+^K} \|y - Hs\|_2^2 + \lambda \mathcal{R}(s)$$

$$\mathcal{R}(s) = \|Ls\|_1 \text{ instead of } \|Ls\|_2^2$$

$$y = \begin{matrix} A \\ x \end{matrix} + n$$

$$s = L^{-1}x \quad A = HL^{-1}$$



C I B M . C H

## HISTORY OF IMAGE RECONSTRUCTION



Tikhonov  
Wiener

Least-squares:  $\min_{\tilde{s} \in \mathbb{R}_+^M} \{\|y - H\tilde{s}\|_2^2\}$

$$\min_{\tilde{s} \in \mathbb{R}_+^K} \underbrace{\|y - H\tilde{s}\|_2^2}_{\text{data consistency}} + \underbrace{\lambda \|L\tilde{s}\|_2^2}_{\text{regularization or Gaussian prior}}$$

"Filtered" backprojection

$$\Rightarrow s = (H^T H + \lambda L^T L)^{-1} H^T y$$



$$\min_{s \in \mathbb{R}_+^K} \|y - Hs\|_2^2 + \lambda \mathcal{R}(s)$$

$$\mathcal{R}(s) = \|Ls\|_1 \text{ instead of } \|Ls\|_2^2$$

$$y = \begin{matrix} A \\ x \end{matrix} + n$$

$$s = L^{-1}x \quad A = HL^{-1}$$



Dimensionality of parameters learnt

- $\lambda$
- Cal.  $H$

C I B M . C H

## HISTORY OF IMAGE RECONSTRUCTION



Tikhonov  
Wiener

Least-squares:  $\min_{\tilde{s} \in \mathbb{R}_+^M} \{\|y - H\tilde{s}\|_2^2\}$

$$\min_{\tilde{s} \in \mathbb{R}_+^K} \underbrace{\|y - H\tilde{s}\|_2^2}_{\text{data consistency}} + \underbrace{\lambda \|L\tilde{s}\|_2^2}_{\text{regularization or Gaussian prior}}$$

"Filtered" backprojection

$$\Rightarrow s = (H^T H + \lambda L^T L)^{-1} H^T y$$




$$\min_{s \in \mathbb{R}_+^K} \|y - Hs\|_2^2 + \lambda \mathcal{R}(s)$$

$$\mathcal{R}(s) = \|Ls\|_1 \text{ instead of } \|Ls\|_2^2$$

$$y = \begin{matrix} A \\ x \end{matrix} + n$$

C I B M . C H



## HISTORY OF IMAGE RECONSTRUCTION



Tikhonov  
Wiener

Least-squares:  $\min_{\tilde{s} \in \mathbb{R}_+^M} \{\|y - H\tilde{s}\|_2^2\}$

$$\min_{\tilde{s} \in \mathbb{R}_+^K} \underbrace{\|y - H\tilde{s}\|_2^2}_{\text{data consistency}} + \underbrace{\lambda \|L\tilde{s}\|_2^2}_{\text{regularization or Gaussian prior}}$$

"Filtered" backprojection

$$\Rightarrow s = (H^T H + \lambda L^T L)^{-1} H^T y$$



$$\min_{s \in \mathbb{R}_+^K} \|y - Hs\|_2^2 + \lambda \mathcal{R}(s)$$

$$\mathcal{R}(s) = \|Ls\|_1 \text{ instead of } \|Ls\|_2^2$$

$$y = \begin{matrix} A \\ x \end{matrix} + n$$




Dimensionality of parameters learnt

- $\lambda$
- L
- Cal. H
- $\mathcal{R}$

C I B M . C H


## HISTORY OF IMAGE RECONSTRUCTION



© CIBM | Center for Biomedical Imaging

12


## STABILITY OF IMAGE RECONSTRUCTION ALGORITHMS



## STABILITY OF IMAGE RECONSTRUCTION ALGORITHMS



### Learning-based reconstruction



© CIBM | Center for Biomedical Imaging


V. Antun, F. Remia, C. Poon, B. Adcock, and A.C. Hansen, "On instabilities of deep learning in image reconstruction and the potential costs of AI," *Proceedings of the National Academy of Sciences* 98 (117), pp. 30088–30095, 2020

14

## STABILITY OF IMAGE RECONSTRUCTION ALGORITHMS



### Learning-based reconstruction




V. Antun, F. Remia, C. Poon, B. Adcock, and A.C. Hansen, "On instabilities of deep learning in image reconstruction and the potential costs of AI," *Proceedings of the National Academy of Sciences* 98 (117), pp. 30088–30095, 2020

14

## STABILITY OF IMAGE RECONSTRUCTION ALGORITHMS



### Learning-based reconstruction



© CIBM | Center for Biomedical Imaging

V. Antun, F. Renia, C. Poon, B. Adcock, and A.C. Hansen, "On instabilities of deep learning in image reconstruction and the potential costs of AI," *Proceedings of the National Academy of Sciences* 115 (17), pp. 30988–30995, 2018.


C I B M . C H

14

## STABILITY OF IMAGE RECONSTRUCTION ALGORITHMS



### Learning-based reconstruction



© CIBM | Center for Biomedical Imaging

V. Antun, F. Renia, C. Poon, B. Adcock, and A.C. Hansen, "On instabilities of deep learning in image reconstruction and the potential costs of AI," *Proceedings of the National Academy of Sciences* 115 (17), pp. 30988–30995, 2018.

C I B M . C H

14

## STABILITY OF IMAGE RECONSTRUCTION ALGORITHMS



### Sparsity-based reconstruction

#### Stability of the reconstruction



Continuity of the solution map

e.g., Lipschitz continuity:

$$\frac{\|\mathbf{s}_{\mathbf{y}_1} - \mathbf{s}_{\mathbf{y}_2}\|}{\|\mathbf{y}_1 - \mathbf{y}_2\|}$$

© CIBM | Center for Biomedical Imaging

C I B M . C H

15

## STABILITY OF IMAGE RECONSTRUCTION ALGORITHMS



### Sparsity-based reconstruction


#### Stability of the reconstruction



Continuity of the solution map

e.g., Lipschitz continuity:

$$\frac{\|\mathbf{s}_{\mathbf{y}_1} - \mathbf{s}_{\mathbf{y}_2}\|}{\|\mathbf{y}_1 - \mathbf{y}_2\|}$$



M. Genzel, J. Macdonald, and M. Marz, "Solving inverse problems with deep neural networks – Robustness included," *IEEE Transactions on Pattern Analysis and Machine Learning – Early Access* (2023)


C I B M . C H

15

## STABILITY OF IMAGE RECONSTRUCTION ALGORITHMS



### Sparsity-based reconstruction



**Figure 3: Both trained and un-trained reconstruction methods are vulnerable to small adversarial perturbations.**  
Performance loss as a function of the perturbation strength,  $\epsilon = \frac{\|\text{perturbation}\|_2}{\|\text{k-space}\|_2}$ , for all methods. In each plot, the perturbations are obtained by attacking one method (specified in the plot title), and are applied to all methods. The results are averaged over 10 randomly-chosen proton density knee images from the fastMRI validation set. Shaded areas denote 95% confidence intervals.



## DISCRETIZATION: REVISITED

Exact discretization  $\Leftrightarrow$  Representer theorems for optimization in abstract spaces

### ■ Very general setting

$$s \in L_2(\mathbb{R}^d) \text{ turns into } f \in \mathcal{X}'$$

$$\min_{f \in \mathcal{X}'} \{E(\mathbf{y}, \boldsymbol{\nu} f) + \psi(\|f\|_{\mathcal{X}'})\}$$

- $\mathcal{X}$ : reflexive Banach space w. dual  $\mathcal{X}'$

## DISCRETIZATION: REVISITED

Exact discretization  $\Leftrightarrow$  Representer theorems for optimization in abstract spaces

### ■ Very general setting

$$s \in L_2(\mathbb{R}^d) \text{ turns into } f \in \mathcal{X}'$$

$$\min_{f \in \mathcal{X}'} \{E(\mathbf{y}, \boldsymbol{\nu} f) + \psi(\|f\|_{\mathcal{X}'})\}$$

## DISCRETIZATION: REVISITED

Exact discretization  $\Leftrightarrow$  Representer theorems for optimization in abstract spaces

### ■ Very general setting

$$s \in L_2(\mathbb{R}^d) \text{ turns into } f \in \mathcal{X}'$$

$$\min_{f \in \mathcal{X}'} \{E(\mathbf{y}, \boldsymbol{\nu} f) + \psi(\|f\|_{\mathcal{X}'})\}$$

- $\mathcal{X}$ : reflexive Banach space w. dual  $\mathcal{X}'$

- $\mathbf{y} \in \mathbb{R}^M$ : measurements

## DISCRETIZATION: REVISITED

Exact discretization  $\Leftrightarrow$  Representer theorems for optimization in abstract spaces

- Very general setting

$$s \in L_2(\mathbb{R}^d) \text{ turns into } f \in \mathcal{X}'$$

$$\min_{f \in \mathcal{X}'} \{E(\mathbf{y}, \boldsymbol{\nu} f) + \psi(\|f\|_{\mathcal{X}'})\}$$

- $\mathcal{X}$ : reflexive Banach space w. dual  $\mathcal{X}'$
- $\mathbf{y} \in \mathbb{R}^M$ : measurements
- $\boldsymbol{\nu}: \mathcal{X}' \rightarrow \mathbb{R}^M$ : forward model
- $\boldsymbol{\nu} = (\nu_1, \dots, \nu_M)$

M. Unser "A unifying representer theorem for inverse problems and machine learning,"  
Foundations of Computational Mathematics (2021), in press.

17

## DISCRETIZATION: REVISITED

Exact discretization  $\Leftrightarrow$  Representer theorems for optimization in abstract spaces

- Very general setting

$$s \in L_2(\mathbb{R}^d) \text{ turns into } f \in \mathcal{X}'$$

$$\min_{f \in \mathcal{X}'} \{E(\mathbf{y}, \boldsymbol{\nu} f) + \psi(\|f\|_{\mathcal{X}'})\}$$

- $\mathcal{X}$ : reflexive Banach space w. dual  $\mathcal{X}'$
- $\mathbf{y} \in \mathbb{R}^M$ : measurements
- $\boldsymbol{\nu}: \mathcal{X}' \rightarrow \mathbb{R}^M$ : forward model
- $\boldsymbol{\nu} = (\nu_1, \dots, \nu_M)$
- $E$  and  $\psi$  are nice

M. Unser "A unifying representer theorem for inverse problems and machine learning,"  
Foundations of Computational Mathematics (2021), in press.

17

## DISCRETIZATION: REVISITED

Exact discretization  $\Leftrightarrow$  Representer theorems for optimization in abstract spaces

- Very general setting

$$s \in L_2(\mathbb{R}^d) \text{ turns into } f \in \mathcal{X}'$$

$$\min_{f \in \mathcal{X}'} \{E(\mathbf{y}, \boldsymbol{\nu} f) + \psi(\|f\|_{\mathcal{X}'})\}$$

- $\mathcal{X}$ : reflexive Banach space w. dual  $\mathcal{X}'$
- $\mathbf{y} \in \mathbb{R}^M$ : measurements
- $\boldsymbol{\nu}: \mathcal{X}' \rightarrow \mathbb{R}^M$ : forward model
- $\boldsymbol{\nu} = (\nu_1, \dots, \nu_M)$
- $E$  and  $\psi$  are nice

M. Unser "A unifying representer theorem for inverse problems and machine learning,"  
Foundations of Computational Mathematics (2021), in press.

17

## DISCRETIZATION: REVISITED

Exact discretization  $\Leftrightarrow$  Representer theorems for optimization in abstract spaces

- Very general setting

$$s \in L_2(\mathbb{R}^d) \text{ turns into } f \in \mathcal{X}'$$

$$\exists! \mathbf{a}_y \in \mathbb{R}^M$$

$$\min_{f \in \mathcal{X}'} \{E(\mathbf{y}, \boldsymbol{\nu} f) + \psi(\|f\|_{\mathcal{X}'})\}$$

$$\implies f_y = \mathcal{J}_{\mathcal{X}} \left( \sum_{m=1}^M a_{y,m} \nu_m \right)$$

- $\mathcal{X}$ : reflexive Banach space w. dual  $\mathcal{X}'$
- $\mathbf{y} \in \mathbb{R}^M$ : measurements
- $\boldsymbol{\nu}: \mathcal{X}' \rightarrow \mathbb{R}^M$ : forward model
- $\boldsymbol{\nu} = (\nu_1, \dots, \nu_M)$
- $E$  and  $\psi$  are nice

M. Unser "A unifying representer theorem for inverse problems and machine learning,"  
Foundations of Computational Mathematics (2021), in press.

17

## DISCRETIZATION: REVISITED

Exact discretization  $\Leftrightarrow$  Representer theorems for optimization in abstract spaces

- Very general setting

$s \in L_2(\mathbb{R}^d)$  turns into  $f \in \mathcal{X}'$

$$\min_{f \in \mathcal{X}'} \{E(\mathbf{y}, \nu f) + \psi(\|f\|_{\mathcal{X}'})\}$$

- $\mathcal{X}$ : reflexive Banach space w. dual  $\mathcal{X}'$
- $\mathbf{y} \in \mathbb{R}^M$ : measurements
- $\nu: \mathcal{X}' \rightarrow \mathbb{R}^M$ : forward model
- $\nu = (\nu_1, \dots, \nu_M)$
- $E$  and  $\psi$  are nice

- $f_y$ : reconstruction for measurements  $\mathbf{y}$

$$\exists! \mathbf{a}_y \in \mathbb{R}^M$$

$$\implies f_y = \mathcal{J}_{\mathcal{X}} \left( \sum_{m=1}^M a_{y,m} \nu_m \right)$$

- $\mathcal{J}_{\mathcal{X}}: \mathcal{X} \rightarrow \mathcal{X}'$ : non-linear "duality map"

### Exact discretization

M. Unser "A unifying representer theorem for inverse problems and machine learning,"

Foundations of Computational Mathematics (2021), in press.

17

## STABILITY OF IMAGE RECONSTRUCTION ALGORITHMS



Classical image reconstruction: Tikhonov regularization

- $\mathcal{X}$ : Hilbert space



© CIBM | Center for Biomedical Imaging

18

## STABILITY OF IMAGE RECONSTRUCTION ALGORITHMS



Classical image reconstruction: Tikhonov regularization

- $\mathcal{X}$ : Hilbert space  $\Rightarrow \mathcal{J}_{\mathcal{X}}$  is linear

## STABILITY OF IMAGE RECONSTRUCTION ALGORITHMS



Classical image reconstruction: Tikhonov regularization

- $\mathcal{X}$ : Hilbert space  $\Rightarrow \mathcal{J}_{\mathcal{X}}$  is linear  $\implies f_y = \sum_{m=1}^M a_{y,m} \varphi_m$  where  $\varphi_m = \mathcal{J}_{\mathcal{X}}\{\nu_m\}$



© CIBM | Center for Biomedical Imaging

18



© CIBM | Center for Biomedical Imaging

18

## STABILITY OF IMAGE RECONSTRUCTION ALGORITHMS



Classical image reconstruction: Tikhonov regularization

- $\mathcal{X}$ : Hilbert space  $\Rightarrow \mathcal{J}_{\mathcal{X}}$  is linear  $\Rightarrow f_{\mathbf{y}} = \sum_{m=1}^M a_{\mathbf{y},m} \varphi_m$  where  $\varphi_m = \mathcal{J}_{\mathcal{X}}\{\nu_m\}$
- For the simplified case:  $E(\mathbf{y}, \tilde{\mathbf{y}}) = \frac{1}{2}\|\mathbf{y} - \tilde{\mathbf{y}}\|^2$  and  $\psi(x) = \lambda x^2$



## STABILITY OF IMAGE RECONSTRUCTION ALGORITHMS

Classical image reconstruction: Tikhonov regularization

- $\mathcal{X}$ : Hilbert space  $\Rightarrow \mathcal{J}_{\mathcal{X}}$  is linear  $\Rightarrow f_{\mathbf{y}} = \sum_{m=1}^M a_{\mathbf{y},m} \varphi_m$  where  $\varphi_m = \mathcal{J}_{\mathcal{X}}\{\nu_m\}$
- For the simplified case:  $E(\mathbf{y}, \tilde{\mathbf{y}}) = \frac{1}{2}\|\mathbf{y} - \tilde{\mathbf{y}}\|^2$  and  $\psi(x) = \lambda x^2 \Rightarrow \mathbf{a}_{\mathbf{y}} = (\mathbf{H} + 2\lambda \mathbf{Id})^{-1}\mathbf{y}$



## STABILITY OF IMAGE RECONSTRUCTION ALGORITHMS



Classical image reconstruction: Tikhonov regularization

- $\mathcal{X}$ : Hilbert space  $\Rightarrow \mathcal{J}_{\mathcal{X}}$  is linear  $\Rightarrow f_{\mathbf{y}} = \sum_{m=1}^M a_{\mathbf{y},m} \varphi_m$  where  $\varphi_m = \mathcal{J}_{\mathcal{X}}\{\nu_m\}$
- For the simplified case:  $E(\mathbf{y}, \tilde{\mathbf{y}}) = \frac{1}{2}\|\mathbf{y} - \tilde{\mathbf{y}}\|^2$  and  $\psi(x) = \lambda x^2 \Rightarrow \mathbf{a}_{\mathbf{y}} = (\mathbf{H} + 2\lambda \mathbf{Id})^{-1}\mathbf{y}$

$$\begin{aligned} \|f_{\mathbf{y}_1} - f_{\mathbf{y}_2}\|_{\mathcal{H}}^2 &= (\mathbf{a}_{\mathbf{y}_1} - \mathbf{a}_{\mathbf{y}_2})^T \mathbf{H} (\mathbf{a}_{\mathbf{y}_1} - \mathbf{a}_{\mathbf{y}_2}) \\ &= (\mathbf{y}_1 - \mathbf{y}_2)^T (\mathbf{H} + 2\lambda \mathbf{Id})^{-T} \mathbf{H} (\mathbf{H} + 2\lambda \mathbf{Id})^{-1} (\mathbf{y}_1 - \mathbf{y}_2) \\ &= (\mathbf{y}_1 - \mathbf{y}_2)^T \mathbf{P} \frac{\Lambda}{(\Lambda + 2\lambda \mathbf{Id})^2} \mathbf{P}^T (\mathbf{y}_1 - \mathbf{y}_2) \\ &\leq \max_{m \in \{1, 2, \dots, M\}} \left\{ \frac{\sigma_m}{(\sigma_m + 2\lambda)^2} \right\} \|\mathbf{y}_1 - \mathbf{y}_2\|_2^2, \end{aligned}$$



## STABILITY OF IMAGE RECONSTRUCTION ALGORITHMS



Classical image reconstruction: Tikhonov regularization

- $\mathcal{X}$ : Hilbert space  $\Rightarrow \mathcal{J}_{\mathcal{X}}$  is linear  $\Rightarrow f_{\mathbf{y}} = \sum_{m=1}^M a_{\mathbf{y},m} \varphi_m$  where  $\varphi_m = \mathcal{J}_{\mathcal{X}}\{\nu_m\}$
- For the simplified case:  $E(\mathbf{y}, \tilde{\mathbf{y}}) = \frac{1}{2}\|\mathbf{y} - \tilde{\mathbf{y}}\|^2$  and  $\psi(x) = \lambda x^2 \Rightarrow \mathbf{a}_{\mathbf{y}} = (\mathbf{H} + 2\lambda \mathbf{Id})^{-1}\mathbf{y}$

$$\begin{aligned} \|f_{\mathbf{y}_1} - f_{\mathbf{y}_2}\|_{\mathcal{H}}^2 &= (\mathbf{a}_{\mathbf{y}_1} - \mathbf{a}_{\mathbf{y}_2})^T \mathbf{H} (\mathbf{a}_{\mathbf{y}_1} - \mathbf{a}_{\mathbf{y}_2}) \\ &= (\mathbf{y}_1 - \mathbf{y}_2)^T (\mathbf{H} + 2\lambda \mathbf{Id})^{-T} \mathbf{H} (\mathbf{H} + 2\lambda \mathbf{Id})^{-1} (\mathbf{y}_1 - \mathbf{y}_2) \\ &= (\mathbf{y}_1 - \mathbf{y}_2)^T \mathbf{P} \frac{\Lambda}{(\Lambda + 2\lambda \mathbf{Id})^2} \mathbf{P}^T (\mathbf{y}_1 - \mathbf{y}_2) \\ &\leq \max_{m \in \{1, 2, \dots, M\}} \left\{ \frac{\sigma_m}{(\sigma_m + 2\lambda)^2} \right\} \|\mathbf{y}_1 - \mathbf{y}_2\|_2^2, \end{aligned}$$

Lipschitz continuity



## STABILITY OF IMAGE RECONSTRUCTION ALGORITHMS




Classical image reconstruction: Tikhonov regularization



## STABILITY OF IMAGE RECONSTRUCTION ALGORITHMS





Classical image reconstruction: Tikhonov regularization



## STABILITY OF IMAGE RECONSTRUCTION ALGORITHMS





Sparsity-based reconstruction



## STABILITY OF IMAGE RECONSTRUCTION ALGORITHMS




Sparsity-based reconstruction



## STABILITY OF IMAGE RECONSTRUCTION ALGORITHMS



Sparsity-based reconstruction



© CIBM | Center for Biomedical Imaging

E. J. Candès, J. K. Romberg, and T. Tao "Stable signal recovery from incomplete and inaccurate measurements." Communications on Pure and Applied Mathematics 59 (8), pp. 1207–1223, 2006


C I B M . C H

20

## STABILITY OF IMAGE RECONSTRUCTION ALGORITHMS



Sparsity-based reconstruction



© CIBM | Center for Biomedical Imaging

E. J. Candès, J. K. Romberg, and T. Tao "Stable signal recovery from incomplete and inaccurate measurements." Communications on Pure and Applied Mathematics 59 (8), pp. 1207–1223, 2006

C I B M . C H

20

## STABILITY OF IMAGE RECONSTRUCTION ALGORITHMS



$L_p$ -regularization

$$\mathcal{R}(\mathbf{s}) = \|\mathbf{L}\mathbf{s}\|_p^p \Leftrightarrow \mathcal{X}' = L_p(\Omega) \text{ and } \psi(x) = \lambda x^p$$



© CIBM | Center for Biomedical Imaging

C I B M . C H


21

## STABILITY OF IMAGE RECONSTRUCTION ALGORITHMS



$L_p$ -regularization

$$\mathcal{R}(\mathbf{s}) = \|\mathbf{L}\mathbf{s}\|_p^p \Leftrightarrow \mathcal{X}' = L_p(\Omega) \text{ and } \psi(x) = \lambda x^p$$



$$\|f_{\mathbf{y}_1} - f_{\mathbf{y}_2}\|_{L_p} \leq \frac{(2r_p(Y))^{2-p}K_p}{\lambda p(p-1)} \|\mathbf{y}_1 - \mathbf{y}_2\|_2$$

© CIBM | Center for Biomedical Imaging

C I B M . C H


21

## STABILITY OF IMAGE RECONSTRUCTION ALGORITHMS



$L_p$ -regularization

$$\mathcal{R}(\mathbf{s}) = \|\mathbf{L}\mathbf{s}\|_p^p \Leftrightarrow \mathcal{X}' = L_p(\Omega) \text{ and } \psi(x) = \lambda x^p$$




## STABILITY OF IMAGE RECONSTRUCTION ALGORITHMS



$L_p$ -regularization

$$\mathcal{R}(\mathbf{s}) = \|\mathbf{L}\mathbf{s}\|_p^p \Leftrightarrow \mathcal{X}' = L_p(\Omega) \text{ and } \psi(x) = \lambda x^p$$



THANK YOU FOR YOUR ATTENTION

