Towards a Formal Definition of Kinase Inhibitor Selectivity: Empirical Characterization of Definitional Instability and Proposed Desiderata

Polina Vinogradova*

Independent Researcher, Ottawa, Canada

E-mail: polina@example.com

Abstract

Kinase inhibitor selectivity is routinely quantified using one of several definitions — including the S-score, selectivity entropy, Gini coefficient, and ratio-based measures — yet these definitions are applied interchangeably without systematic justification. We characterize, across two large kinase inhibitor profiling datasets (Davis, $n = 68$; Klaeger, $n = 222$), the extent and structural sources of disagreement between definitions as their parameters vary. Spearman correlations between median rankings reveal that ratio-based definitions cluster separately from entropy- and distribution-based definitions ($r = 0.27$–0.48 vs. $r = 0.75$–0.91). Rank instability is not random: three mechanistically distinct sources are identified. Drugs with no detected binding above a minimum affinity threshold exhibit maximal instability (mean rank $\sigma \approx 74$) due to definitional noise. Among active compounds, near-tied top targets drive ratio-specific instability ($r = -0.293$, $p < 0.001$), while broad flat binding profiles drive instability within distribution-based families. Attempts to correlate selectivity scores with clinical adverse event rates from FAERS ($n = 46$) and FDA label discontinuation rates ($n = 33$) yield null results, which we attribute to confounding by indication severity and prescription volume. From these findings we derive four desiderata toward a well-formed selectivity measure: a reliability threshold condition, bounded gap sensitivity, distributional consistency under parameter perturbation, and monotonicity under weak off-target addition. These desiderata are intended as the empirical foundation for a future formal axiomatization, in the tradition of Shannon's characterization of entropy.

Introduction

Kinase inhibitors are among the most intensely investigated compound classes in drug discovery, with over 80 FDA-approved agents and several hundred in clinical development.[1] Because the human kinome comprises more than 500 proteins sharing a highly conserved ATP-binding site,[2] the design of inhibitors with controlled target profiles remains a central challenge. Selectivity — the degree to which a compound's binding activity is concentrated on intended rather than off-target kinases — is a primary criterion in lead optimization and a key determinant of the therapeutic window of approved drugs.[3]

Despite its importance, selectivity is not formally defined. The field employs at least four distinct quantitative measures: the S-score,[4] selectivity entropy,[5] Gini coefficient,[6] and ratio-based measures including the CATDS score.[7] These definitions are applied interchangeably in the literature without systematic justification, and their parameter values — the activity threshold in the S-score, the baseline in the entropy and Gini coefficient, the reference off-target in ratio-based measures — are chosen by convention rather than principle. Bosc et al.[8] showed that these definitions can produce substantially different compound rankings on the same dataset, and the KInhibition platform[9] explicitly acknowledged that conflicting results across metrics are common. However, the structural sources of this disagreement — which binding profile properties determine when definitions agree or conflict, and why — have not been characterized.

The absence of a principled basis for selectivity measurement is not merely a practical inconvenience. Go/no-go decisions worth hundreds of millions of dollars in development costs are made on the basis of selectivity assessments whose sensitivity to definitional choice is unknown. When a compound is described as "selective" in the literature, the claim is implicitly conditioned on a specific definition and parameterization that is rarely stated explicitly and almost never justified. This situation is analogous to measuring temperature with instruments whose scales have not been calibrated to a common reference: the measurements are internally consistent but not directly comparable.

The analogy to measurement theory suggests a remedy. In information theory, Shannon[10] resolved the question of how to measure uncertainty by identifying axioms that uniquely characterize the entropy functional. In social choice theory, Arrow[11] resolved — negatively — the question of how to aggregate individual rankings by proving that no aggregation rule can simultaneously satisfy a small set of reasonable requirements. In both cases, the path from empirical observation to formal resolution required first characterizing the behavior of competing approaches and then identifying the properties that distinguish good measures from bad ones.

This paper takes the first step of that program for kinase inhibitor selectivity. We systematically characterize, across two large profiling datasets, the extent and structural sources of disagreement between existing definitions as their parameters vary. We identify three mechanistically distinct sources of rank instability and show that instability is predictable from the shape of a compound's binding profile. We then propose four desiderata — necessary properties of any well-formed selectivity measure — motivated directly by these failure modes.

Specifically, the contributions of this work are as follows. First, we show that existing definitions cluster into two families — distribution-based (S-score, entropy, Gini) and ratio-based (CATDS and variants) — that correlate poorly with one another ($r = 0.27$–0.48) and measure fundamentally different properties of binding profiles. Second, we identify three sources of instability: definitional noise for compounds below the assay detection threshold, ratio-specific instability driven by near-tied top targets, and distribution-ratio disagreement for compounds with one dominant target and broad moderate off-target activity. Third, we propose four desiderata — a reliability threshold condition, bounded gap sensitivity, distributional consistency, and monotonicity under weak off-target addition — and show that no existing definition satisfies all four simultaneously. Fourth, we demonstrate that correlations between selectivity scores and clinical adverse event outcomes are null under current data sources, and identify the confounders responsible.

The desiderata proposed here are intended as the empirical foundation for a future formal

axiomatization of kinase inhibitor selectivity, in the tradition of Shannon's characterization of entropy and Arrow's analysis of social choice functions.

Related Work

Existing Selectivity Metrics and Their Comparison

Several metrics have been proposed to quantify kinase inhibitor selectivity from profiling panel data. Karaman et al.[4] introduced the selectivity score (S-score), defined as the fraction of panel kinases inhibited above a fixed concentration threshold. Uitdehaag and Zaman[5] proposed the selectivity entropy, which treats the normalized binding profile as a probability distribution and computes its Shannon entropy; low entropy indicates activity concentrated on few targets. Graczyk[6] defined a thermodynamics-based partition index estimating the fraction of total binding directed to a reference kinase, suited primarily to hit-to-lead optimization against a specified target. Klaeger et al.[7] introduced the concentration- and target-dependent selectivity (CATDS) score, which quantifies the reduction in binding to a given target relative to the summed reduction across all detected targets at a specified inhibitor concentration. Bosc et al.[8] subsequently proposed two further metrics — the window score and ranking score — and performed a comparative analysis of all five definitions on the Davis and Anastassiadis datasets, showing that compound rankings differ substantially depending on the metric applied and that certain compounds cannot be scored under some definitions when threshold conditions are not satisfied.

Reconciliation attempts have remained limited in scope. Bajorath et al.[12] compared CATDS-based selectivity assessments from Klaeger et al. with activity data mined from ChEMBL for the same compounds, finding broad consistency between the two sources but noting that no simple relationship between selectivity and clinical performance was apparent. The KInhibition platform[9] explicitly acknowledged that multiple metrics yield conflicting results for the same compound but addressed this through a unified query interface rather than theoretical analysis.

A common limitation of these studies is that each metric is evaluated at fixed parameter values. The sensitivity of rankings to parameter choice — the activity threshold in the

S-score, the baseline in the entropy and Gini coefficient, the reference off-target in ratio-based measures — has not been systematically characterized. The structural properties of binding profiles that determine when definitions agree or disagree have not been identified. The present work addresses both gaps.

Axiomatic Approaches to Ranking and Measurement

Axiomatic characterization of measurement functions is well-established in adjacent fields. Shannon[10] demonstrated that the entropy of a probability distribution is uniquely determined, up to a multiplicative constant, by four requirements: continuity, maximality on the uniform distribution, expansibility, and additivity over independent subsystems. Khinchin[13] subsequently strengthened this uniqueness result. The selectivity entropy of Uitdehaag and Zaman applies the Shannon formula to normalized binding profiles but provides no analogous axiomatic justification; it is not known whether it is the unique function satisfying reasonable selectivity-specific desiderata, or one of many equivalent alternatives.

Arrow[11] established that no social welfare function satisfies simultaneously the requirements of unrestricted domain, Pareto efficiency, independence of irrelevant alternatives, and non-dictatorship. The theorem is relevant here in two respects: it demonstrates that axiomatic analysis of a ranking concept can yield impossibility rather than uniqueness, and it shows that competing ranking functions reflect genuine tradeoffs between properties that cannot all be achieved simultaneously. Pavillidis et al.[14] applied this methodology to truth discovery — the problem of identifying true facts from conflicting reports of unknown reliability — posing it as a social choice problem, formulating candidate axioms, and proving that a subset cannot be jointly satisfied. The structure of that problem is closely analogous to selectivity measurement, where multiple definitions produce conflicting rankings and the question of which to trust is unresolved.

Formal methods have been applied to biological systems primarily in the context of process dynamics. Cardelli[15] used the pi-calculus to model molecular biology, and the

Beacon Calculus[16] addresses stochastic reaction networks; both verify properties of temporal execution rather than static measurements. Type-theoretic approaches to scientific units of measure, including Kennedy's[17] F# units-of-measure system and ongoing formalization in Lean 4 Mathlib,[18] enforce dimensional consistency for physical quantities but do not extend to biochemical properties. To our knowledge, axiomatic analysis has not previously been applied to any selectivity or promiscuity measure in drug discovery. The present work proposes empirically motivated desiderata as a foundation for future formal axiomatization.

Methods

Datasets

Two publicly available kinase inhibitor profiling datasets were used. The Davis dataset[19] comprises binding affinities (K_d) for 68 small-molecule kinase inhibitors measured against 433 human kinase domains. Affinity values are reported as dissociation constants and were converted to $\mathrm{p}K_d = -\log_{10}(K_d/10^9)$ prior to analysis, following the convention of the DeepDTA benchmark.[20] Pairs for which no binding was detected at the screening concentration of 10 μM were assigned $\mathrm{p}K_d = 5.0$, corresponding to a K_d of 1 mM, which represents the detection floor of the assay. The resulting affinity matrix is complete (68×433, 29,444 entries). No censored or greater-than values are present in the Davis dataset: the assay reports a quantitative K_d for every compound–kinase pair where activity was detected at 10 μM, and assigns the detection floor value otherwise. All non-binding pairs are therefore treated as exactly equivalent at $\mathrm{p}K_d = 5.0$, which slightly underestimates true non-binding affinity but is the standard convention for this dataset.[8,20]

The Klaeger dataset[7] comprises binding affinities for 243 clinical kinase inhibitors profiled against the human kinome using Kinobeads chemical proteomics. Apparent dissociation constants (K_d^{app}) were derived from dose–response curves across eight inhibitor concentrations in cancer cell lysates. Only high-confidence measurements (as annotated by the original authors) were retained, yielding 5,123 drug–target pairs for 222 drugs and 343 kinase targets. Affinity values were converted to $\mathrm{p}K_d$ using the same formula as above. Missing entries, representing kinase–inhibitor combinations not detected in the assay, were assigned the background value $\mathrm{p}K_d = 5.0$. The resulting matrix (222×343) has 17.9% active entries (pKd > 6, i.e. $K_d < 1$ μM) for Davis and 3.6% for Klaeger, reflecting the more stringent detection threshold of the chemoproteomic assay.

Selectivity Definitions and Parameterization

Four selectivity definitions were implemented, each controlled by one or more parameters whose values are not standardized in the literature.

The **S-score**[4] is defined as

$$S(\tau) = \frac{1}{N} \sum_{i=1}^{N} \mathbf{1}[\mathrm{p}K_{d,i} > \tau], \tag{1}$$

where N is the number of kinases in the panel and τ is an activity threshold. Lower values indicate greater selectivity. We swept $\tau \in \{5.5, 5.75, \ldots, 8.0\}$ (11 values), spanning K_d cutoffs from approximately 300 nM to 10 nM.

The **selectivity entropy**[5] treats the baseline-subtracted binding profile as a probability distribution:

$$H(\beta) = -\sum_{i=1}^{N} p_i \log_2 p_i, \quad p_i = \frac{\max(\mathrm{p}K_{d,i} - \beta,\, 0)}{\sum_j \max(\mathrm{p}K_{d,j} - \beta,\, 0)}, \tag{2}$$

where β is a baseline affinity below which binding is considered negligible. Lower entropy indicates greater selectivity. We swept $\beta \in \{5.0, 5.25, \ldots, 6.5\}$ (7 values).

The **Gini coefficient**[6] of the baseline-subtracted profile is

$$G(\beta) = \frac{2 \sum_{i=1}^{N} i \cdot x_{(i)}}{N \sum_{i=1}^{N} x_{(i)}} - \frac{N+1}{N}, \tag{3}$$

where $x_{(i)} = \max(\mathrm{p}K_{d,i} - \beta,\, 0)$ sorted in ascending order. Higher values indicate greater selectivity. We swept β over the same range as for the entropy.

The **ratio** definition measures the gap between the primary target and a specified off-target:

$$R(k) = \mathrm{p}K_{d,(1)} - \mathrm{p}K_{d,(k+1)}, \tag{4}$$

where $\mathrm{p}K_{d,(j)}$ denotes the j-th largest affinity in the profile and $k \in \{1, 2, 3, 4, 5\}$ indexes the off-target used as reference. Higher values indicate greater selectivity.

Rank Stability Analysis

For each definition and each parameter value, compounds were ranked from most selective (rank 1) to least selective (rank n, where n is the number of compounds in the dataset). Ties were broken by average rank. This yielded 30 rank vectors per dataset: 11 for the S-score, 7 for entropy, 7 for Gini, and 5 for ratio.

For each compound, rank instability was quantified as the standard deviation of its rank across all 30 configurations (overall instability) and separately across configurations within each definition family (within-family instability). Cross-definition agreement was assessed using Spearman rank correlation between the median rank vectors of each definition family. The median was taken across parameter values within each family prior to correlation to obtain a single representative ranking per definition.

Associations between rank instability and binding profile features were assessed using Spearman correlation. Profile features computed for each compound included: the number of active kinases ($\mathrm{p}K_d > 6$), the gap between the top and second-ranked affinities, the standard deviation and range of active affinities, and the absolute affinity of the second-ranked target. All correlations are reported with two-tailed p-values; significance thresholds of $p < 0.05$, $p < 0.01$, and $p < 0.001$ are denoted *, **, and *** respectively. Analyses were performed in Python 3.10 using NumPy, pandas, and SciPy.[21–23]

Clinical Outcome Data

For the subset of Klaeger compounds that are FDA-approved ($n = 46$ after excluding drugs with no FAERS records), adverse event report counts were retrieved from the FDA Adverse Event Reporting System (FAERS) via the openFDA API. Total report counts, serious event counts, and death and hospitalization subcounts were retrieved per drug. Serious event rates and death rates were computed by normalizing subcounts by total reports per drug.

Treatment discontinuation rates due to adverse events were extracted from FDA drug labels retrieved via the openFDA label API for 33 approved drugs with available adverse

reactions sections. Label text was reviewed manually to identify the primary clinical trial discontinuation rate attributable to adverse reactions.

Associations between selectivity scores and each clinical outcome measure were assessed using Spearman correlation across drugs with available data.

Results

Selectivity Definitions Cluster into Two Families

Across 30 parameter configurations (11 S-score thresholds, 7 entropy baselines, 7 Gini baselines, 5 ratio off-target indices), compound rankings were computed for all drugs in both datasets. Spearman correlations between the median ranking of each definition family reveal a consistent two-cluster structure (Table 1). On the Davis dataset, S-score, entropy, and Gini are mutually correlated ($r = 0.91$–0.999), while ratio correlates substantially less with each ($r = 0.34$–0.45). On the larger Klaeger dataset, the same structure is preserved but with lower within-cluster correlations ($r = 0.75$–0.91), reflecting greater diversity of binding profiles among the 222 clinical compounds. The ratio definition is the consistent outlier in both datasets, with cross-family correlations of $r = 0.27$–0.62.

Table 1: Spearman correlations between median rankings by definition family. Upper triangle: Davis ($n = 68$). Lower triangle: Klaeger ($n = 222$).

	S-score	Entropy	Gini	Ratio
S-score	—	0.949	0.954	0.454
Entropy	0.910	—	0.999	0.343
Gini	0.745	0.869	—	0.343
Ratio	0.265	0.480	0.617	—

This two-family structure reflects a fundamental difference in what the definitions measure. The S-score, entropy, and Gini all quantify the concentration of activity across the full kinase panel; they differ in functional form but respond to the same underlying property of the binding profile. The ratio definition, by contrast, quantifies the gap between the primary target and a specific off-target, ignoring the remainder of the binding profile entirely. A compound that strongly inhibits one kinase and moderately inhibits twenty others will be ranked as highly selective by ratio (large gap to the second target) but as moderately promiscuous by entropy and Gini (activity distributed across many targets).

Rank Instability Is Structured and Predictable



Figure 1: Best predictor of rank instability within each definition family (Klaeger dataset, active compounds, $n = 206$). Ratio instability is best predicted by the top_1–top_2 affinity gap; entropy, Gini, and S-score instability are best predicted by features of the active affinity distribution.

Overall rank instability — measured as the standard deviation of a compound's rank across all 30 configurations — varies substantially across compounds (Klaeger: mean $\sigma = 35.4$, range 7.1–91.6; Davis: mean $\sigma = 8.5$, range 1.6–19.3). Instability is not uniformly distributed: within each definition family, specific binding profile features predict which compounds receive unstable rankings (Table 2).

Within the ratio family, the gap between the top and second-ranked affinities (top_1–top_2 gap) is the strongest predictor of instability ($r = -0.446$, $p < 0.001$ on Davis; $r = -0.293$, $p < 0.001$ on Klaeger). Compounds with near-identical affinities for their two best-bound kinases receive highly variable ratio rankings because the choice of reference off-target ($k = 1$ vs. $k = 2, \ldots, 5$) determines whether the compound appears selective or promiscuous.

Within the entropy and Gini families, the number of active kinases ($\text{p}K_d > 6$) negatively predicts instability ($r = -0.279$, $p < 0.05$ for entropy on Davis; $r = -0.474$, $p < 0.001$ on Klaeger). Compounds with many active kinases have well-defined broad profiles whose entropy and Gini scores are stable across baseline choices; compounds with few active kinases are sensitive to the baseline because small changes can move targets in or out of the active set, altering the distribution substantially. The active affinity range is also a significant predictor across all three distribution-based definitions on Klaeger ($r = -0.31$ to -0.37, all $p < 0.001$), consistent with the interpretation that heterogeneous profiles are easier to rank stably than

flat ones.

S-score instability is not significantly predicted by any profile feature on either dataset, suggesting that it is driven primarily by where compounds fall relative to the threshold rather than by intrinsic profile shape.

Table 2: Spearman correlations between binding profile features and within-family rank instability on the Klaeger dataset ($n = 206$ active drugs). Significance: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Feature	S-score	Entropy	Gini	Ratio
n_{active}	-0.263^{***}	-0.474^{***}	-0.332^{***}	-0.091
top_1–top_2 gap	$+0.044$	$+0.108$	$+0.068$	-0.293^{***}
Active affinity std	-0.302^{***}	-0.263^{***}	-0.319^{***}	-0.131
Active affinity range	-0.366^{***}	-0.350^{***}	-0.309^{***}	-0.141^{*}
top_2 pK_d	-0.312^{***}	-0.309^{***}	-0.390^{***}	$+0.112$

Three Mechanistically Distinct Sources of Instability



Figure 2: Representative binding profiles illustrating the three instability types. Left: Drug 64 (Davis), most stable ($\sigma = 1.6$), two active kinases. Center: Drug 42 (Davis), most unstable ($\sigma = 19.3$), 22 active kinases at near-identical affinities. Right: Drug 23 (Davis), most ratio-sensitive, five active kinases including two near-tied top targets ($\Delta \text{p}K_d = 0.019$). Dashed line: p$K_d = 6$ activity threshold; dotted line: p$K_d = 5$ assay floor.

Examination of the most and least unstable compounds in the Klaeger dataset reveals three structurally distinct sources of rank instability, each implicating a different aspect of how the definitions are constructed.

Type 1 — Zero-active compounds. The five most unstable Klaeger compounds (e.g. AZD-8055, BMS-911543, Filgotinib; rank $\sigma > 80$) all have zero active kinases at the p$K_d > 6$

threshold. These compounds have no meaningful binding to any profiled kinase at sub-micromolar affinity, yet all four definitions still assign them numerical selectivity scores based on the detection floor values. The resulting rankings are arbitrary: different parameterizations produce different orderings of compounds that are, in practice, indistinguishable from one another. Mean rank instability for zero-active compounds is $\sigma \approx 74$, compared with $\sigma \approx 32$ for compounds with at least one active kinase ($p < 0.001$, Mann–Whitney U test). This pattern is observed only in the Klaeger dataset; all Davis compounds have at least one active kinase, consistent with Davis's fixed-concentration screening design, which reports a K_d only when activity is first detected at 10 μM.

Type 2 — Near-tied top targets. Among active compounds, ratio-specific instability is strongly predicted by a small top_1–top_2 gap ($r = -0.293$, $p < 0.001$). Compounds such as Dovitinib (top_1–top_2 gap = 1.935 pKd units) and BMS-690514 (gap = 1.985) show large entropy-ratio disagreement despite having well-defined primary targets, because the ratio definition is sensitive to whether the reference off-target is the second or a lower-ranked kinase. By contrast, Drug 23 in the Davis dataset (top_1–top_2 gap = 0.019) shows maximal ratio instability because its two best-bound kinases are essentially indistinguishable, making the ratio highly sensitive to which is designated the primary target.

Type 3 — Broad profiles with a dominant primary target. The largest entropy-ratio disagreements in the Klaeger dataset arise for compounds with high n_{active} and a large top_1–top_2 gap (e.g. Dovitinib: $n_{\text{active}} = 26$, gap = 1.935; BMS-690514: $n_{\text{active}} = 20$, gap = 1.985). These compounds have one clearly dominant target but also bind many kinases at moderate affinity. The ratio definition scores them as highly selective (large gap between primary and secondary target), while the entropy and Gini definitions score them as moderately promiscuous (activity spread across many targets). Neither characterization is incorrect; they reflect genuinely different questions about selectivity.

Conversely, the most stable compounds in both datasets (e.g. XL-228, $\sigma = 7.1$; PF-03814735, $\sigma = 8.1$) are highly promiscuous, with 40–57 active kinases. When a compound

binds nearly the entire profiled kinome, all definitions rank it as non-selective regardless of parameterization, producing low instability despite — or rather because of — high promiscuity.

Clinical Outcome Correlations

Spearman correlations between median selectivity rankings and FAERS-derived adverse event rates (serious event rate, death rate, hospitalization rate) across 46 FDA-approved Klaeger compounds were uniformly weak and non-significant (all $|r| < 0.18$, all $p > 0.14$). Similarly, correlations between selectivity rankings and FDA label discontinuation rates due to adverse events ($n = 33$ drugs) were non-significant across all four definition families (all $|r| < 0.27$, all $p > 0.13$).

We attribute these null results to confounding inherent in both data sources rather than absence of a biological relationship between selectivity and toxicity. FAERS report counts are driven primarily by prescription volume and indication prevalence rather than drug toxicity per se; drugs approved for common conditions (e.g. tofacitinib for rheumatoid arthritis, palbociclib for breast cancer) accumulate large report counts regardless of their adverse event profiles relative to alternative therapies. Discontinuation rates from drug labels reflect the clinical trial populations and indication-specific toxicity expectations for those populations, making cross-indication comparisons unreliable. A rigorous test of the relationship between in vitro selectivity and clinical toxicity would require indication-stratified, treatment-related adverse event rates from randomized controlled trials with matched comparators, which are not available in a standardized form across the breadth of kinase inhibitors studied here.

Panel Size Dependence of Selectivity Rankings

To assess the stability of selectivity rankings as a function of kinase panel size, we performed a subsampling analysis on the Klaeger dataset. The Davis dataset was not used for this analysis for two reasons: its complete matrix design (every compound tested against every kinase by construction) does not reflect the sparse coverage of real profiling panels, and

subsampling kinases from Davis changes the apparent number of active compounds in ways that confound the panel size effect with the zero-active instability characterized in Section . The Klaeger dataset, derived from chemoproteomic dose–response profiling with genuine sparse kinase coverage, is the more appropriate substrate for this question. For each panel size $p \in \{50, 80, \ldots, 320, 343\}$, we drew 50 random subsets of p kinases from the full panel of 343, computed each selectivity definition on the subpanel, and measured agreement with the full-panel ranking using Spearman correlation.

The four definitions differ substantially in their panel size requirements (Figure 3). Selectivity entropy reaches mean $r = 0.90$ against the full-panel ranking at approximately 110 kinases, and S-score at approximately 140 kinases. The Gini coefficient requires approximately 290 kinases to reach the same threshold. The ratio definition does not reach $r = 0.90$ until the panel contains 320 or more kinases, and achieves $r = 0.888$ even at 320 kinases — only 23 below the full panel size.

This result has a direct practical implication. Standard commercial kinase selectivity panels typically profile 50–100 kinases.[8] At these panel sizes, entropy- and S-score-based rankings are reasonably stable ($r \approx 0.76$–0.85), but Gini and ratio rankings are effectively uncorrelated with what would be obtained on a full kinome panel ($r < 0.14$ for Gini, $r < 0.00$ for ratio at $p = 50$). Ratio rankings in particular are unreliable at any panel size commonly used in practice, because the identity of the reference off-target (k-th ranked kinase) changes substantially as panel composition changes. This constitutes a fifth source of evidence that ratio-based definitions measure a qualitatively different property from distribution-based definitions, and reinforces D2 (bounded gap sensitivity) as a necessary desideratum.



Figure 3: Mean Spearman correlation between rankings computed on random kinase subpanels and rankings on the full Klaeger panel (343 kinases), as a function of subpanel size. Shaded bands indicate ± 1 standard deviation across 50 subsamples. The dashed line marks $r = 0.90$. Entropy and S-score rankings stabilize at approximately 110–140 kinases; Gini requires approximately 290 kinases; ratio does not reach $r = 0.90$ at any practically used panel size.

Desiderata for a Well-Formed Selectivity Measure

The empirical findings of Section motivate four properties that any well-formed kinase inhibitor selectivity measure should satisfy. We present these as desiderata rather than axioms: they are necessary conditions motivated by observed failure modes of existing definitions, but they are not yet proven to be jointly sufficient to characterize a unique measure, nor is it known whether they are jointly satisfiable. Establishing these properties formally is the subject of future work.

Let $\mathbf{x} = (x_1, \ldots, x_N) \in \mathbb{R}^N$ denote a compound's binding profile across N kinases, where $x_i = \mathrm{p}K_{d,i}$. A selectivity measure is a function $f : \mathbb{R}^N \to \mathbb{R}$, where higher values indicate greater selectivity. A selectivity ranking over a set of compounds is induced by ordering their f-scores.

D1 — Reliability threshold. There exists a threshold $\tau^* > 0$ such that $f(\mathbf{x})$ is undefined, or explicitly flagged as unreliable, when $\max_i x_i \leq \tau^*$.

Motivation. Compounds with no detected binding above the assay detection floor receive arbitrary selectivity scores under all four definitions studied here, producing rank instability with mean $\sigma \approx 74$ compared with $\sigma \approx 32$ for active compounds (Section , Type 1). Selectivity is a property of binding; a compound that does not bind cannot be meaningfully ranked relative to compounds that do. No existing definition incorporates an explicit reliability condition: both the S-score and selectivity entropy assign scores to compounds with no sub-micromolar binding, treating detection floor values as informative.

D2 — Bounded gap sensitivity. For any compound profile $\mathbf{x}$ satisfying D1, a small perturbation to the identity of the reference off-target should produce a bounded change in $f(\mathbf{x})$. Formally, if f depends on the k-th ranked affinity $x_{(k)}$, then $|f(\mathbf{x}; k) - f(\mathbf{x}; k+1)|$ should be bounded by a function of $x_{(k)} - x_{(k+1)}$.

Motivation. Ratio-based definitions violate this property when $x_{(1)} \approx x_{(2)}$: a compound

with near-identical affinities for its top two kinases receives a large ratio score when $k = 1$ (the gap to the next off-target may be large) but a near-zero score when $k = 2$ (the gap between the tied targets is negligible). This produces the Type 2 instability identified in Section , where top_1–top_2 gap is the strongest predictor of ratio instability ($r = -0.293$, $p < 0.001$). Distribution-based definitions (entropy, Gini) satisfy a weaker version of this property because they do not privilege any single pairwise comparison.

D3 — Distributional consistency. The ranking induced by f should be stable under small perturbations to the activity baseline β used to define the active binding distribution. Formally, for any two compounds $\mathbf{x}, \mathbf{y}$ and any $\epsilon > 0$, there exists $\delta > 0$ such that if $|\beta_1 - \beta_2| < \delta$ then the ranking of $\mathbf{x}$ relative to $\mathbf{y}$ under $f(\cdot; \beta_1)$ agrees with that under $f(\cdot; \beta_2)$.

Motivation. The entropy and Gini definitions are parameterized by a baseline β that determines which affinities contribute to the binding distribution. Compounds with few active kinases are sensitive to this choice because small changes in β can move individual kinases in or out of the active set, substantially altering the distribution. The negative correlation between n_{active} and entropy/Gini instability ($r = -0.47$ on Klaeger, $p < 0.001$) reflects this sensitivity. D3 requires that the ranking be robust to the choice of β within a reasonable neighborhood, which existing implementations do not guarantee.

D4 — Monotonicity under weak off-target addition. If compound $\mathbf{x}$ has a weak additional binding interaction added — formally, if $\mathbf{x}' = (x_1, \ldots, x_N, x_{N+1})$ where $x_{N+1} \leq \tau^*$ — then $f(\mathbf{x}') \leq f(\mathbf{x})$. Adding a sub-threshold binding interaction should not increase a compound's selectivity score.

Motivation. Ratio-based definitions can violate this property: if the new weak interaction becomes the reference off-target ($k = 1$), the ratio to the primary target may increase, artifactually improving the apparent selectivity score. This failure mode is practically relevant when profiling panels differ in size or composition across experiments, as is common when comparing selectivity across different assay platforms or across datasets such as Davis and

Klaeger. Distribution-based definitions satisfy D4 naturally, since adding a sub-threshold term to the probability distribution decreases or preserves entropy and Gini.

Table 3 summarizes which existing definitions satisfy each desideratum. No existing definition satisfies all four simultaneously. Whether a definition satisfying all four exists, and whether the four desiderata are jointly consistent, are open questions that we identify as the primary targets of a formal axiomatization.

Table 3: Satisfaction of desiderata by existing selectivity definitions. ✓: satisfied by construction. ∼: partially satisfied or satisfied under additional assumptions. ×: violated in general.

Desideratum	S-score	Entropy	Gini	Ratio
D1 (reliability threshold)	×	×	×	×
D2 (bounded gap sensitivity)	∼	✓	✓	×
D3 (distributional consistency)	∼	×	×	✓
D4 (monotonicity, weak off-target)	✓	✓	✓	×

Discussion

The central finding of this work is that existing kinase inhibitor selectivity definitions do not measure the same quantity. The ratio definition and the distribution-based family (S-score, entropy, Gini) produce substantially different compound rankings ($r = 0.27$–0.48), and the sources of disagreement are mechanistically interpretable rather than random. This has practical consequences: a compound that ratio-based analysis designates as selective may be designated as moderately promiscuous by entropy-based analysis, and vice versa, depending on the shape of its binding profile. Go/no-go decisions in early drug discovery are routinely made on the basis of selectivity assessments without acknowledgment that the choice of definition materially affects the outcome.

The Two-Family Structure Reflects a Conceptual Distinction

The clustering of definitions into two families is not a statistical artifact but reflects a genuine difference in what is being measured. Distribution-based definitions ask: how concentrated is the compound's binding activity across the full profiled kinome? The ratio definition asks: how much more potently does the compound bind its primary target than a specified off-target? These are distinct questions with distinct answers for a large class of compounds — specifically, those with one clearly dominant target that also bind many kinases at moderate affinity (Type 3 profiles in our taxonomy). For such compounds, ratio-based methods report high selectivity while distribution-based methods report moderate promiscuity, and both characterizations are correct given their respective questions.

The disagreement is therefore not a defect to be corrected by choosing the "right" definition, but a signal that the field has not agreed on what selectivity means. The desiderata proposed in Section are an attempt to make this question precise enough that it can be answered formally.

Practical Recommendations

Pending a formal resolution, three practical recommendations follow from our findings.

First, D1 (reliability threshold) should be adopted immediately as a reporting standard. Compounds with no detected binding above the assay detection floor should not receive selectivity scores, or should have their scores explicitly flagged as unreliable. Of the 222 Klaeger compounds, 16 (7.2%) fall into this category and currently receive scores that are indistinguishable from noise.

Second, selectivity assessments should report the sensitivity of rankings to parameter choice rather than a single score at a single parameter value. Our analysis shows that rank standard deviation across the parameter sweep is a compound-specific, interpretable quantity that identifies which compounds have robust selectivity assessments and which do not. This is analogous to reporting confidence intervals in statistical estimation.

Third, when comparing selectivity across studies that use different definitions, the comparison should be restricted to the subset of compounds and parameter values for which the definitions agree. The high within-family correlations ($r > 0.74$) suggest that S-score, entropy, and Gini are interchangeable for most purposes when parameters are chosen consistently; ratio-based measures should not be compared directly with distribution-based measures without acknowledging the conceptual distinction.

Limitations

The analysis is limited to two profiling datasets that, while large and well-characterized, represent a small fraction of the chemical and kinase space relevant to drug discovery. Davis covers 68 compounds and 433 kinases using a fixed-concentration competition binding assay; Klaeger covers 222 clinical compounds and 343 kinases using chemoproteomic dose–response profiling. The two assay technologies are not directly comparable, and the difference in active compound fractions (17.9% for Davis vs. 3.6% for Klaeger) reflects both genuine selectivity differences and assay-specific detection thresholds. Findings should be interpreted in light of

these technological differences.

The clinical outcome analysis is limited by the observational nature of FAERS data and the heterogeneity of FDA label discontinuation rates across indications. The null results reported in Section are consistent with a genuine absence of correlation between in vitro selectivity and clinical toxicity, but they are equally consistent with the confounders identified therein. A properly controlled study would require prospective collection of treatment-related adverse event rates stratified by indication, which is not currently available in a standardized form across the breadth of approved kinase inhibitors.

The desiderata proposed in Section are necessary conditions motivated by empirical failure modes; they are not proven to be sufficient to uniquely characterize a selectivity measure, and it is not known whether they are jointly satisfiable. The satisfaction assessments in Table 3 are based on analysis of the functional forms of each definition and the empirical findings of this study; a formal proof of satisfaction or violation for each desideratum–definition pair is left for future work.

Relation to Formal Methods

The approach taken here — characterizing the empirical behavior of competing definitions to motivate formal desiderata — follows a methodology well established in adjacent fields. Shannon's derivation of entropy began with the observation that several reasonable-seeming measures of uncertainty existed and proceeded to identify the axioms that single out one uniquely.[10] Arrow's impossibility theorem began with the observation that several reasonable voting rules existed and proved that no rule satisfies all desirable properties simultaneously.[11]

The present work occupies the empirical stage of this process: we have characterized where and why existing definitions disagree, and proposed properties that any satisfactory definition should have. Whether a definition satisfying all four desiderata exists, whether the desiderata are jointly consistent, and whether an impossibility result holds, are questions for a formal axiomatization. We anticipate that such an axiomatization, potentially implemented in a

proof assistant such as Agda or Lean, would constitute a natural and productive extension of this work.[18]

Conclusion

We have shown that commonly used kinase inhibitor selectivity definitions cluster into two families that measure fundamentally different properties of binding profiles, that rank instability within and between families is mechanistically predictable from binding profile shape, and that three structurally distinct sources of instability can be identified and characterized. These findings motivate four desiderata — a reliability threshold condition, bounded gap sensitivity, distributional consistency, and monotonicity under weak off-target addition — that any well-formed selectivity measure should satisfy. No existing definition satisfies all four simultaneously.

The immediate practical implication is that selectivity assessments should report the sensitivity of rankings to parameter choice, and should explicitly flag compounds with no sub-threshold binding as unreliable. The longer-term implication is that the field requires a principled basis for choosing among selectivity definitions, analogous to the axiomatic foundations that have been established for entropy in information theory and for ranking functions in social choice theory. The desiderata proposed here are intended as the empirical foundation for that formal program.

Acknowledgement

The authors thank the developers of the Davis and Klaeger datasets for making their data publicly available.

Supporting Information Available

Code and data are available at `https://github.com/polinavino/kinase-selectivity-definitions`.

References

(1) Roskoski, R. Properties of FDA-approved small molecule protein kinase inhibitors: a 2024 update. *Pharmacol. Res.* **2024**, *200*, 107059.

(2) Manning, G.; Whyte, D. B.; Martinez, R.; Hunter, T.; Sudarsanam, S. The protein kinase complement of the human genome. *Science* **2002**, *298*, 1912–1934.

(3) Bhullar, K. S.; others Kinase-targeted cancer therapies: progress, challenges and future directions. *Mol. Cancer* **2018**, *17*, 48.

(4) Karaman, M. W.; others A quantitative analysis of kinase inhibitor selectivity. *Nat. Biotechnol.* **2008**, *26*, 127–132.

(5) Uitdehaag, J. C. M.; Zaman, G. J. R. A theoretical entropy score as a single value to express inhibitor selectivity. *BMC Bioinformatics* **2011**, *12*, 94.

(6) Graczyk, P. P. Gini coefficient: a new way to express selectivity of kinase inhibitors against a family of kinases. *J. Med. Chem.* **2007**, *50*, 5773–5779.

(7) Klaeger, S.; others The target landscape of clinical kinase drugs. *Science* **2017**, *358*, eaan4368.

(8) Bosc, N.; Meyer, C.; Bonnet, P. The use of novel selectivity metrics in kinase research. *BMC Bioinformatics* **2017**, *18*, 17.

(9) Dranchak, P.; others KInhibition: a kinase inhibitor selection portal. *iScience* **2018**, *8*, 29–37.

(10) Shannon, C. E. A mathematical theory of communication. *Bell Syst. Tech. J.* **1948**, *27*, 379–423.

(11) Arrow, K. J. *Social Choice and Individual Values*; Wiley: New York, 1951.

(12) Miljkovic, F.; Bajorath, J. Reconciling selectivity trends from a comprehensive kinase inhibitor profiling campaign with known activity data. *ACS Omega* **2018**, *3*, 17196–17206.

(13) Khinchin, A. Y. *Mathematical Foundations of Information Theory*; Dover: New York, 1957.

(14) Pavillidis, P.; others Towards an axiomatic approach to truth discovery. *Auton. Agents Multi-Agent Syst.* **2022**, *36*, 43.

(15) Cardelli, L. Brane calculi. *Lect. Notes Comput. Sci.* **2004**, *3082*, 257–278.

(16) Jones, B.; others The Beacon Calculus: a formal method for the coarse-grained modelling of biological systems. *PLOS Comput. Biol.* **2020**, *16*, e1007646.

(17) Kennedy, A. J. Dimension types. Programming Languages and Systems – ESOP 1994. 1994; pp 348–362.

(18) The Mathlib Community Mathlib4. 2024; `https://github.com/leanprover-community/mathlib4`.

(19) Davis, M. I.; others Comprehensive analysis of kinase inhibitor selectivity. *Nat. Biotechnol.* **2011**, *29*, 1046–1051.

(20) Öztürk, H.; Özgür, A.; Ozkirimli, E. DeepDTA: deep drug–target binding affinity prediction. *Bioinformatics* **2018**, *34*, i821–i829.

(21) Harris, C. R.; others Array programming with NumPy. *Nature* **2020**, *585*, 357–362.

(22) McKinney, W. Data structures for statistical computing in Python. Proceedings of the 9th Python in Science Conference. 2010; pp 56–61.

(23) Virtanen, P.; others SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nat. Methods* **2020**, *17*, 261–272.