

The Complement Rule Is Excluded Middle: Machine-Checked Probability over Intuitionistic Logic

Anonymous Author(s)

Abstract

Kolmogorov’s axioms fix not only an arithmetic of chance but a *logic* of events, a Boolean algebra, and quantum mechanics long ago showed that the logic is a modeling choice. We mechanize, in Lean 4 over mathlib, what probability becomes when the event logic is *intuitionistic*: valuations on frames, the algebras of verifiable propositions. The complement rule $v(\neg A) = 1 - v(A)$ holds for every valuation exactly when excluded middle holds in the algebra, and Dempster conditioning agrees with Bayesian conditioning exactly when the evidence satisfies excluded middle: probability’s classicality, static and dynamic, is one logical axiom. On finite frames every valuation is a classical mass function on points, in general it splits into atomic and diffuse parts, and its Boolean core carries a totally monotone Dempster–Shafer belief function. The “slack” replacing the complement rule is boundary mass in the measure model, and removing it from a semidecidable event solves the halting problem: slack is the price of a computable epistemic state. A machine-checked dichotomy delimits the probability-monad question, the naive Cox statement is unsatisfiable, and the Aczél/Hölder analytic core is built from scratch, in about 4,300 sorry-free lines.

Keywords: intuitionistic logic, probability, Cox’s theorem, belief functions, locales, Lean, mathlib, formalized mathematics

1 Introduction

Classical logic underlies Kolmogorov’s probability. The axioms of this probability specify an arithmetic of chance (non-negativity, normalization, additivity). Beneath that arithmetic sits a logic: the events form a Boolean σ -algebra, closed under complement and satisfying excluded middle. The complement rule, $P(\neg A) = 1 - P(A)$, is where the two layers touch. It is an arithmetical equation about a logical operation, obeyed by every assignment of degrees of certainty. In the limiting case of all-or-nothing probabilities, it becomes equivalent to the law of excluded middle.

There is no intrinsic reason why probability should be compatible with exactly one logic as different situations are described by different logics. Intuitionistic logic describes computation, verification, and finite-precision observation, where a proposition may be confirmable but not refutable. Linear logic describes resources, where using an assumption consumes it [Girard 1987]. Quantum logic describes measurement, where propositions fail to distribute. If probability is, as the Cox–Jaynes tradition has it, *extended logic* [Cox

1946; Jaynes 2003], the calculus of degrees of certainty over a space of propositions, then the logic of those propositions is a *parameter* of the theory, and Kolmogorov’s calculus is the value of that parameter at Boolean logic.

This is not a speculative proposal, because physics has already forced the move once. Quantum probability lives on orthomodular lattices of projections rather than Boolean algebras [Birkhoff and von Neumann 1936], and Gleason’s theorem [Gleason 1957] plays the role there that this paper’s representation theorems play here: it identifies the states compatible with the event logic. Nobody concludes that quantum probability is “not really probability.” The conclusion is that the Boolean substrate was a modeling choice. Our question is what happens at a different value of the same parameter, motivated by the logic used to reason about computation: what does probability become when the logic of events is *intuitionistic*?

Intuitionistic logic is the natural first non-Boolean instance to study using the proofs-as-programs paradigm. The choice of (in our case, intuitionistic) logic sits upstream of statistics the way a type system sits upstream of programs: the same syntax of operations, a different set of valid transformations, and some derivations that typecheck classically can either fail or split in two. This logic’s algebraic semantics is mature order theory, well supported by mathlib [mathlib Community 2020]: frames, that is, complete Heyting algebras, modelled on the lattices of open sets. The content of the original (classical) probability definitions and results is not lost: complete Boolean algebras are exactly the frames where excluded middle holds, so the classical theory embeds as a special case rather than being replaced.

The choice of logic fixes the event algebra. What remains is to fix the arithmetic on top of it, keeping only the axioms that can be stated without a complement. A *valuation* on a frame Ω is a monotone, normalized, modular map $v : \Omega \rightarrow [0, \infty]$. Modularity is inclusion–exclusion, $v(a) + v(b) = v(a \sqcup b) + v(a \sqcap b)$, and no axiom mentions complement. In a Heyting algebra the pseudo-complement satisfies $a \sqcap \neg a = \perp$ but not, in general, $a \sqcup \neg a = \top$. The complement rule therefore degrades to a derived *inequality* $v(a) + v(\neg a) \leq 1$, with a gap

$$\text{slack}(a) = 1 - v(a) - v(\neg a)$$

measuring exactly the valuation of the region where excluded middle fails for a . The term *slack*, and this logical reading, are introduced in this paper. The quantity itself is not new: it is the width of the classical Dempster–Shafer belief/plausibility interval [Dempster 1967; Shafer 1976]. The

two align on the frame's Boolean core. Dempster–Shafer reads that width as an agent's ignorance. Here it needs no agent: in the concrete model where Ω is the opens of a space carrying a probability measure μ , the slack of an open U is precisely $\mu(\partial U)$, the measure of its topological boundary. The slack of an open set is a structural quantity, not a measure of ignorance. The theory that results is recognizably the *shape* of Dempster–Shafer belief functions, but derived from the underlying logic rather than postulated.

The statistical *operations* are logic-uniform. The conditional is still $v(a \sqcap b)/v(b)$, and the product rule and Bayes symmetry hold verbatim. What varies is the set of *identities* those operations satisfy, and with it the licensed computations. The classical case split over $\{A, \neg A\}$, and with it the usual denominator of Bayes' theorem $P(B) = P(B \mid A)P(A) + P(B \mid \neg A)P(\neg A)$, degrades to a lower bound whose deficit is a conditional slack. Marginalization is exact only over *genuine* partitions, families that actually join to $\top$. The complement trick $P(A) = 1 - P(\neg A)$ degrades to the interval $[v(A), 1 - v(A^c)]$ of width $\text{slack}(A)$, so point statistics becomes interval statistics exactly on the events the logic leaves undecided, and collapses back to points on complemented ones. Updating itself forks: Dempster's rule and Bayesian conditioning, classically indistinguishable, give different posteriors from the same evidence unless its slack vanishes.

The choice of logic changes the query being answered: the value of an event A is $P(\Box A)$, i.e. the probability of its verifiable part. In the measure model, slack is boundary mass, so a nonatomic distribution queried on intervals behaves entirely classically, since an interval's boundary is two points, which carry no mass. The differences are observable precisely at events with charged boundaries and at semidecidable hypotheses.

A theory this close to excluded middle must be developed with unusual care, because classicality is easy to smuggle in: a single informal case split over $\{A, \neg A\}$ silently decides an undecidable event. The risk is documented even classically, as Cox's original theorem required repair after Halpern's counterexample [Cox 1946; Halpern 1999; Horn 2003], and the constructive setting multiplies the opportunities. The direct transcription of the classical Cox statement within an intuitionistic logic setting is not merely unprovable but *unsatisfiable*. The sum rule is irreducible: a disjunction's value is provably not a function of the values of its disjuncts, so modularity must be posited rather than derived. Section 8 justifies each appearance of classical logic in this work, including at the meta-theory level.

Contributions. Our contributions are as follows:

1. **The two hinges** (Sections 3 and 5). The complement rule holds for *every* valuation iff excluded middle holds in the frame, an equivalence isolating the exact content of Van Horn's classical negation axiom R3 [Horn

2003]. Dempster and Bayesian conditioning agree iff the evidence has zero slack;

2. **Structure theorems** (Sections 4 and 5). On finite frames a valuation is a classical mass function on points. In general it splits into atomic and diffuse parts, and on its Boolean core of regular elements it is a belief function;
3. **The Cox program, dissected** (Section 6, Appendix A). The product-rule half is independent of the logic, proved by an Aczél/Hölder generator built from scratch. The sum-rule half (a theorem in classical Cox) admits no such derivation here, as shown by a five-element counterexample, and must therefore be included among the defining axioms of a valuation;
4. **Two grounding models** (Section 7). Any classical probability measure, read on the opens, is a valuation whose slack is boundary mass. On the Sierpiński frame, a slack-free classical readout of halting is uncomputable;
5. **The monad dichotomy** (Section 10). Product valuations are unique on chain products and provably non-unique on the powerset frame, where Fubini fails. The countable fragment composes associatively and commutatively;
6. **Mechanization** (Section 8). Everything above is mechanized in (Lean 4 using mathlib), and is (to our knowledge) the first mechanization of non-additive probability over intuitionistic logic.

2 Valuations on Frames

2.1 The event logic

A *frame* is a complete lattice in which finite meets distribute over arbitrary joins. Equivalently, it is a complete Heyting algebra. Frames are the algebraic semantics of intuitionistic logic and the point-free abstraction of topological spaces. The same objects with morphisms reversed form the category of *locales*, so the distinction is invisible at the level of a single algebra. We say *frame* throughout and reserve *locale* for the spatial and categorical statements (points of a locale, classifying locales) where that word is the standard one. The motivating example throughout is $\Omega = \mathcal{O}(X)$, the opens of a space, with $\sqcap, \sqcup$ as intersection and union and the pseudo-complement a^c of an open given by the *interior* of its set complement. In mathlib this is the typeclass `Order.Frame`, and the Heyting operations come with the key asymmetry already in place: $a \sqcap a^c = \perp$ holds in any Heyting algebra (`inf_compl_eq_bot`), while $a \sqcup a^c = \top$ is available only in Boolean algebras. An element with $a \sqcup a^c = \top$ is *complemented*: it satisfies its own instance of excluded middle.

2.2 The valuation structure

The central object assigns a degree of certainty to each element of the frame, with inclusion–exclusion as the only interaction axiom:

Definition 1 (Valuation). A *valuation* on a frame Ω is a function $v : \Omega \rightarrow [0, \infty]$ with $v(\perp) = 0$, $v(\top) = 1$, monotonicity, and *modularity*:

$$v(a) + v(b) = v(a \sqcup b) + v(a \sqcap b).$$

```
structure Valuation (Ω : Type*) [Order.
  Frame Ω] where
  toFun : Ω → ℝ≥0∞
  map_bot' : toFun ⊥ = 0
  map_top' : toFun ⊤ = 1
  mono' : Monotone toFun
  modular' : ∀ a b, toFun a + toFun b
    = toFun (a ⊔ b) + toFun (a ⊓ b)
```

Values live in $[0, \infty]$ (ENNReal, extended non-negative reals) following mathlib’s measure-theoretic convention, and normalization then confines them to $[0, 1]$ (Valuation.le_one). The definition is Kolmogorov’s minus one thing: *no axiom mentions complement*. Classically that omission is invisible, because the complement rule is derivable from additivity and $a \vee \neg a = \top$. Constructively it is the whole story.

2.3 Slack, and its two independent obstructions

Modularity applied to the disjoint pair (a, a^c) collapses to $v(a) + v(a^c) = v(a \sqcup a^c)$ (add_compl_eq_sup): the sum measures a ’s instance of excluded middle. Since that join need not be $\top$:

Theorem 2 (add_compl_le_one). *For every valuation and every a : $v(a) + v(a^c) \leq 1$.*

This is the Dempster–Shafer sub-additivity inequality, here *derived* from the logic rather than postulated. The deficit

$$\text{slack}(a) = 1 - (v(a) + v(a^c)) = 1 - v(a \sqcup a^c)$$

is the valuation of the undecided region. It decomposes canonically. The pair (a^{cc}, a^c) , double negation against negation, is always disjoint but is a partition only under *weak* excluded middle. Running it through modularity splits the slack into two logically independent gaps:

Theorem 3 (slack_eq_dnGap_add_deMorganGap). *$\text{slack}(a) = \text{dnGap}(a) + \text{deMorganGap}(a)$, where $\text{dnGap}(a) = v(a^{cc}) - v(a)$ vanishes when a is $\neg\neg$ -stable (regular), and $\text{deMorganGap}(a) = 1 - v(a^{cc} \sqcup a^c)$ vanishes when weak excluded middle holds at a .*

The single Dempster–Shafer “ignorance” number is thus resolved into a regularity defect plus a De Morgan defect, each tracking a named intermediate logic. An informal aside on

what the first gap measures: intuitionistically the contrapositive of a universal hypothesis returns only its double-negated form, so in Hempel’s raven paradox [Hempel 1945] the white shoe confirms at most a^{cc} , and dnGap is exactly the distance that indoor evidence cannot cross.

2.4 What survives without excluded middle

Three classical pillars survive intact, and one fails in a precisely delimited way. *Disjoint additivity* is unconditional: $a \sqcap b = \perp$ implies $v(a \sqcup b) = v(a) + v(b)$. *Conditioning* is native: for $v(b) \neq 0$ the posterior $v(a \mid b) = v(a \sqcap b) / v(b)$ is again a valuation (the proof is frame distributivity), and the product rule $v(a \mid b) \cdot v(b) = v(a \sqcap b)$ and Bayes symmetry $v(a \mid b) v(b) = v(b \mid a) v(a)$ hold (condVal_mul, condVal_symm). *Total probability* holds over any genuine partition, meaning $a \sqcap a' = \perp$ and $a \sqcup a' = \top$: $v(b) = v(a \sqcap b) + v(a' \sqcap b)$. What fails is total probability over $\{a, \neg a\}$. That family does not tile $\top$, and the conditional masses fall short of $v(b)$ by a conditional slack (cond_add_compl_le). Bayesian updating survives. The assumption that A and $\neg A$ exhaust the world does not.

The classical fragment. We characterize the elements of Ω on which the classical complement rule holds. The $\neg\neg$ -stable (*regular*) elements, which form a Boolean algebra by Glivenko’s theorem, are a natural candidate, but they do not suffice: modularity constrains the frame join $\sqcup$, whereas the Boolean join of regular elements is the regularized $(a \sqcup b)^{cc}$, for which modularity survives only as the inequality $v(a) + v(b) \leq v((a \sqcup b)^{cc}) + v(a \sqcap b)$, the 2-monotonicity of Section 5. A regular but uncomplemented element (the open ray $(-\infty, 0)$ in the opens of $\mathbb{R}$, under an atomic measure at 0) still carries slack. The correct classical fragment is the *complemented* elements, where $v(a) + v(a^c) = 1$ holds on the nose (add_compl_eq_one_of_complemented). On a connected locale that fragment can be as small as $\{\perp, \top\}$. In the Boolean limit every element is complemented, the slack vanishes identically, and Kolmogorov is recovered (classical_additivity, classical_slack_zero).

Mixtures. Valuations are closed under finite convex combination (Valuation.mix), with every axiom inherited termwise, because modularity is linear. This innocuous fact becomes a characterization in Section 4.

3 Where Classicality Lives: the R3 Hinge

Van Horn’s repaired Cox theorem [Horn 2003] axiomatizes plausibility over classical propositions. Its negation axiom, called R3, posits that the plausibility of $\neg A$ is a fixed non-increasing function S of the plausibility of A . Consistency forces $S \circ S = \text{id}$, an involution, and thence the complement rule $p \mapsto 1 - p$. An involutive negation is double-negation elimination, so R3 is where classical logic enters every known Cox-style derivation. We isolate the exact content of R3’s

conclusion as a predicate on valuations and prove it equivalent to excluded middle.

```
def HasClassicalNegation (v : Valuation Ω)
  : Prop :=
  ∀ a, v a + v ac = 1
```

Theorem 4 (the R3 hinge). *For a frame Ω , every valuation on Ω has classical negation iff $a \sqcup a^c = \top$ for every $a \in \Omega$.*

The forward direction (`hasClassicalNegation_of_em`) is a calculation. The content is the converse (`em_of_forall_hasClassicalNegation`), where excluded middle fails, a valuation with genuine slack must be *manufactured*. Suppose $a \sqcup a^c \neq \top$. The prime ideal separation theorem (`mathlib's DistribLattice.prime_ideal_of_disjoint_filter_ideal`) yields a prime ideal J containing $a \sqcup a^c$ but not $\top$. The indicator of J 's complement,

$$v(x) = \begin{cases} 0 & x \in J \\ 1 & x \notin J, \end{cases}$$

is a valuation. Modularity in the only nontrivial case ($x, y \notin J$) is *exactly* primeness, $x \sqcap y \in J \Rightarrow x \in J \vee y \in J$. And this valuation assigns $v(a) + v(a^c) = 0 \neq 1$. So the complement rule is not one convention among many. Quantified over all valuations, it is a statement *about the algebra*, and the statement is excluded middle.

Sharp valuations are the points of the locale. The construction above is not ad hoc. It is one half of an exact correspondence. Call v *sharp* if it is $\{0, 1\}$ -valued, a state of full certainty. Sharpness characterizes exactly the prime-ideal indicators:

Theorem 5 (`sharp_iff_point`). *v is sharp iff v is the complement-indicator of a prime ideal. The maps are mutually inverse: the ideal recovered from v is its zero-set, prime exactly because v is sharp.*

In locale-theoretic terms, the certainty limit of the probability theory recovers the *points* of the locale. (These are the finitely prime points. The completely prime points would correspond to sharp valuations that are also Scott-continuous, a refinement we note but do not formalize.) The hinge and the correspondence together are the paper's static answer to "where does classicality live": in one axiom, whose semantic shadow is point-hood.

4 Representation: Points, Masses, Mixtures

Is intuitionistic probability a new animal, or classical probability seen at an angle? The answer is exact: on finite frames the latter, in general the former, with Scott continuity marking the boundary.

4.1 Finite frames: every valuation is a PMF on points

Every finite frame is the lattice of *lower sets* of a finite poset P , by Birkhoff duality. We work with `LowerSet P` directly

and cite the duality rather than mechanize it. Define the *mass* of a point as the discrete derivative of v :

```
def Valuation.mass (v : Valuation (
  LowerSet P)) (p : P) :
  ℝ≥0∞ := v (LowerSet.Iic p) - v (
  LowerSet.Iio p)
```

The mass function accounts for the valuation completely, and the masses form a probability:

Theorem 6 (finite representation, `eq_sum_mass`, `sum_mass`). *For finite P and any valuation v on $\text{Low}(P)$: $v(U) = \sum_{p \in U} \text{mass}(p)$ and $\sum_p \text{mass}(p) = 1$.*

The proof peels a maximal point p of U at a time. Modularity applied to $(U \setminus \uparrow p, \downarrow p)$ shows the increment is exactly $\text{mass}(p)$, and a strong induction on $|U|$ finishes. The arithmetic is in $[0, \infty]$ with truncated subtraction, where the bookkeeping is real work. Two packagings make the classicality literal. `Valuation.toPMF` exhibits mass as a `mathlib` PMF on P . And since lower sets are the opens of the Alexandrov topology, where interior is the identity, the theorem reads: every valuation on a finite frame is a classical probability seen through the interior operator, $v = \mu \circ \square$. This is the converse, in the finite case, of the measure-theoretic bridge of Section 7.

Dually, with δ_p the sharp *point-valuation* $\delta_p(U) = [p \in U]$:

Theorem 7 (mixture characterization, `eq_mix_deltaPoint`). *On a finite frame, $v = \sum_p \text{mass}(p) \cdot \delta_p$: valuations are exactly the finite convex mixtures of point-valuations.*

Modularity is thereby *characterized* as closure under mixing certainties. This is the constructive answer to what the sum rule axiomatizes, given that it cannot be derived (Section 6).

4.2 The infinite boundary: enter Scott continuity

Both halves of the finite theorem fail at the first infinite frame, and the failure is informative.

Theorem 8 (`obstruction_exists_valuation_not_point_representable`). *On $\text{Low}(\mathbb{N})$, the indicator of $\top$ is a valuation with $\text{mass}(p) = 0$ at every point yet $v(\top) = 1$. The unit of mass escapes to the missing point at infinity.*

Approximating the whole frame from below is exactly what the counterexample refuses to do, and requiring that approximation recovers the representation:

Theorem 9 (`recovery_eq_tsum_mass_of_scott`, generalized by `isPurelyAtomic_of_scott`). *On $\text{Low}(\mathbb{N})$, Scott continuity restores point-representability: if $v(\top) = \bigsqcup_n v(\text{Iic } n)$ then $v(\top) = \sum'_n \text{mass}(n)$. The counterexample of Theorem 8 is itself a proved failure of Scott continuity (`topIndicator_not_scott`). More generally, on any poset all of whose principal downsets are finite, Scott continuity implies zero diffuse part.*

With no hypotheses at all, the point masses still account for at most the whole:

Theorem 10 (general decomposition, `tsum_mass_1e`). *On $\text{Low}(P)$ for arbitrary P , with no finiteness and no continuity assumption: $\sum_p \text{mass}(p) \leq v(\top)$.*

Every valuation thus splits unconditionally into an *atomic* part carried by points, always a sub-probability, and a *diffuse* remainder $v(\top) - \sum_p \text{mass}(p) \geq 0$. This is a constructive Lebesgue-style decomposition. The finite and Scott cases are the purely atomic extreme, and the $\top$ -indicator is the purely diffuse one. The proof of Theorem 10 re-runs the maximal-point peel over arbitrary finite subsets, bounded by the whole, and passes to the supremum. Its proof script makes no excluded-middle case split. Classicality enters only through `mathlib`'s baseline (Section 8). Representing the diffuse part on non-spatial frames, where, tellingly, the obstruction is *too few points*, the same phenomenon as undecidability in Section 7, is the open frontier (Section 10).

5 The Dempster–Shafer Reading, Completed

Sections 2–4 keep meeting the *shape* of Dempster–Shafer (DS) theory [Dempster 1967; Shafer 1976]: sub-additive complements, a belief/plausibility interval, mass functions. This section makes the identification exact in both statics and dynamics, and locates, each time, the precise point where excluded middle is the difference.

5.1 Statics: total monotonicity via constructive inclusion–exclusion

The $\neg\neg$ -stable (*regular*) elements of a frame form a Boolean algebra with meet $\sqcap$ and join $(a \sqcup b)^{\text{cc}}$ (Glivenko). Call it the *Booleanization*. A DS belief function is a capacity on a Boolean algebra that is n -monotone for every n : $\text{Bel}(\bigvee_i A_i) \geq \sum_{\emptyset \neq T \subseteq s} (-1)^{|T|+1} \text{Bel}(\bigwedge_{i \in T} A_i)$. The 2-monotone case is a one-liner from modularity (`two_monotone`). The full tower follows from a fact that is interesting in its own right:

Theorem 11 (inclusion–exclusion, `inclusion_exclusion`). *For every valuation on every frame and every finite family $a : \iota \rightarrow \Omega$, inclusion–exclusion holds with equality for the frame join:*

$$v\left(\bigsqcup_{i \in s} a_i\right) = \sum_{\emptyset \neq T \subseteq s} (-1)^{|T|+1} v\left(\bigsqcap_{i \in T} a_i\right).$$

(Stated in Lean additively, with the odd-cardinality half of the signed sum set against the even-cardinality half, so that no $[0, \infty]$ subtraction occurs. The proof is induction with modularity and frame distributivity.)

Reading the same signed sum against the Booleanization join turns the equality into the full belief-function tower:

Theorem 12 (total monotonicity, `infty_monotone`). *For every valuation, finite family, and n :*

$$\sum_{\emptyset \neq T} (-1)^{|T|+1} v\left(\bigsqcap_T a_i\right) \leq v\left(\left(\bigsqcup_i a_i\right)^{\text{cc}}\right).$$

Read on the Booleanization, where $(\cdot)^{\text{cc}} \circ \sqcup$ is the join and meets of regulars are regular, this is verbatim the ∞ -monotone tower: v restricted to its Booleanization is a bona fide belief function.

The proof of Theorem 12 from Theorem 11 is one line: monotonicity along $a \leq a^{\text{cc}}$. The conceptual point runs opposite to the usual telling. Inclusion–exclusion does *not* fail constructively. It is exact for the intuitionistic disjunction, and total monotonicity is what that equality looks like through the double-negation nucleus: rounding a join up to its regular hull can only add value, and the defect is exactly the double-negation gap of Section 2. Additive probability does *not become* a belief function by losing an axiom. It becomes one by being *read through* $\neg\neg$. With $\text{Pl}(a) = 1 - v(a^c)$, the belief–plausibility interval has width exactly the slack (`plausibility_sub_self`). The DS “ignorance” is boundary measure, not psychology.

5.2 Dynamics: Dempster against Bayes, separated by one number

DS theory carries two update rules that coincide classically. *Geometric* conditioning renormalizes belief, and under the DS dictionary it is the localic posterior $v(a \sqcap b) / v(b)$ of Section 2. *Dempster’s rule of conditioning* [Shafer 1976] discards the mass incompatible with the evidence and renormalizes by its plausibility:

$$v(a \parallel b) = \frac{v(a \sqcup b^c) - v(b^c)}{1 - v(b^c)}.$$

Both are again valuations (`dempsterCond`, whose modularity is the frame distributivity $(a \sqcup b^c) \sqcap (a' \sqcup b^c) = (a \sqcap a') \sqcup b^c$). Their disagreement is governed by a single structural quantity, the *excluded-middle gap* $\text{emGap}(a, b) = v(a) - v(a \sqcap (b \sqcup b^c))$, the mass of a stranded outside b ’s instance of excluded middle:

Lemma 13 (`sup_compl_decomp`). $v(a \sqcup b^c) = v(b^c) + v(a \sqcap b) + \text{emGap}(a, b)$. *Dempster’s numerator exceeds Bayes’s by exactly the gap.*

The gap, and with it the disagreement of the two update rules, is controlled by one number, the slack at the evidence:

Theorem 14 (the conditioning hinge, `dempsterCond_eq_condVal14iff_s`). *For $v(b) \neq 0$: Dempster conditioning on b agrees with Bayesian/geometric conditioning on b at every proposition iff $\text{slack}(b) = 0$.*

Zero slack at the evidence kills the gap for every a and makes the two normalizers coincide, since $1 - v(b^c) = v(b)$. Conversely, testing at $a := b$, geometric conditioning gives

certainty 1 while Dempster gives $v(b)/(1 - v(b^c))$, which is 1 only if the slack vanishes. Theorem 14 is the dynamic companion of Theorem 4. The static hinge locates classicality in the complement rule, and the dynamic hinge locates it in the agreement of the two DS update rules: one instance of excluded middle, at the evidence. In the measure model, Dempster and Bayes disagree exactly when the evidence has a charged boundary.

6 The Cox Program, Dissected

Cox's theorem [Cox 1946; Jaynes 2003] derives the probability calculus from desiderata on plausibility. After Halpern's counterexample [Halpern 1999], its repaired forms [Arnborg and Sjödin 2000; Horn 2003] fix the calculus up to monotone regratuation. Every known derivation is classical, and Section 3 showed why: the negation axiom R3 is excluded middle. What survives of the program when R3 is dropped? Mechanization forced a sharper answer than we had conjectured, in three parts. The product rule is logic-free. The sum rule is irreducible. And the naive constructive statement is unsatisfiable.

6.1 A Cox structure without negation

```

structure CoxModel (Ω : Type*) [Order.
  Frame Ω] where
  pl : Ω → Ω → ℝ          -- pl a b :
    plausibility of a given b
  mono_left : ∀ c, Monotone fun a => pl a
    c
  pl_bot : ∀ c, pl ⊥ c = 0
  pl_top : ∀ c, pl ⊤ c = 1
  F : ℝ → ℝ → ℝ          -- conjunction
    functional
  conj : ∀ a b c, pl (a ⊔ b) c = F (pl a (
    b ⊔ c)) (pl b c)
  F_assoc : ∀ x y z, F (F x y) z = F x (F
    y z)
  F_strictMono_left : ∀ y, 0 < y →
    StrictMono fun x => F x y

```

The deliberate omission is any field relating $pl(a^c, b)$ to $pl(a, b)$. That missing field is exactly R3. Two details are themselves checker-forced. Strict monotonicity of F must be restricted to $0 < y$, because the boundary axioms force $F(x, 0) = 0$ for all x , so the unrestricted axiom makes the structure *uninhabited* and every theorem about it vacuous. We guard against this permanently: `coxModelENNReal` is an explicit model on the non-Boolean chain $[0, \infty]$, with no appeal to excluded middle at the object level. Deciding $a = \top$ there is a constructor check, and the formalization supplies the computable Decidable instance witnessing this (`decEqTopENNReal`), so the axiom class is inhabited by a genuinely non-classical instance.

6.2 The sum rule is irreducible

Classically, additivity is *derived*: R3 turns disjunction into negated conjunction via De Morgan, and the product rule does the rest. Constructively that route is closed, since $a \vee b \neq \neg(\neg a \wedge \neg b)$ in general. And this is not an artifact of one proof strategy:

Theorem 15 (`modularity_irreducible, no_disjunction_functional`)

There is a five-element frame (the lower sets of the poset $o < a, o < b$ with a, b incomparable) and a monotone, normalized assignment q on it that is additive on disjoint joins yet not modular. Moreover no binary operation S on values satisfies $q(x \sqcup y) = S(q(x), q(y))$ for all x, y . The pairs $(\downarrow a, \downarrow a)$ and $(\downarrow a, \downarrow b)$ have identical value-pairs $(\frac{1}{2}, \frac{1}{2})$ but joins valued $\frac{1}{2}$ and 1.

A disjunction's plausibility is not a function of its disjuncts' plausibilities, unlike a conjunction's, which the `conj` field reduces to values via conditioning. So a constructive Cox theorem *must* posit the sum-rule half. Modularity is the posit, and Theorem 7 is its independent justification: modularity says exactly that the calculus is closed under mixing certainities.

6.3 The theorem, corrected and proved

Our original target statement asked for a strictly monotone regratuation $g : \mathbb{R} \rightarrow [0, \infty]$ with $g(0) = 0$. The checker refused, for a reason no referee had caught on paper: strict monotonicity forces $g(-1) < g(0) = 0$, impossible in $[0, \infty]$. The statement was not too hard. It was *unsatisfiable*, and the repair (strictness only on $[0, 1]$, where plausibilities live) is exactly the domain hygiene that Halpern's critique of the classical theorem turns on. The corrected statement is a theorem:

Theorem 16 (`constructive_cox`). *Every Cox model whose unconditional plausibility is modular regratuates to a valuation: there exist $v : \text{Valuation } \Omega$ and g strictly monotone on $[0, 1]$ with $g(0) = 0$, $g(1) = 1$, and $v(a) = g(pl(a, \top))$ for all a .*

The regratuation for the sum-rule half is undemanding ($g = \text{ENNReal.ofReal}$). The substance lives in the product-rule half, showing F regratuates to multiplication, which is Aczél's associativity theorem [Aczél 1966], a statement of pure analysis whose type never mentions Ω . The constructive/classical divide is therefore *entirely* in the sum rule. `mathlib` has no Aczél theorem, no Hölder-style embedding for ordered semigroups on intervals, and no Cauchy functional equation under monotonicity (only under continuity), so we build the analytic core from scratch ($\approx 1,100$ lines): a bundled `Scale` of Aczél hypotheses, a generator defined as the dyadic limit $g(b) = \bigcup_n (\text{count of } 2^n\text{-th roots under } b) / 2^n$, and the Hölder-style conclusion, a normalized, strictly monotone

additive generator and hence a strictly monotone multiplicative one, on the positive cone (`exists_mul_generator`, `aczelStatement_cone`). The Archimedean property is *derived* from continuity and positivity rather than assumed, and the interval form of divisibility is likewise a consequence of the intermediate value theorem (`exists_diag_eq`), though the bundled `Scale` takes divisibility as a hypothesis. Appendix A gives the construction. The residual analytic gap, continuity of the generator and extension below the unit, is stated honestly there and in Section 10, and is witnessed non-vacuously by the log-sum-exp archetype, for which the global continuous generator is exp.

7 Two Grounding Models

Two concrete models anchor the abstract theory. One shows every classical measure *contains* an intuitionistic probability. The other shows the intuitionistic theory cannot collapse back without solving the halting problem.

7.1 Measures through the interior operator

Gödel–McKinsey–Tarski interpret intuitionistic logic in modal S4, semantically: propositions are *open* sets, and $\Box$ is topological interior [Gödel 1933; McKinsey and Tarski 1944]. Our valuations are the quantitative face of that translation:

Theorem 17 (`Measure.toValuationOpens`). *Let μ be a probability measure on a space X . Then $U \mapsto \mu(U)$ on the frame $O(X)$ is a valuation. Equivalently, $v = \mu \circ \text{int}$ as a function of arbitrary sets, that is, $v(A) = P(\Box A)$.*

Modularity is $\mu(U \cup V) + \mu(U \cap V) = \mu(U) + \mu(V)$. The slack of an open U is then $1 - \mu(U) - \mu(\text{int } U^C)$, which is the measure of the boundary ∂U . *Slack is boundary mass.* (The general boundary identity is elementary measure theory. The development proves the complement sub-additivity inequality in general, `interiorMeasure_add_compl_le`, and mechanizes the boundary identity end to end in the Sierpiński model below.) The DS “ignorance” of Section 5 is, in this model, the measure charged to the frontier where a finite-precision observer can never decide membership, a structural quantity with no epistemic subject. This also exhibits the modal identification $\text{Bel} = P(\Box \cdot)$ of the DS literature [Ruspini 1986] arising from the logic itself. Theorem 6 is the converse in the finite case, and the general converse is the open frontier of Section 4.

7.2 The computability guard

The design constraint on every axiom set in this paper is that it must not silently force slack $\equiv 0$. Otherwise the “constructive” theory is Kolmogorov in disguise. Our guard is a model in which collapse is not merely false but *uncomputable*.

A semidecidable proposition is an open in the Sierpiński topology on `Prop`: verifiable by a finite computation, never refutable by one. On the three-open frame

$O(\mathbb{S}) = \{\perp, \text{halts}, \top\}$ we prove the Σ_1 asymmetry as a theorem, $\text{halts}^c = \perp$ (every nonempty open of $\mathbb{S}$ contains `True`, so nothing partially refutes a semidecidable event), and build the valuation from a genuine measure $p \cdot \delta_{\text{True}} + (1 - p) \cdot \delta_{\text{False}}$ through the bridge above. The slack computes end to end (`slack_eq_boundary`):

$$\text{slack}(\text{halts}) = 1 - p = \mu(\partial\{q \mid q\}) = \mu\{\text{False}\},$$

the mass of the point where the computation stays silent forever.

The guard itself is grounded in `mathlib`’s computability theory. For a code c and input n , the *sharp classical readout* assigns the slack-free valuation answering “does c halt on n ?” with certainty, weight 1 or 0 by the actual halting fact. The family is classically well defined, but no algorithm can read it off:

Theorem 18 (`sharpReadout_not_computable`). *Deciding whether the sharp classical readout assigns belief 1 to the halting event is exactly the halting problem, hence not computable* (`ComputablePred.halting_problem`).

Any axiom strong enough to force classical collapse therefore demands of its models an uncomputable epistemic state. Slack is not a defect the constructive theory should apologize for. It is the price of a computable one. This guard plays for the sum rule the role the explicit `CoxModel` instance plays for the product rule (Section 6): each axiom set in the paper ships with a machine-checked witness that it has non-degenerate models.

8 Formalization Design and Reuse

Shape of the development. Fifteen Lean 4 modules, about 4,300 lines, over a pinned `mathlib` (both at v4.31.0): the core valuation theory and GMT bridge, the DS layer (2-monotonicity, inclusion–exclusion and the ∞ -monotone tower, the two conditionings), the representation suite (finite, obstruction, Scott, general decomposition), the Cox layer (`CoxModel`, sum-irreducibility, the corrected theorem), the Aczél development, the two guards (halting chain, Sierpiński model), the product rigidity theorem on chain products, the countable-mixture composition laws, and the ultrafilter couplings that separate the iteration orders. The build is clean, with no sorry and no warnings, and every theorem named in this paper reports exactly [`propext`, `Classical.choice`, `Quot.sound`] under `#print axioms`.

Meta-logic against object logic. Lean’s logic is classical, while our *object* of study is the frame. This division of labor is deliberate and, we found, the pragmatic sweet spot. Constructivity claims attach to the algebra (no theorem assumes $a \sqcup a^c = \top$), while the meta-theory may reason classically *about* it. The counterexample valuations of Sections 4 and 6 use classical case splits legitimately, since a counterexample is a classical fact about the theory. Where the distinction has

content we track it: the non-vacuity witness for `CoxModel` was re-based from `Prop` (which imports excluded middle through the back door) to the chain $[0, \infty]$, where the needed case split is a constructor check, witnessed by a computable Decidable instance rather than the classical one, and the maximal-point peel behind the general decomposition was arranged to make no excluded-middle case split in its script.

The division also bites in a place the theory itself predicts. Every valuation in the development whose *definition* requires classical logic is an indicator: the point valuations, the prime-ideal indicators behind `sharp_iff_point`, the $\top$ -indicator, and the halting weights. An indicator into $[0, \infty]$ must decide membership, so the sharp states are exactly the objects a constructive meta-theory could not build. Likewise the hard direction of the R3 hinge rests on prime-ideal separation, a choice principle. The formalization thus enacts its own thesis one level up: certainty and points are classical resources, inside the theory and above it.

What the checker contributed. Beyond the usual assurance, five findings in this paper originate in failed proofs. The naive `constructive_cox` statement was unsatisfiable (Section 6). Unrestricted strict monotonicity for the conjunction functional was vacuous (Section 6). The regular-elements guess for the classical fragment was false (Section 2). Our first Aczél hypothesis bundle omitted order-preservation, which the additivity proof needs (Appendix A). And our expected counterexample to product uniqueness was refuted by the rigidity that became Section 10's theorem. We regard these as the paper's best evidence that this subject specifically repays mechanization.

mathlib surface. The development consumes order theory (`Order`, `Frame`, Heyting algebras, prime ideal separation), `ENNReal` arithmetic, measure theory for the bridge, PMF, and the computability library (`Nat.Partrec.Code`, `ComputablePred`). Candidates for upstreaming, in decreasing generality: the Aczél/Hölder generator construction of Appendix A (`mathlib` has no associativity-representation theorem, no monotone Cauchy functional equation, and no t -norm theory), valuations on frames with the mass/representation API, and the inclusion-exclusion identity for modular functions on distributive lattices (Theorem 11), which is independent of probability. $[0, \infty]$'s truncated subtraction deserves a note. It forces additive normal forms, with every identity stated so that no subtraction occurs. This reads as a constraint but functioned as a correctness discipline: several informal manipulations of the DS literature do not survive it unchanged.

Artifact. The artifact (anonymized supplement) builds with a two-command recipe on the pinned toolchain and re-checks every theorem, including the axiom audit.

9 Related Work

Informal precedents. That probability might be paired with intuitionistic logic is an old if scattered thought. Weatherston develops intuitionistic probability axiomatically and defends it with Dutch-book arguments [Weatherston 2003]. Paris treats non-classical uncertain reasoning broadly [Paris 1994]. The identification $\text{Bel} = P(\Box \cdot)$ connecting belief functions to modal logic is classical [Dempster 1967; Ruspini 1986; Shafer 1976]. On the localic side, valuations on frames are standard objects of constructive measure theory, with Vickers's work and the Coquand–Spitters development as canonical references [Coquand and Spitters 2009; Vickers 2008], normally with Scott continuity built in and with additive, Riesz-flavored goals. What we take from Section 4 is that Scott continuity is a substantive assumption (it decides point-representability on the frames we examine), so our base structure omits it. The Cox line, through Cox, Jaynes, Halpern's counterexample, and the Van Horn and Arnborg–Sjödín repairs [Arnborg and Sjödín 2000; Cox 1946; Halpern 1999; Horn 2003; Jaynes 2003], is classical throughout. Van Horn himself observes that dropping R3 yields a two-number theory, which is exactly the belief/plausibility pair our slack separates. To our knowledge, no prior work, informal or formal, states either hinge theorem: complement rule $\iff$ excluded middle (Theorem 4), or Dempster = Bayes $\iff$ zero slack at the evidence (Theorem 14).

Prior formalizations. Three bodies of mechanized work are adjacent, and none overlaps. ALEA [Audebaud and Paulin-Mohring 2009] formalizes a discrete distribution monad in Coq for randomized programs: additive probability, no event logic in our sense. Recent work in cubical Agda formalizes synthetic probability via Markov categories [Sargsyan et al. 2023], where again the objects are additive distributions. Closest is the HoTT/Coq line of Faissole and Spitters on *valuations* over lower reals [Faissole and Spitters 2017], done constructively with Scott continuity and aimed at integration (Riesz) and a Giry-style valuation *monad*. Their valuations are the additive, continuous cousins of ours, and their monad is the Scott-continuous instance of the question examined in Section 10. The delta of this paper is not the word “valuation” but the development on top of it: the non-additive complement regime, slack and its decomposition, both hinge theorems, the DS bridge with total monotonicity and the two conditionings, the representation/mixture/Scott-atomic suite, the corrected Cox theorem with its irreducibility counterexamples, the Aczél construction, product rigidity, and the computability guard. We claim first mechanization for that development, not for valuations on locales as such.

Non-Boolean probability elsewhere. Quantum probability on orthomodular lattices, with Gleason's theorem [Birkhoff and von Neumann 1936; Gleason 1957],

is the long-standing precedent for re-basing probability on a non-Boolean event logic. Free probability, fuzzy measures, and imprecise probability [Walley 1991] vary other parameters (independence, additivity, single-valuedness). Our variation is specifically the *logic*, holding the Cox-style methodology fixed. Within DS theory, the two conditionings and their inequivalence are classical [Fagin and Halpern 1991; Shafer 1976]. What is new in Theorem 14 is the exact logical characterization of their agreement, and its mechanization.

10 Conclusion: Probability with the Logic as a Parameter

We set out from the observation that Kolmogorov's axioms fix a logic before they fix an arithmetic, and that nothing in the concept of graded belief privileges the Boolean choice. The machine-checked results bear the framing out with a precision informal treatments have not reached. The entire classical surplus is one axiom, whose statical face is the complement rule (Theorem 4) and whose dynamical face is the agreement of Dempster and Bayesian conditioning (Theorem 14). Everything else survives the intuitionistic substrate intact: conditioning, Bayes symmetry, disjoint additivity, total probability over genuine partitions, and inclusion-exclusion itself (Theorem 11). What the weaker logic adds is structure, not damage: slack with its two-gap decomposition, the atomic/diffuse divide, the belief-function reading through $\neg\neg$, and a computability-theoretic reason the additions are not optional.

Open problems. Three are mathematical. First, representing the diffuse part on non-spatial frames (the localic Riesz direction). Second, closing the analytic residue of the Aczél development, namely continuity of the generator and the off-cone extension (the cone results and the log-sum-exp archetype are done). Third, a genuine uniqueness theorem: deriving modularity from a more primitive desideratum, knowing (Theorem 15) that it cannot come from disjunction data alone, with the mixture characterization (Theorem 7) as the current best justification. A fourth is foundational hygiene at scale: a machine-checked Dutch-book theorem in the style of [Weatherson 2003], which to our knowledge no proof assistant has for *any* logic.

The functor is literal, the monad is delimited. Within a fixed doctrine the connection to the categorical picture is concrete. Pushforward along frame homomorphisms preserves all four valuation axioms, making $\text{Val}(-)$ a functor on locales, the localic face of the functor part of the Giry monad [Giry 1982]. This is a routine verification that we state but have not mechanized. The point valuations δ_p of Section 4 are the evident unit (these *are* formalized), and Theorem 6 shows that on finite frames Val agrees, on objects, with the classical finite-distribution monad on points.

The *multiplication* is where the substance is. Flattening a valuation of valuations is integration, hence product valuations and Fubini, and by Kock's identification of commutative monads with Fubini theorems [Kock 2012], whether statistics-as-kernel-composition transports to a weaker logic is precisely the question whether that logic's valuation functor is a commutative monad. Two machine-checked results now delimit the multiplication from both sides.

On the chain product the question of *which* product valuation to take does not arise. The frame $\text{Low}(\mathbb{N} \times \mathbb{N})$ is *rigid*. Every lower set is a *finite* union of rectangles: beyond a stabilization height every occupied column is full, so the full columns and finitely many rows account for everything (exists_rect_decomposition). The rigidity mechanism is general: by the inclusion-exclusion equality, two valuations agreeing on any $\sqcap$ -closed family agree on all finite joins of its members (eq_on_finsetSup_of_eq_on), and the chain product is the instance where that family is the rectangles. So the product valuation there, whenever it exists, is unique, with no continuity assumed (Valuation.product_unique). We had expected the opposite, diffuse mass free to choose among the infinities of the product, and our candidate counterexamples all died on the same rock: lower sets of chain products detect only finitely much of the boundary.

They were right in spirit and wrong in venue. On the powerset frame $\text{Set}(\mathbb{N} \times \mathbb{N})$, whose elements are far from finite unions of rectangles, non-uniqueness is a theorem. The indicator of an ultrafilter is a valuation (ultrafilterValuation), and any ultrafilter on a product splits rectangles through its marginal pushforwards (rect_mem_iff). Taking a free ultrafilter $\mathcal{U}$ on $\mathbb{N}$, the two iteration orders of its tensor with itself and its diagonal pushforward all carry identical values on *every* rectangle, yet they differ on the upper triangle $\{(m, m') \mid m < m'\}$ (product_valuation_not_unique). In particular the two orders of iterated integration of the same marginal data disagree: *Fubini fails*, concretely (fubini_fails_for_valuations). Compare the Fubini pathologies of non-additive integration [Denneberg 1994; Ghirardato 1997]. By Kock's correspondence, no canonical commutative multiplication exists for unrestricted valuations. On the powerset frame these valuations are finitely additive probability charges in the sense of [Rao and Rao 1983], so this half of the dichotomy is not about intuitionism at all: the freedom lives already in the classical finitely additive world, and dropping continuity is what lets it in. Anyone who wants a monad must therefore restrict, and the dichotomy says along which axis: the frames, or the valuations.

The hard problem. The question *upstream* of the multiplication deserves to have its hardness named rather than implied, and what follows is our assessment rather than a

theorem. A monad needs a category for Val to be an endofunctor on. The space of valuations must itself be an object, a “space” whose opens one can speak of. For Scott-continuous valuations that space is a locale and monad structure is available, constructively, through the probabilistic powerdomain line [Coquand and Spitters 2009; Jones and Plotkin 1989; Vickers 2008]. Yet *even there*, exhibiting a cartesian closed category closed under the construction is the Jung–Tix problem, open since 1998 [Jung and Tix 1998]. Dropping continuity makes the difficulty prior rather than merely worse: the standard presentations of the valuation space use continuity essentially. One route to a well-posed question does exist. The localic treatments present the space of continuous valuations as the classifying locale of a propositional geometric theory [Coquand and Spitters 2009; Vickers 1989], with generators asserting $q < v(a)$ for rational q and axioms for monotonicity, normalization, modularity (expressible with rational witnesses and countable disjunctions), and continuity. Deleting the continuity axiom leaves a geometric theory, so a classifying locale still exists on general grounds, and classically its points are exactly the valuations of this paper. We have verified none of the structure this candidate would need. The unit looks unproblematic, but the multiplication requires integrating the evaluation maps against a merely modular, discontinuous valuation, and there the sum of two lower-real-valued maps is an infinite join, exactly where the diffuse part of Theorem 8 leaks. We offer the formulation as a target, not a result.

The strategy with the best modern track record is to carve a fragment rather than to conquer the whole. On the measure-theoretic side, the failure of cartesian closure for measurable spaces was answered by quasi-Borel spaces [Heunen et al. 2017]. On the domain-theoretic side, commutativity of the full valuations monad on DCPO is open, and the response is a hierarchy of commutative *submonads* generated from simple valuations [Jia et al. 2021]. Our decomposition theorem names the corresponding fragment here: the atomic part. As a first machine-checked step in that direction we prove that the countably-presented fragment composes with no space structure at all. A countable mixture of valuations is a valuation (`mixCountable`), a mixture of mixtures flattens to a mixture with product weights, which is discrete Chapman–Kolmogorov (`mixCountable_mixCountable`), and the one-point mixture is the identity (`mixCountable_unique`), all by the unconditional Fubini of $[0, \infty]$ sums over sigma types. And in exact contrast with the failure above, the fragment is *commutative*: iterated countable mixing is order-independent (`mixCountable_comm`). Commutative exactly where countable. This is associative, unital, commutative kernel composition for countably-atomic data. It is not yet a monad: the presentations are not canonical, we do not quotient by “presents the same valuation,” and the diffuse part is exactly what escapes every countable presentation. The ledger, then. The unit, the finite level, countable kernel composition, and

its commutativity are proved. Rigidity on chain products and non-uniqueness on the powerset frame are proved, so the general multiplication is settled: it is not canonically determined. What remains is the ambient category, a formulation problem before it is anything else.

Future work on the monad. Two continuations are concrete. First, quotient the countable presentations by “presents the same valuation” and package the result as a Kleisli category of countably-atomic kernels, checking that mixing descends to the quotient. That would upgrade composition-on-data to an honest categorical object, and we expect it to be a development of its own. Second, mechanize the classifying-locale formulation sketched above, which requires infrastructure `mathlib` currently lacks entirely: classifying locales of propositional geometric theories. Both continuations inherit the boundary drawn here. Whatever structure exists must live on a fragment, because the unrestricted multiplication is now provably not canonical.

Across doctrines, by contrast, we make no functoriality claim. There is no ambient category in which “logic $\mapsto$ its probability theory” is a functor, and each column requires re-deriving what a state *is* from the desiderata. What is uniform across columns is the *method*, and this paper’s execution of it is the Heyting column.

The program. If the logic is a parameter, this paper computed one new value of it. The table wants filling in. Linear logic is the most inviting column for this community. Its event algebras are quantales, its conjunction is resource-sensitive, and a “valuation” there should grade the *consumption* of evidence. The analogue of Gleason’s theorem or of our representation theorems (which states are compatible with the algebra?) is wide open, as is the right analogue of slack for the failure of contraction. The probabilistic semantics of linear logic that already exist, probabilistic coherence spaces [Danos and Ehrhard 2011], start from the additive side. A Cox-style derivation over a quantale would start from the logic. The methodology of this paper, to state the classical desiderata, delete the axioms the logic no longer licenses, and let a proof assistant referee what survives, what splits, and what becomes unsatisfiable, is, we suggest, the right instrument for that program. Each cell of the table is exactly the kind of delicate, counterexample-rich boundary work that mechanization is for.

References

- János Aczél. 1966. *Lectures on Functional Equations and Their Applications*. Academic Press.
- Stefan Arnborg and Gunnar Sjödin. 2000. Bayes Rules in Finite Models. *Proceedings of the European Conference on Artificial Intelligence (ECAI)* (2000), 571–575.
- Philippe Audebaud and Christine Paulin-Mohring. 2009. Proofs of Randomized Algorithms in Coq. In *Science of Computer Programming*, Vol. 74. 568–589.

Garrett Birkhoff and John von Neumann. 1936. The Logic of Quantum Mechanics. *Annals of Mathematics* 37, 4 (1936), 823–843.

Thierry Coquand and Bas Spitters. 2009. Integrals and Valuations. *Journal of Logic and Analysis* 1, 3 (2009), 1–22.

Richard T. Cox. 1946. Probability, Frequency and Reasonable Expectation. *American Journal of Physics* 14, 1 (1946), 1–13.

Vincent Danos and Thomas Ehrhard. 2011. Probabilistic Coherence Spaces as a Model of Higher-Order Probabilistic Computation. *Information and Computation* 209, 6 (2011), 966–991.

Arthur P. Dempster. 1967. Upper and Lower Probabilities Induced by a Multivalued Mapping. *Annals of Mathematical Statistics* 38, 2 (1967), 325–339.

Dieter Denneberg. 1994. *Non-Additive Measure and Integral*. Kluwer.

Ronald Fagin and Joseph Y. Halpern. 1991. A New Approach to Updating Beliefs. *Uncertainty in Artificial Intelligence* 6 (1991), 347–374.

Florian Faissolle and Bas Spitters. 2017. Synthetic Topology in Homotopy Type Theory for Probabilistic Programming. In *Mathematically Structured Functional Programming (MSFP), and: Preuves et Programmes (JFLA)*. See also Faissolle–Spitters, “Synthetic topology in HoTT for probabilistic programming,” MSCS 2021..

Paolo Ghirardato. 1997. On Independence for Non-Additive Measures, with a Fubini Theorem. *Journal of Economic Theory* 73, 2 (1997), 261–291.

Jean-Yves Girard. 1987. Linear Logic. *Theoretical Computer Science* 50, 1 (1987), 1–101.

Michèle Giry. 1982. A Categorical Approach to Probability Theory. In *Categorical Aspects of Topology and Analysis*. Lecture Notes in Mathematics, Vol. 915. Springer, 68–85.

Andrew M. Gleason. 1957. Measures on the Closed Subspaces of a Hilbert Space. *Journal of Mathematics and Mechanics* 6, 6 (1957), 885–893.

Kurt Gödel. 1933. Eine Interpretation des intuitionistischen Aussagenkalküls. *Ergebnisse eines mathematischen Kolloquiums* 4 (1933), 39–40.

Joseph Y. Halpern. 1999. A Counterexample to Theorems of Cox and Fine. *Journal of Artificial Intelligence Research* 10 (1999), 67–85.

Carl G. Hempel. 1945. Studies in the Logic of Confirmation (I). *Mind* 54, 213 (1945), 1–26.

Chris Heunen, Ohad Kammar, Sam Staton, and Hongseok Yang. 2017. A Convenient Category for Higher-Order Probability Theory. In *Logic in Computer Science (LICS)*.

Kevin S. Van Horn. 2003. Constructing a Logic of Plausible Inference: A Guide to Cox’s Theorem. *International Journal of Approximate Reasoning* 34, 1 (2003), 3–24.

Edwin T. Jaynes. 2003. *Probability Theory: The Logic of Science*. Cambridge University Press.

Xiaodong Jia, Bert Lindenhovius, Michael Mislove, and Vladimir Zamdzhiev. 2021. Commutative Monads for Probabilistic Programming Languages. In *Logic in Computer Science (LICS)*.

Claire Jones and Gordon D. Plotkin. 1989. A Probabilistic Powerdomain of Evaluations. In *Logic in Computer Science (LICS)*. 186–195.

Achim Jung and Regina Tix. 1998. The Troublesome Probabilistic Powerdomain. *Electronic Notes in Theoretical Computer Science* 13 (1998), 70–91. Proceedings of Comprou III. DOI 10.1016/S1571-0661(05)80216-6..

Anders Kock. 2012. Commutative Monads as a Theory of Distributions. *Theory and Applications of Categories* 26 (2012), 97–131.

The mathlib Community. 2020. The Lean Mathematical Library. In *Certified Programs and Proofs (CPP)*. 367–381.

J. C. C. McKinsey and Alfred Tarski. 1944. The Algebra of Topology. *Annals of Mathematics* 45, 1 (1944), 141–191.

Jeff B. Paris. 1994. *The Uncertain Reasoner’s Companion: A Mathematical Perspective*. Cambridge University Press.

K. P. S. Bhaskara Rao and M. Bhaskara Rao. 1983. *Theory of Charges: A Study of Finitely Additive Measures*. Pure and Applied Mathematics, Vol. 109. Academic Press.

Enrique H. Ruspini. 1986. The Logical Foundations of Evidential Reasoning. In *Technical Note 408, SRI International*.

Gagik Sargsyan et al. 2023. Synthetic Probability Theory via Markov Categories in Cubical Agda. Preprint. Verify exact citation before submission..

Glenn Shafer. 1976. *A Mathematical Theory of Evidence*. Princeton University Press.

Steven Vickers. 1989. *Topology via Logic*. Cambridge University Press.

Steven Vickers. 2008. A Localic Theory of Lower and Upper Integrals. In *Mathematical Logic Quarterly*. Vol. 54. 109–123.

Peter Walley. 1991. *Statistical Reasoning with Imprecise Probabilities*. Chapman and Hall.

Brian Weatherston. 2003. From Classical to Intuitionistic Probability. *Notre Dame Journal of Formal Logic* 44, 2 (2003), 111–123.

A The Aczél Development

This appendix sketches the from-scratch construction behind the product-rule half of Theorem 16. The body stands without it. The target shape is Aczél’s associativity theorem [Aczél 1966]: a suitably regular associative $F : \mathbb{R} \rightarrow \mathbb{R} \rightarrow \mathbb{R}$ reg graduates to multiplication, $G(F x y) = G(x) \cdot G(y)$ for a strictly monotone G . mathlib offers no route. It has no associativity representation, no Hölder-style embedding of ordered semigroups on intervals, and the Cauchy functional equation only under continuity, not monotonicity.

Hypotheses. A Scale bundles: associativity of F , divisibility $\forall c \exists x, F x x = c$, positivity on the cone $x < F x c$, joint order-preservation, joint continuity, a unit u , and identity at $-\infty$, meaning $F x z \rightarrow x$ as $z \rightarrow -\infty$. Order-preservation was absent from our first bundle, and the additivity proof needs it, a definition-check finding (Section 8). Divisibility is taken as a hypothesis in the bundle, although its interval form is a consequence of the intermediate value theorem (exists_diag_eq). Non-vacuity: $\log\text{-sum-exp}$, $F x y = \log(e^x + e^y)$ with $u = 0$, satisfies every field (logSumExpScale), and for it the intended generator is globally exp.

Construction. Write $a^{(n)}$ for n right-associated copies of a under F (Fpow). Associativity alone makes $n \mapsto a^{(n)}$ a semigroup morphism from $(\mathbb{N}, +)$, with no commutativity needed. Commutativity is a *corollary* of the generator, not a prerequisite. One fact usually assumed is here *derived*: unboundedness of the iterates. Were they bounded, monotone convergence would give a fixed point $F L a = L$, contradicting positivity, and unboundedness yields the Archimedean property. Divisibility supplies a tower of 2^n -th roots $u \succ u^{1/2} \succ u^{1/4} \succ \dots$ (droot) with the coherence law that 2^ℓ fine copies recombine one coarse mark. Counting how many copies of the n -th root fit under b (dcount, well defined by the Archimedean property) gives dyadic approximants $g_n(b) = \text{dcount}(b, n)/2^n$. The refinement sandwich $2N + 1 \leq N' \leq 2N + 2$ makes them monotone and Cauchy, and the generator is their limit, $g(b) = \bigsqcup_n g_n(b)$.

Results (all sorry-free, standard axioms). On the cone $\{x \mid u \leq x\}$ we prove three things. Normalization: $g(u) = 1$. Additivity: $g(F x y) = g(x) + g(y)$, by squeezing super- and

sub-additive counting bounds ($\text{dcount } x + \text{dcount } y + 1 \leq \text{dcount}(F x y) \leq \text{dcount } x + \text{dcount } y + 3$) against a vanishing 2^{-n} slack. And *strict* monotonicity, proved without any continuity of g by a count-amplification argument: a shrinking root inserted between $x < y$ contributes a full extra block of 2^{n-M} copies. Exponentiating gives the multiplicative form, a strictly monotone G with $G(F x y) = G(x) G(y)$ on the cone (`exists_mul_generator`), which is the conclusion shape of Aczél's theorem restricted to the cone (`aczelStatement_cone`). A bounded reorientation $\bar{G} = \exp(-g)$ transports it to the $[0, 1]$, conjunctive orientation of Section 6.

Honest residue. The constructed g is discontinuous at the unit (it is 0 below u), so matching the classical statement verbatim still needs the off-cone extension (a group completion) and continuity of the extension. That is genuine analysis, open here, but *witnessed*: for the log-sum-exp archetype the global continuous generator exists (`hasOrderedGenerator_logSumExp`). Of Hölder's classical hypotheses, the Archimedean property is a theorem in this development rather than an assumption, and the interval form of divisibility follows from the intermediate value theorem. Only the embedding's global regularity remains.