

Outline for Topological Combinatorics

The premise of this talk would be to discuss some of the basics of topology and then eventually state (and prove, maybe¹) the Borsuk-Ulam Theorem, Sperner's Lemma, and Brouwer's fixed point theorem. Following which, we present a variety of applications in combinatorics. Most of the material comes from [1].

Topological Results:

Theorem (Brouwer Fixed Point Theorem)

Let $K \subseteq \mathbb{R}^n$ be convex and compact, and let $f : K \rightarrow K$ be continuous. Then, there exists $k \in K$ such that $f(k) = k$.

Theorem (Borsuk-Ulam Theorem)

Suppose $f : \mathbb{S}^n \rightarrow \mathbb{R}^n$ was continuous. Then, there is some $x \in \mathbb{S}^n$ such that $f(x) = f(-x)$.

Combinatorial Results:

Theorem

Every (open) necklace with d kinds of jewels can be divided between two thieves using no more than d cuts.

Theorem (Kneser Conjecture)

For all $k > 0$ and $n \geq 2k - 1$, the chromatic number of the Kneser graph $KG_{n,k}$ is $\chi(KG_{n,k}) = n - 2k + 2$.

The game of Hex is played between two players, red and green. Each of them takes turns coloring one hexagon in a grid. After all hexagons are filled, red wins if there is a red path connecting the top and bottom, and green wins if there is a green path connecting the left and right.

Theorem

The game of Hex cannot end in a draw.

¹For the sake of time, we might limit the discussion of Borsuk-Ulam to a proof sketch.

References

- [1] Jiří Matoušek, Anders Björner, Günter M Ziegler, et al. *Using the Borsuk-Ulam theorem: lectures on topological methods in combinatorics and geometry*. Springer, 2003.