

6

**Susan Porter**

---

**From:** diannehu@optusnet.com.au  
**Sent:** Tuesday, 26 June 2012 6:20 PM  
**To:** Print Repository; library.fishill@sydney.edu.au  
**Subject:** University of Sydney: Storage Canberra request form

The form below was submitted by  
([diannehu@optusnet.com.au](mailto:diannehu@optusnet.com.au)) on Tuesday, June 26, 2012 at 18:20:10

---

email: [diannehu@optusnet.com.au](mailto:diannehu@optusnet.com.au)

call number: DST03389

author: C. R. Forrester

item title: Journal of the Fisheries Research Board of Canada,

article\_chapter title: Sinistrality in Platichthys stellatus off British Columbia

year: 1969

publication details: Vol. 26, No. 1 : pp. 191-196

(doi: 10.1139/f69-023)

name: Dianne Hughes

phone: 96671879 0423 804 470

borrower number: 009120549020908

faculty: The Australian Centre for Microscopy & Microanalysis

pickup\_library: Badham

Copyright: I agree with the Copyright declaration

---

HTTP\_USER\_AGENT: Mozilla/5.0 (Windows NT 6.1; WOW64; rv:12.0) Gecko/20100101 Firefox/12.0  
REMOTE\_ADDR: 211.30.226.215

Commonwealth of Australia

*Copyright Act 1968*

**Notice for paragraph 49 (7A) (c) of the *Copyright Act 1968***

**Warning**

This material has been provided to you under section 49 of the *Copyright Act 1968* (the *Act*) for the purposes of research or study. The contents of the material may be subject to copyright protection under the Act.

Further dealings by you with this material may be a copyright infringement. To determine whether such a communication would be an infringement, it is necessary to have regard to the criteria set out in Division 3 of Part III of the Act.

**orientalis in British  
es**

f Canada  
no, B.C.  
  
orientalis in British Columbia bivalves.  
  
Mytilicola orientalis has little economic  
of natural infestation of Saxidomus

dori is found in the recurrent  
ote summarizes a study of the  
acific oyster Crassostrea gigas  
Linné for the period 1963-67.  
  
icance of Mytilicola infestation  
952) attributed heavy losses of  
(1946) showed that infestation  
CF) of Ostrea lurida. It has not  
copepod is invariably associat-  
ster, for calculations of the CF  
Ladysmith Harbour failed to

metaplastic changes occurred  
er with a tendency to fibrosis  
nce of gross damage has been  
ever, in oysters that have been  
endency to evacuate contained  
hat the thoracic protuberances  
  
the immediate vicinity where  
ult stock, has been introduced.  
e the oyster population is con-

a single extended reproductive  
situation being similar to that  
intestinalis in Germany. Experi-  
arval stages are short and do

not travel far. Oysters and mussels raised above the substrate are not infested.  
The most frequently infested host is *M. edulis*, followed by *C. gigas* and  
*O. lurida*. Less often *M. californianus* (Chew et al., 1964) and the butter clam  
*Saxidomus giganteus* may be host to the copepod.  
Up to the present there are no indications that *M. orientalis* is of any  
economic significance in the British Columbia shellfish industry.

REFERENCES

BERNARD, F. R. MS, 1968. Incidence and effect of the copepod *Mytilicola orientalis* Mori in the Pacific oyster of British Columbia. Fish. Res. Bd. Canada MS Rept. 993. 14 p.  
CHEW, K. K., A. K. SPARKS, AND S. C. KATKANSKY. 1964. First record of *Mytilicola orientalis* Mori in the California mussel *Mytilus californianus* Conrad. J. Fish. Res. Bd. Canada 21: 205-207.  
KORINGA, P. 1950. De aanval van de parasiet *Mytilicola intestinalis* op de Zeeuwse Mossel cultuur. Vissevig-Nieuws 7.  
1952. Epidemiological observations on the mussel parasite *Mytilicola intestinalis* Steur, carried out in the Netherlands, 1951. Ann. Biol. Copenhagen 8: 182-185.  
MEYER, P. F., AND H. MANN. 1951. Recherches allemandes relative au "*Mytilicola*" copepode parasite de la moule, existent dans les Whatten allemands 1950-51. Rev. Trav. Office Sci. Tech. Pêches Maritimes 17: 63-75.  
ODLAUG, T. O. 1946. The effect of the copepod *Mytilicola orientalis* upon the Olympia oyster, *Ostrea lurida*. Trans. Am. Microscop. Soc. 65: 311-317.  
SPARKS, A. K. 1962. Metaplasia of the gut of the oyster *Crassostrea gigas* (Thunberg) caused by infection with the copepod *Mytilicola orientalis* Mori. J. Insect Pathol. 4: 57-62.

**Sinistrality in *Platichthys stellatus* off British Columbia**

C. R. FORRESTER

Fisheries Research Board of Canada  
Biological Station, Nanaimo, B.C.

FORRESTER, C. R. 1969. Sinistrality in *Platichthys stellatus* off British Columbia. J. Fish. Res. Bd. Canada 26: 191-196.

Percentage sinistrality in the starry flounder (*Platichthys stellatus*) off British Columbia was found to increase proceeding from south to north, and on the average (66%) was greater than that found off the western coast of the United States, and close to that found off Alaska.

Received September 16, 1968

FLATFISHES OF the family Pleuronectidae in the North Pacific are normally dextral, i.e., have eyes and coloration on the right side of the fish. Most frequent exceptions to this pattern occur in the starry flounder, *Platichthys stellatus*,

Printed in Canada

and several authors have shown that reversal in this species is widespread (Norman, 1934; Gudger, 1935; Townsend, 1937). Summaries of sampling conducted from Monterey Bay, California, north through the Gulf of Alaska and off Japan showed that incidence of sinistral specimens of *P. stellatus* ranged from about 50% off Oregon to 100%, or close to 100%, off Japan (Hubbs and Kuroshima, 1942; Orcutt, 1950).

Information on sinistrality or "handedness" in flounders from the British Columbia coast has been collected at intervals since 1945. This previously unpublished material shows that the percentage of sinistral specimens off British Columbia (66.4%) lies above that found for samples taken off Oregon and close to that for samples taken off Kodiak Island and the Alaska Peninsula (Table 1).

TABLE 1. Percentage occurrences<sup>a</sup> of sinistral specimens of *Platichthys stellatus* in the North Pacific Ocean.

Locality	No. of specimens	Sinistral (%)
California	509	55.2
California (Orcutt, 1950)	1439	59.5
Oregon outer coast	65	49.2
Columbia River mouth	225	60.4
Washington outer coast	247	56.3
Puget Sound	8972	51.6
British Columbia	7671	66.4
Southeastern Alaska	2498	58.2
Kodiak Island and Alaska Peninsula	5129	68.0
Japan	476	100.0

<sup>a</sup>All data except that by Orcutt (1950) for California and that for British Columbia are from table 5 of Hubbs and Kuroshima (1942).

However, use of a single figure to describe sinistrality for a large region may be misleading. Off British Columbia there is a general trend to a higher percentage of sinistral specimens proceeding from southernmost to northern areas (Fig. 1). Off the Fraser River estuary in the southern Strait of Georgia (about 49°N lat) samples of *P. stellatus* are about 51% sinistral, virtually identical to the percentage found in samples from Puget Sound immediately to the south. On the Two Peaks ground of northern Hecate Strait (54°20'N lat) samples are about 78% sinistral.

The most anomalous data in Fig. 1 are those from grounds off the west coast of Vancouver Island (about 49°N lat). However, there is more reason to consider the west coast samples in conjunction with offshore samples rather than with the inshore and Strait of Georgia samples. In this context the mean of 65.4% for the west coast lies between the 50-60% sinistral observed off the Oregon and Washington coasts, respectively, and the 76% for all of Hecate Strait.




FIG. 1. Percentage sinistral specimens of *Platichthys stellatus* off British Columbia. Data from various sources.

The British Columbia variation within an area of p (2) the between-area variation intervals (percentages were within-year variation in an

Analysis of variance on year variations in an area we but neither between-year no (0.05). All four tests of betw between areas. There were c of Vancouver Island and H Vancouver Island and the Strait and the Fraser River the Fraser River estuary. Cl by length-groups within san total chi-squares were not differential viability between

For the tests to be co between areas should repr exist between areas it migh

sal in this species is widespread (1937). Summaries of sampling north through the Gulf of Alaska al specimens of *P. stellatus* ranged to 100%, off Japan (Hubbs and

ss" in flounders from the British vals since 1945. This previously e of sinistral specimens off British mples taken off Oregon and close d the Alaska Peninsula (Table 1).

ns of *Platichthys stellatus* in the North

S	Sinistral (%)
	55.2
	59.5
	49.2
	60.4
	56.3
	51.6
	66.4
	58.2
	68.0
	100.0

and that for British Columbia are from

strality for a large region may be eral trend to a higher percentage nmost to northern areas (Fig. 1). trait of Georgia (about 49°N lat) virtually identical to the percenti ately to the south. On the Two 0'N lat) samples are about 78%

those from grounds off the west However, there is more reason to on with offshore samples rather mples. In this context the mean 0-60% sinistral observed off the and the 76% for all of Hecate




FIG. 1. Percentage sinistrality in *Platichthys stellatus* among samples collected off British Columbia. Data from west coast of Vancouver Island are shown by open circles.

The British Columbia samples were examined for: (1) the between-year variation within an area of percentage sinistral by comparable length intervals; (2) the between-area variation in percentage sinistral by comparable length intervals (percentages were transformed to angles prior to calculation); (3) within-year variation in an area of percentage sinistral by size groups.

Analysis of variance on seven sets of within-area data showed that between-year variations in an area were greater than between-length variations (Table 2) but neither between-year nor between-length variations were significant ( $P < 0.05$ ). All four tests of between-area variations showed significant differences between areas. There were differences between samples taken off the west coast of Vancouver Island and Hecate Strait, between those from the west coast of Vancouver Island and the Fraser River estuary, between those from Hecate Strait and the Fraser River estuary, and between those from Cape Lazo and the Fraser River estuary. Chi-square tests on the numbers sinistral and dextral by length-groups within samples showed no apparent trend within samples and total chi-squares were not significant ( $P < 0.05$ ; Table 4). This suggests no differential viability between dextral and sinistral forms of *P. stellatus*.

For the tests to be completely meaningful, the length groups compared between areas should represent the same year-classes. If growth differences exist between areas it might weaken the comparison. Further, comparisons of

TABLE 3. Percentage sinistral by size groups in *Platichthys stellatus* off British Columbia. Number in a size group was 20 or greater.

	Lower west coast Vancouver Is.	Johnstone Strait	Fraser R.	Cape Lazo	Upper Hecate Strait
Midpoint of length,	1026 1027 1028 1029 1030 1031	1020 1021 1022 1023 1024	1027 1029	1022 1025 1031 1037 1061	1016 1017 1061 1067 1063 1064

sinistrality within an area to determine differences in viability should be made on fish of the same year-class or classes in successive time intervals. The sampling was not intensive enough to permit such a study. Although the conclusions are tentative, they do suggest the presence of a number of discrete local stocks of *P. stellatus*.

n-year and between-area variations in off British Columbia.

f	df	F ratio
ars	3	1.06
gths	4	0.31
L)	12	
ars	1	1.51
gths	3	0.21
L)	3	
ars	1	3.75
gths	3	0.32
L)	3	
ars	2	1.93
gths	3	0.25
L)	6	
ars	1	0.55
gths	2	0.09
L)	2	
ars	1	0.26
gths	3	0.32
L)	3	
ars	1	1.35
gths	4	0.96
L)	4	
as	1	11.27
gths	7	0.55
L)	7	
as	1	15.33
gths	4	2.35
L)	4	
as	1	31.49
gths	4	2.36
L)	4	
as	1	9.01
gths	4	4.91
L)	4	

ces in viability should be made ve time intervals. The sampling r. Although the conclusions are nber of discrete local stocks of

TABLE 3. Percentage sinistral by size groups in *Platichthys stellatus* off British Columbia. Number in a size group was 20 or greater.

Midpoint of length interval (cm)	Lower west coast Vancouver Is.					Johnstone Strait					Fraser R.			Cape Lazo					Upper Hecate Strait				
	1955	1959	1960	1961	1962	1964	1960	1961	1964	1957	1962	1955	1958	1961	1962	1964	1946	1947	1961	1962	1963	1964	
32																							
35																							
38																							
41																							
44																							
47																							
50																							
53																							
56																							
59																							
62																							
65																							
68																							
No. of fish	117	273	414	176	169	125	285	645	155	177	262	209	144	199	134	350	215	249	679	374	529	166	

TABLE 4. Results of chi-square tests of numbers sinistral and dextral in various length groups of *Platichthys stellatus*.

Area	Year	No. of length intervals <sup>a</sup>	No. of fish	Chi-square	df
Lower west coast of Vancouver Is.	1955	5	117	3.79	4
	1959	7	273	5.24	6
	1960	8	414	2.01	7
	1961	5	176	1.70	4
	1962	5	169	1.89	4
Johnstone Strait	1964	4	125	1.03	3
	1960	7	285	4.03	6
	1961	8	645	7.65	7
Fraser R.	1964	4	155	4.13	3
	1957	4	177	0.69	3
Cape Lazo	1962	6	262	4.23	5
	1955	6	209	9.90	5
	1958	4	144	4.20	3
	1961	4	199	2.21	3
	1962	4	134	2.46	3
Upper Hecate Strait	1964	6	350	6.97	5
	1946	4	215	1.08	3
	1947	5	249	3.09	4
	1961	8	679	3.74	7
	1962	6	374	9.47	5
	1963	9	529	12.11	8
	1964	5	166	4.45	4

<sup>a</sup>See Table 3.

*Acknowledgments*—The assistance of Dr L. V. Pienaar in the statistical treatment is greatly appreciated.

## REFERENCES

- GUDGER, E. W. 1935. Abnormalities in flatfishes (Heterosomata). J. Morphol. 58: 1-39.
- HUBBS, C. L., AND K. KURONUMA. 1942. Hybridization in nature between two genera of flounders in Japan. Mich. Acad. Sci. Papers (1941) 27: 267-306.
- NORMAN, J. R. 1934. A systematic monograph of the flatfishes (Heterosomata). Vol. 1. Psettodidae, Bothidae, Pleuronectidae. British Museum, London. viii + 459 p.
- ORCUTT, H. G. 1950. The life history of the starry flounder, *Platichthys stellatus* (Pallas). Calif. Fish Game Fish Bull. 78. 64 p.
- TOWNSEND, L. D. 1937. Geographical variation and correlation in Pacific flounders. Copeia 1937: 92-103.

Recent Articles  
inStudies  
No.

1263. PARKER, ROBERT R. 1968. Bull. Brit. Museum (Nat. Hist.) 19: 1-11.
1264. IDLER, D. R., AND P. WISEMAN. 1968. The king crab and the North Pacific. J. Nat. Hist. Museum 19: 1113-1117.
1265. ACKMAN, R. G. 1968. The breakdown and fundamental aspects of the LDD and DDD disease. Biochem. Biophys. Res. Commun. 31: 1-11.
1266. ELDJARN, L., K. TRY, R. G. ACKMAN, AND R. G. ACKMAN. 1968. The LDD and DDD disease. Biochem. Biophys. Res. Commun. 31: 1-11.
1267. MARGOLIS, L. 1968. Review of the genus *Rhabdochona* (Nematoda: Rhabdochonidae). Museum 2: 23-44.
1268. RICKER, W. E., RENE MALOUF, AND J. G. MALOUF. 1968. The stoneflies of Quebec (Plecoptera). Bull. Brit. Museum (Nat. Hist.) 19: 1-11.

Copies of individual reprints, in the form of microfiche, are available from the establishments through which the individual reprints are distributed. Reprints of these articles are in the form of microfiche. These are distributed to a select group of libraries in the Studies for 1968, Part 1.)