

# ASSIGNMENT-7

P.Kalpana

## 1 QUESTION NO-2.71A (QUADRATIC FORMS)

Find the coordinates of the foci, the vertices, the length of major axis, the minor axis, the eccentricity and the length of the latus rectum of the ellipse

$$\mathbf{x}^T \begin{pmatrix} \frac{1}{36} & 0 \\ 0 & \frac{1}{16} \end{pmatrix} \mathbf{x} = 1.$$

## 2 SOLUTION

Given equation of the ellipse,

$$\mathbf{x}^T \begin{pmatrix} \frac{1}{36} & 0 \\ 0 & \frac{1}{16} \end{pmatrix} \mathbf{x} = 1 \quad (2.0.1)$$

we have,

$$\mathbf{V} = \begin{pmatrix} \frac{1}{36} & 0 \\ 0 & \frac{1}{16} \end{pmatrix} \quad (2.0.2)$$

$$\mathbf{u}^T \mathbf{V}^{-1} \mathbf{u} - f = 1 \quad (2.0.3)$$

$$\mathbf{c} = -\mathbf{V}^{-1} \mathbf{u} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad (2.0.4)$$

$$\lambda_1 = \frac{1}{36}, \lambda_2 = \frac{1}{16} \quad (2.0.5)$$

Axes of ellipse is given by: Length of semi major axis, a is

$$a = \sqrt{\frac{\mathbf{u}^T \mathbf{V}^{-1} \mathbf{u} - f}{\lambda_1}} \quad (2.0.6)$$

substituting the values in (2.0.6), we get

$$a = 6 \quad (2.0.7)$$

Length of major axis is  $2a = 12$   
and the length of semi minor axis, b is

$$b = \sqrt{\frac{\mathbf{u}^T \mathbf{V}^{-1} \mathbf{u} - f}{\lambda_2}} \quad (2.0.8)$$

substituting the values in (2.0.8), we get

$$b = 4 \quad (2.0.9)$$

Length of the minor axis is  $2b = 8$

The vertices are given as

$$\pm \begin{pmatrix} 6 \\ 0 \end{pmatrix} \quad (2.0.10)$$

Coordinates of foci are given by,

$$\mathbf{F} = \pm \left( \sqrt{\frac{(\mathbf{u}^T \mathbf{V}^{-1} \mathbf{u} - f)(\lambda_2 - \lambda_1)}{\lambda_1 \lambda_2}} \right) \mathbf{p}_1 \quad (2.0.11)$$

where,  $\mathbf{p}_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$  since the equation of ellipse is in standard form. Substituting the values in (2.0.11) we have,

$$\mathbf{F} = \pm \begin{pmatrix} 2\sqrt{5} \\ 0 \end{pmatrix}. \quad (2.0.12)$$

Eccentricity of the ellipse is given by,

$$e = \frac{\sqrt{\frac{(\mathbf{u}^T \mathbf{V}^{-1} \mathbf{u} - f)(\lambda_2 - \lambda_1)}{\lambda_1 \lambda_2}}}{\sqrt{\frac{\mathbf{u}^T \mathbf{V}^{-1} \mathbf{u} - f}{\lambda_1}}} \quad (2.0.13)$$

substituting the values in (2.0.13), we have

$$e = \frac{\sqrt{5}}{3}. \quad (2.0.14)$$

Length of the latus rectum is given by,

$$l = \frac{2 \left( \sqrt{\frac{f - \mathbf{u}^T \mathbf{V}^{-1} \mathbf{u}}{\lambda_2}} \right)^2}{\sqrt{\frac{\mathbf{u}^T \mathbf{V}^{-1} \mathbf{u} - f}{\lambda_1}}} \quad (2.0.15)$$

substituting the values in (2.0.15), we have

$$l = \frac{4}{3} \quad (2.0.16)$$

The plot of the ellipse is given below




Fig. 2.1: Plot of standard ellipse