

Assignment-17

Pooja H
AI20MTECH14003

Abstract—In this document, we solve for basis of the vector space of a transformation matrix.

Download all latex-tikz codes from

https://github.com/poojah15/EE5609_AI20MTECH14003/tree/master/Assignment_17

Hence, matrix of T is:

$$\begin{pmatrix} 2 & 0 & 2 & 0 \\ 0 & 2 & 0 & 6 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 2 \end{pmatrix} \quad (2.0.6)$$

3 ANSWERS FOR DIFFERENT CASES

1 PROBLEM STATEMENT

Let \mathbf{P}_3 be the vector space of polynomials with real coefficients and of at most degree 3. Consider the linear map $T : \mathbf{P}_3 \rightarrow \mathbf{P}_3$ defined by $T(p(x)) = p(x+1) + p(x-1)$. Which of the following does the matrix of T (with respect to the standard basis $\mathbf{B} = \{1, x, x^2, x^3\}$ of \mathbf{P}_3) satisfy?

- 1) $\det(T) = 0$
- 2) $(T - 2\mathbf{I})^4 = 0$ but $(T - 2\mathbf{I})^3 \neq 0$
- 3) $(T - 2\mathbf{I})^3 = 0$ but $(T - 2\mathbf{I})^2 \neq 0$
- 4) 2 is an eigenvalue with multiplicity 4.

2 SOLUTION

Given

$$T(p(x)) = p(x+1) + p(x-1). \quad (2.0.1)$$

The matrix of T with respect to the standard basis $\mathbf{B} = \{1, x, x^2, x^3\}$ is given by:

$$\begin{aligned} p(x) = 1 &\implies T(1) = 1 + 1 \\ &= 2 \end{aligned} \quad (2.0.2)$$

$$\begin{aligned} p(x) = x &\implies T(x) = x + 1 + x - 1 \\ &= 2x \end{aligned} \quad (2.0.3)$$

$$\begin{aligned} p(x) = x^2 &\implies T(x^2) = (x+1)^2 + (x-1)^2 \\ &= 2 + 2x^2 \end{aligned} \quad (2.0.4)$$

$$\begin{aligned} p(x) = x^3 &\implies T(x^3) = (x+1)^3 + (x-1)^3 \\ &= 6x + 2x^3 \end{aligned} \quad (2.0.5)$$

$\det(T) = 0$	False. From (2.0.6), it is found that the determinant is not zero as the eigenvalues are nonzero.
$(T - 2\mathbf{I})^4 = 0$ but $(T - 2\mathbf{I})^3 \neq 0$	False. $(T - 2\mathbf{I}) = \begin{pmatrix} 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 6 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$ $\implies (T - 2\mathbf{I})^2 = 0$ and hence $(T - 2\mathbf{I})^4 = 0$ and $(T - 2\mathbf{I})^3 = 0$
$(T - 2\mathbf{I})^3 = 0$ but $(T - 2\mathbf{I})^2 \neq 0$	False. Because $(T - 2\mathbf{I})^3 = 0$ and $(T - 2\mathbf{I})^2 = 0$
2 is an eigenvalue with multiplicity 4.	True. It is noted that the matrix of T is an upper triangular matrix having the value 2 along its principal diagonal and hence 2 is an eigenvalue with algebraic multiplicity 4.