

NAME:	Pooja Patil
UID:	2022301011
SUBJECT	Design and Analysis of Algorithms
EXPERIMENT NO :	1A
DATE OF PERFORMANCE	30.01.2023
DATE OF SUBMISSION	05.02.2023
AIM:	<p>To implement the various functions e.g., linear, non-linear, quadratic, exponential etc.</p> <p>1) Print the values of each function value for all n starting 0 to 100 in tabular format for both aforementioned cases</p> <p>2) Draw two 2D plot of all functions such that x-axis represents the values of n and y-axis represent the function value for different n values using LibreOffice Calc/MS Excel.</p>
THEORY	<p>A function is a relation between a set of inputs and a set of permissible outputs with the property that each input is related to exactly one output. Let A & B be any two non-empty sets; mapping from A to B will be a function only when every element in set A has one end, only one image in set B.</p> <p>1) n</p> <p>2) n^3</p> <p>3) $\log n$</p> <p>4) $n \cdot 2^n$</p> <p>5) $\log (\log n)$</p>

	6) 2^n 7) e^n 8) $3/2.n$ 9) $(\log n)^{1/2}$ 10) $n.(\log n)$
ALGORITHM	<p>➤ Initialize variables n and result.</p> <p>1. n Take the value of n from 0-100 and print all the values.</p> <p>2. n^3</p> <ul style="list-style-type: none"> • $result = n * n * n$ • Apply a for loop for values of n from 0-100 and print all the values for result. <p>3. $\log(n)$</p> <ul style="list-style-type: none"> • $result = \log(n)$ • Apply a for loop for values of n from 0-100 and print all the values for result. <p>4. $n * 2^n$</p> <ul style="list-style-type: none"> • $result = n * \text{pow}(2, n)$ • Apply a for loop for values of n from 0-100 and print all the values for result. <p>5. $(3/2)^n$</p> <ul style="list-style-type: none"> • $result = \log(\log(n))$ • Apply a for loop for values of n from 0-100 and print all the values for result.

6. e^n

- `result = pow(2,n)`
- Apply a for loop for values of n from 0-100 and print all the values for result.

7. 2^n

- `result = exp(n)` (e^n)
- Apply a for loop for values of n from 0-100 and print all the values for result.

8. $\lg(\lg n)$

- `result = 3/2*n`
- Apply a for loop for values of n from 0-100 and print all the values for result.

9. $(\log n)^{1/2}$

- `result = pow(log(n),0.5)`
- Apply a for loop for values of n from 0-100 and print all the values for result.

10. $n \cdot \log(n)$

- `result = n*log(n)`
- Apply a for loop for values of n from 0-100 and print all the values for result.

11.

i. Initialize a variable n.

ii. Create a function to find the factorial.

iii. `factorial(n)`

`if(n==1 || n==0)`

`return i`

`else`

`return n*factorial(n-1)`

iv. Apply a for loop for values of n from 0-19 and print all the values for result in the main function.

PROGRAM:

```
#include <stdio.h>
#include <stdlib.h>
#include <math.h>


long double factorial(int n){
    if(n==0 || n==1){
        return 1;
    }
    else{
        return n*factorial(n-1);
    }
}

int main(){
    int n,f1,f2,x=0;
    long double f4,f7,f11;
    double f3,f5,f6,f8,f9,f10;
    while(x<=11){
        printf("Enter the function no.: \n");
        scanf("%d",&x);
        if(x==1){
            printf("n\n");
            for(int i=0; i<=100; i++){
                f1=i;
                printf("%d\n",f1);
            }
        }
        else if(x==2){
            printf("n^3\n");
            for(int i=0; i<=100; i++){
                f2=i*i*i;
                printf("%d\n",f2);
            }
        }
        else if(x==3){
            printf("log(n)\n");
            for(int i=0; i<=100; i++){
                f3=log(i);
                printf("%.2lf\n",f3);
            }
        }
        else if(x==4){
            printf("n.2^n\n");
            for(int i=0; i<=100; i++){
                f4=i*(pow(2,i));
                printf("%.Lf\n",f4);
            }
        }
        else if(x==5){
```


```
printf("(3/2)n\n");
for(int i=0; i<=100; i++){
f5=1.5*i;
printf("%.2lf\n",f5);
}
}
else if(x==6){
printf("e^n\n");
for(int i=0; i<=100; i++){
f6=exp(i);
printf("%.2lf\n",f6);
}
}
else if(x==7){
printf("2^n\n");
for(int i=0; i<=100; i++){
f7=(pow(2,i));
printf("%.Lf\n",f7);
}
}
else if(x==8){
printf("log(log(n))\n");
for(int i=0; i<=100; i++){
f8=log(log(i));
printf("%.2lf\n",f8);
}
}
else if(x==9){
printf("(log n)^1/2\n");
for(int i=0; i<=100; i++){
f9=pow(log(i),0.5);
printf("%.2lf\n",f9);
}
}
else if(x==10){
printf("n*log(n)\n");
for(int i=0; i<=100; i++){
f10=i*log(i);
printf("%.2lf\n",f10);
}
}
else if(x==11){
printf("n!\n");
for(int i=0; i<20; i++){
f11=factorial(i);
printf("%.2Lf\n",f11);
}
}
else{
```

RESULT:


```
printf("Invalid function number entered. Please enter a number from 1 to 11.\n");
}
}
return 0;
}
```




The observation from the output is that the function $f1(n)$ increases linearly as n increases, with each value of $f1(n)$ being equal to n .



The data represents the cube of numbers from 0 to 100. The values increase rapidly as n increases, indicating that the cube function grows quickly. Additionally, the graph of the data would show a smooth curve that increases at an increasing rate, with a concave upward shape.




As n increases, the logarithms increase as well, but at a decreasing rate. In other words, the function $\log(n)$ grows more slowly than n itself. Additionally, the logarithm of 1 is 0, and the logarithm of 0 is negative infinity.




From this graph we observe that, for the values 0-90 the curve increases steadily and sort of linearly but from 90-100 it shoots rapidly

$$\left(\frac{3}{2}\right)^n$$




From this graph we can observe that the value of function increases linearly as the value of n increases

$$e^n$$




As n increases, the value of e^n also increases rapidly.

2^n




It increases steadily in the beginning and from 94 or 95 increases sharply.

$\log(\log(n))$




In this graph, we observe that for values 2-25, the graph grows exponentially and then increases linearly and steadily. Here we don't consider $n=0,1$ as $\log 0$ is not defined and as $\log 1$ is 0 then $\log(\log 1)$ cannot be defined too.

$(\log n)^{1/2}$ inf




This graph increases rapidly from 1 to 12-13 and then becomes steady.

$n \cdot \log(n)$



The curve in this graph increases somewhat linearly as the value of n increases

<p>CONCLUSION:</p>	<div data-bbox="483 193 1409 766">  </div> <p>The curve for factorial is linear from 0-17 and then shoots rapidly from 18-19.</p> <p>Thus we have implemented and plotted graphs for different functions and observed their growth for various cases.</p>
---------------------------	---