

GLOBAL HANTAVIRUS SURVILLANCE DASHBOARD

2K

Total Cases

1K

Hospitalized

155

Fatalities

24.26


Average Recovery

[OVERVIEW]


[ENVIRONMENTAL
ANALYSIS]

[CONCLUSIONS]


Total Cases by country




Total Cases by virus_strain



Total Cases by transmission_type



Total Cases by Month Name



Year	gender	virus_strain
	☐ Female ☐ Male	All
2025	hospitalization	country
	☐ No ☐ Yes	All
2026		

ENVIRONMENTAL FACTORS ANALYSIS

[OVERVIEW]

[ENVIRONMENTAL
ANALYSIS]

[CONCLUSIONS]

23.83

Average Temper...


65.30

Average Humidity


164.51

Average AQI


Temperature vs Humidity




Air Quality Index by Country



Rodent population by country



Recovery Trend



gender ☐ Female ☐ Male	country All ▼	Year ☐ 2025 ☐ 2026	virus_strain Dobrava ▼
---	--	---	---

GLOBAL HANTAVIRUS SURVEILLANCE DASHBOARD

CONCLUSION & KEY INSIGHTS

[OVERVIEW]

[ENVIRONMENTAL
ANALYSIS]

[CONCLUSIONS]

DISEASE BURDEN

2K+

Total reported cases analyzed globally

TRANSMISSION

50:50

Human-to-human and rodent-to-human spread

VIRUS STRAINS

5

Major strains identified across regions

RECOVERY PERIOD

24.26

Average recovery days observed

ENVIRONMENT

3

Temperature, Humidity and AQI factors

RECOMMENDATIONS

Rodent Monitoring

Environmental Data

Early Detection

High Risk Areas

Data-driven Decisions

FUTURE SCOPE

ML Prediction

Real-time Dashboard

Risk Mapping

Alert System

Advanced Analytics

The analysis indicates that environmental conditions influence rodent population patterns, which affect virus transmission risk and ultimately impact human health outcomes.

Environmental Factors
(Temperature | Humidity | AQI)



Rodent Population
(Reservoir Host)



Virus Transmission
(Rodent-to-Human / Human-to-Human)



Human Infection
(Cases & Hospitalization)



Recovery Outcome
(Recovery Period & Fatalities)