

MATB42: Assignment #8

1. A surface S is obtained by rotation the given figure in the xy -plane about the z -axis. (The arc is part of a circle of radius 1 centered at $(2,0)$.)

- (a) Paratetrize S (in pieces) and compute the surface area.

We have that the upper line when rotated, can be parametrized by a restricted cone and similarly for the bottom. The top and bottom respectively can be written as



$$\Phi(u, \theta) = ((1 - u)3 \cos \theta, (1 - u)3 \sin \theta, 3u), \quad 0 \leq u \leq 1, \quad 0 \leq \theta \leq 2\pi$$

$$\Phi(u, \theta) = ((1 + u)3 \cos \theta, (1 + u)3 \sin \theta, -3u), \quad -1 \leq u \leq 0, \quad 0 \leq \theta \leq 2\pi$$

For the circular portion to the left, when rotated around, it will be the inner half of a torus, so the equation will be

- (b) Use a computer algebra system to sketch S .

2. Let S be the cone with vertex $(2,3,3)$ and base the circle $x^2 + y^2 = 1$ in the xy -plane.

(a) Parametrize S

Starting with a base of a circle, we get $(\cos \theta, \sin \theta, 1)$ with $0 \leq \theta \leq 2\pi$. To change into a cone multiply x and y by $(1 - u)$ with $0 \leq u \leq 1$ and finally to shift the vertex, add $(2u, 3u, 2u)$ where $z = 2u$ since the base equation already has a 1, so $1 + ku \leq 3 \implies k \leq 2$.

$$\implies \Phi(u, \theta) = ((1 - u) \cos \theta + 2u, (1 - u) \sin \theta + 3u, 1 + 2u)$$

(b) Use a computer algebra system to sketch S .

(c) Write down the integral that would give the surface area of S . (You are not expected to evaluate the integral.)

$$\begin{aligned} \phi_\theta &= (-(1 - u) \sin \theta, (1 - u) \cos \theta, 0) \\ \phi_u &= (-\cos \theta + 2, -\sin \theta + 3, 2) \\ \phi_\theta \times \phi_u &= ((2(1 - u) \cos \theta), (2(1 - u) \sin \theta), \\ &\quad (-(1 - u) \sin \theta)(-\sin \theta + 3) - ((1 - u) \cos \theta)(-\cos \theta + 2)) \\ &= ((2 - 2u) \cos \theta, (2 - 2u) \sin \theta, (1 - u) \sin^2 \theta - (3 - 3u) \sin \theta + (1 - u) \cos^2 \theta - (2 - 2u) \cos \theta) \\ &= ((2 - 2u) \cos \theta, (2 - 2u) \sin \theta, (1 - u) - (3 - 3u) \sin \theta - (2 - 2u) \cos \theta) \\ \|\phi_\theta \times \phi_u\| &= \sqrt{(2 - 2u)^2 \cos^2 \theta + (2 - 2u)^2 \sin^2 \theta + ((1 - u) - (3 - 3u) \sin \theta - (2 - 2u) \cos \theta)^2} \\ &= \sqrt{(2 - 2u) + ((1 - u) - (3 - 3u) \sin \theta - (2 - 2u) \cos \theta)^2} \\ \implies \mathcal{A}(S) &= \int_0^1 \int_0^{2\pi} \sqrt{(2 - 2u) + ((1 - u) - (3 - 3u) \sin \theta - (2 - 2u) \cos \theta)^2} d\theta du \end{aligned}$$

3. Let S be the self-intersecting rectangle in \mathbb{R}^3 given by the implicit equation $x^2 - y^2z = 0$.

(a) Give a parametrization of S and use a computer algebra system to provide a sketch.

$$x^2 - y^2z = 0 \implies y^2z = x^2 \implies z = \left(\frac{x}{y}\right)^2$$

$$\Phi(x, y) = \left(x, y, \left(\frac{x}{y}\right)^2\right)$$

(b) Is your parametrization one-to-one? Explain.

Yes, if $\Phi(x_0, y_0) = \Phi(x_1, y_1)$ then $\Phi_1(x_0, y_0) = \Phi_1(x_1, y_1) \implies x_0 = x_1$, and $\Phi_2(x_0, y_0) = \Phi_2(x_1, y_1) \implies y_0 = y_1$. This means that $\Phi(x_0, y_0) = \Phi(x_1, y_1) \implies (x_0, y_0) = (x_1, y_1)$ so it is one to one.

(c) Find the equation of the tangent plane to S at $\left(\frac{1}{4}, \frac{1}{2}, \frac{1}{4}\right)$.

$$\phi_x = \left(1, 0, \frac{2x}{y^2}\right), \phi_y = \left(0, 1, \frac{-2x^2}{y^3}\right)$$

$$\phi_x \times \phi_y = \left(\frac{-2x}{y^2}, \frac{2x^2}{y^3}, 1\right)$$

$$\begin{aligned} (\phi_x \times \phi_y)\left(\frac{1}{4}, \frac{1}{2}\right) &= \left(\frac{-1/2}{1/4}, \frac{1/8}{1/8}, 1\right) \\ &= (-2, 1, 1) \end{aligned}$$

So the tangent plane is defined by the equation

$$(-2)(x - 1/4) + (y - 1/2) + (z - 1/4) = 0 \Leftrightarrow -2x + y + z = 1/4$$

4. Let S be the surface defined by $x^2 + y^2 = 1$ for $0 \leq z \leq 1$ and by $x^2 + y^2 = z^2$ for $1 \leq z \leq 2$.

(a) Use symbolic algebra software to sketch S .

(b) Evaluate $\int_S \mathbf{F} \cdot d\mathbf{S}$ where $\mathbf{F}(x, y, z) = (-y, x, z)$ and S is oriented by outward pointing normals.

S can be parametrized piecewise by $\Phi_1(u, \theta) = (\cos \theta, \sin \theta, u)$ for $0 \leq u \leq 1$ and $\Phi_2(u, \theta) = (u \cos \theta, u \sin \theta, u)$ for $1 \leq u \leq 2$

The normals are each given by

$$\phi_{1_u} = (0, 0, 1)$$

$$\phi_{1_\theta} = (-\sin \theta, \cos \theta, 0)$$

$$\phi_{2_u} = (\cos \theta, \sin \theta, 1)$$

$$\phi_{2_\theta} = (-\sin \theta, \cos \theta, 0)$$

(a) Evaluate the (vector) surface integral $\int_S \mathbf{F} \cdot d\mathbf{S}$ in each of the following cases.

i. $\mathbf{F}(x, y, z) = (1, x, z)$, S is the upper hemisphere $x^2 + y^2 + z^2 = 1$, $z \geq 0$, with \mathbf{n} pointing upward.

ii. $\mathbf{F}(x, y, z) = (2, x, z + y)$, S is that part of the plane $x + y + z = 1$ which lies in the first octant and \mathbf{n} points upward.

iii. Marsden & Tromba, page 425, #22.

5. Let S be the portion of the plane $x - 2y + z = 1$ that is cut off by the coordinate planes and the plane $x + y = 1$. Let \mathbf{V} be the velocity field $\mathbf{V}(x, y, z) = (y, z, x^2)$. Find the flow across S when \mathbf{n} points upward. Explain your answer.

6. Let S be the closed surface that consists of the hemisphere $x^2 + y^2 + z^2 = 1$, $z \geq 0$, and its base $x^2 + y^2 \leq 1$, $z = 0$. let \mathbf{E} be the electric field $\mathbf{E}(x, y, z) = (2x, 2y, 2z)$. Directly calculate the electric flux across S .