

MATB42: Assignment #8

1. A surface  $S$  is obtained by rotation the given figure in the  $xy$ -plane about the  $z$ -axis. (The arc is part of a circle of radius 1 centered at  $(2,0)$ .)

- (a) Paratetrize  $S$  (in pieces) and compute the surface area.

We have that the upper line when rotated, can be parametrized by a restricted cone and similarly for the bottom. The top and bottom respectively can be written as



$$\Phi(u, \theta) = ((1 - u)3 \cos \theta, (1 - u)3 \sin \theta, 3u), \quad 0 \leq u \leq 1, \quad 0 \leq \theta \leq 2\pi$$

$$\Phi(u, \theta) = ((1 + u)3 \cos \theta, (1 + u)3 \sin \theta, -3u), \quad -1 \leq u \leq 0, \quad 0 \leq \theta \leq 2\pi$$

For the circular portion to the left, when rotated around, it will be the inner half of a torus, so the equation will be

- (b) Use a computer algebra system to sketch  $S$ .

2. Let  $S$  be the cone with vertex  $(2,3,3)$  and base the circle  $x^2 + y^2 = 1$  in the  $xy$ -plane.

(a) Parametrize  $S$

Starting with a base of a circle, we get  $(\cos \theta, \sin \theta, 1)$  with  $0 \leq \theta \leq 2\pi$ . To change into a cone multiply  $x$  and  $y$  by  $(1 - u)$  with  $0 \leq u \leq 1$  and finally to shift the vertex, add  $(2u, 3u, 2u)$  where  $z = 2u$  since the base equation already has a 1, so  $1 + ku \leq 3 \implies k \leq 2$ .

$$\implies \Phi(u, \theta) = ((1 - u) \cos \theta + 2u, (1 - u) \sin \theta + 3u, 1 + 2u)$$

(b) Use a computer algebra system to sketch  $S$ .

(c) Write down the integral that would give the surface area of  $S$ . (You are not expected to evaluate the integral.)

$$\begin{aligned} \phi_\theta &= (-(1 - u) \sin \theta, (1 - u) \cos \theta, 0) \\ \phi_u &= (-\cos \theta + 2, -\sin \theta + 3, 2) \\ \phi_\theta \times \phi_u &= ((2(1 - u) \cos \theta), (2(1 - u) \sin \theta), \\ &\quad (-(1 - u) \sin \theta)(-\sin \theta + 3) - ((1 - u) \cos \theta)(-\cos \theta + 2)) \\ &= ((2 - 2u) \cos \theta, (2 - 2u) \sin \theta, (1 - u) \sin^2 \theta - (3 - 3u) \sin \theta + (1 - u) \cos^2 \theta - (2 - 2u) \cos \theta) \\ &= ((2 - 2u) \cos \theta, (2 - 2u) \sin \theta, (1 - u) - (3 - 3u) \sin \theta - (2 - 2u) \cos \theta) \\ \|\phi_\theta \times \phi_u\| &= \sqrt{(2 - 2u)^2 \cos^2 \theta + (2 - 2u)^2 \sin^2 \theta + ((1 - u) - (3 - 3u) \sin \theta - (2 - 2u) \cos \theta)^2} \\ &= \sqrt{(2 - 2u) + ((1 - u) - (3 - 3u) \sin \theta - (2 - 2u) \cos \theta)^2} \\ \implies \mathcal{A}(S) &= \int_0^1 \int_0^{2\pi} \sqrt{(2 - 2u) + ((1 - u) - (3 - 3u) \sin \theta - (2 - 2u) \cos \theta)^2} d\theta du \end{aligned}$$

3. Let  $S$  be the self-intersecting rectangle in  $\mathbb{R}^3$  given by the implicit equation  $x^2 - y^2z = 0$ .

(a) Give a parametrization of  $S$  and use a computer algebra system to provide a sketch.

$$x^2 - y^2z = 0 \implies y^2z = x^2 \implies z = \left(\frac{x}{y}\right)^2$$

$$\Phi(x, y) = \left(x, y, \left(\frac{x}{y}\right)^2\right)$$

(b) Is your parametrization one-to-one? Explain.

Yes, if  $\Phi(x_0, y_0) = \Phi(x_1, y_1)$  then  $\Phi_1(x_0, y_0) = \Phi_1(x_1, y_1) \implies x_0 = x_1$ , and  $\Phi_2(x_0, y_0) = \Phi_2(x_1, y_1) \implies y_0 = y_1$ . This means that  $\Phi(x_0, y_0) = \Phi(x_1, y_1) \implies (x_0, y_0) = (x_1, y_1)$  so it is one to one.

(c) Find the equation of the tangent plane to  $S$  at  $\left(\frac{1}{4}, \frac{1}{2}, \frac{1}{4}\right)$ .

$$\phi_x = \left(1, 0, \frac{2x}{y^2}\right), \phi_y = \left(0, 1, \frac{-2x^2}{y^3}\right)$$

$$\phi_x \times \phi_y = \left(\frac{-2x}{y^2}, \frac{2x^2}{y^3}, 1\right)$$

$$\|\phi_x \times \phi_y\| = \left(\frac{-2x}{y^2}, \frac{2x^2}{y^3}, 1\right)$$

4. Let  $S$  be the surface defined by  $x^2 + y^2 = 1$  for  $0 \leq z \leq 1$  and by  $x^2 + y^2 = z^2$  for  $1 \leq z \leq 2$ .

(a) Use symbolic algebra software to sketch  $S$ .

(b) Evaluate  $\int_S \mathbf{F} \cdot d\mathbf{S}$  where  $\mathbf{F}(x, y, z) = (-y, x, z)$  and  $S$  is oriented by outward pointing normals.

(a) Evaluate the (vector) surface integral  $\int_S \mathbf{F} \cdot d\mathbf{S}$  in each of the following cases.

i.  $\mathbf{F}(x, y, z) = (1, x, z)$ ,  $S$  is the upper hemisphere  $x^2 + y^2 + z^2 = 1$ ,  $z \geq 0$ , with  $\mathbf{n}$  pointing upward.

ii.  $\mathbf{F}(x, y, z) = (2, x, z + y)$ ,  $S$  is that part of the plane  $x + y + z = 1$  which lies in the first octant and  $\mathbf{n}$  points upward.

iii. Marsden & Tromba, page 425, #22.

5. Let  $S$  be the portion of the plane  $x - 2y + z = 1$  that is cut off by the coordinate planes and the plane  $x + y = 1$ . Let  $\mathbf{V}$  be the velocity field  $\mathbf{V}(x, y, z) = (y, z, x^2)$ . Find the flow across  $S$  when  $\mathbf{n}$  points upward. Explain your answer.

6. Let  $S$  be the closed surface that consists of the hemisphere  $x^2 + y^2 + z^2 = 1$ ,  $z \geq 0$ , and its base  $x^2 + y^2 \leq 1$ ,  $z = 0$ . Let  $\mathbf{E}$  be the electric field  $\mathbf{E}(x, y, z) = (2x, 2y, 2z)$ . Directly calculate the electric flux across  $S$ .