

For postmenopausal women and adult men with *ESR1*-mutated, ER+/HER2- advanced or metastatic breast cancer whose disease has progressed on endocrine therapy

WHEN IT'S ***ESR1m***,
IT MAY BE THE



Not an actual patient.

FOR ME

A guide to treatment with ORSERDU

A targeted treatment for *ESR1*-mutated,
ER+/HER2- advanced or metastatic breast cancer

INDICATION

ORSERDU (elacestrant) is a prescription medicine to treat women who have gone through menopause and adult men with estrogen receptor (ER)-positive, human epidermal growth factor receptor 2 (HER2)-negative, *ESR1*-mutated advanced breast cancer or breast cancer that has spread to other parts of the body (metastatic), **and** whose disease has progressed after endocrine therapy.

Your healthcare provider will perform a test to make sure that ORSERDU is right for you.

It is not known if ORSERDU is safe and effective in children.

ER+, estrogen receptor-positive; *ESR1*, estrogen receptor 1; *ESR1m*, estrogen receptor 1 mutation; HER2-, human epidermal growth factor receptor 2-negative.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information, including Important Facts, for ORSERDU.

IMPORTANT SAFETY INFORMATION

ORSERDU may cause serious side effects, including:

- Increased fat (lipid) levels in your blood (hypercholesterolemia and hypertriglyceridemia).

Your healthcare provider will do blood tests to check your lipid levels before and during your treatment with ORSERDU

ONCE-DAILY TABLET

 **ORSERDU**[®]
elacestrant
345 mg • 86 mg tablets

Understanding treatment with ORSERDU

If your advanced or metastatic breast cancer changes, understanding what has changed, and why, may help you and your doctor determine what comes next. If your cancer develops an *ESR1* mutation, **ORSERDU may be the 1 for you**. ORSERDU is a treatment for postmenopausal women and adult men with ER+/HER2- advanced or metastatic breast cancer that has developed an *ESR1* mutation following disease progression on hormone therapy. It's a **once-daily pill** that can be taken at home or on the go.



**In this guide,
you'll learn about:**

- How cancer can change over time
- *ESR1* mutations
- ORSERDU information
- ORSERDU results
- ORSERDU safety
- Taking ORSERDU
- Helpful resources
- Savings and support

IMPORTANT SAFETY INFORMATION (continued)

Before taking ORSERDU, tell your healthcare provider about all your medical conditions, including if you:

- Have liver problems
- Are pregnant or plan to become pregnant. ORSERDU can harm your unborn baby

Females who are able to become pregnant:

- Your healthcare provider may do a pregnancy test before you start treatment with ORSERDU


Please see additional Important Safety Information throughout and accompanying full Prescribing Information, including Important Facts, for ORSERDU.

Advanced or metastatic breast cancer **can change over time**

Mutations are one way cancer may change

A mutation is a change in your cancer makeup. These changes can affect how the cancer behaves and which treatments may work best, even though it is still the same type of cancer.

Because cancer can change over time, mutations can happen at different times. They can be found at diagnosis, or they can develop later as cancer cells change.



**Mutations give your doctors
important clues about your cancer.
They can be hints for which treatment
may be right for your cancer**



Consider asking your healthcare team:

- How do we determine if my cancer has a mutation?
- What mutations can my cancer develop over time?

IMPORTANT SAFETY INFORMATION (continued)

Before taking ORSERDU, tell your healthcare provider about all your medical conditions, including if you: (continued)


Females who are able to become pregnant: (continued)

- You should use effective (contraception) birth control during treatment with ORSERDU and for 1 week after the last dose

Please see additional Important Safety Information throughout and accompanying full Prescribing Information, including Important Facts, for ORSERDU.

The difference between **mutations** and **progression**

When mutations develop during or after treatment, it means the cancer is changing its makeup to get around treatment. But this does not always mean the cancer has progressed (spread or gotten worse).



Current treatment may still be working to control the cancer—and it may keep working for some time.

Doctors use several factors, including imaging results, symptoms, and overall clinical evaluation, to determine if cancer has progressed and is in need of a treatment change. Treatment change is typically considered when disease progression is confirmed.



Consider asking your healthcare team:

- Is my current treatment still working to control my cancer?

Talk with your doctor about making the most of every treatment option available to you

IMPORTANT SAFETY INFORMATION (continued)

Before taking ORSERDU, tell your healthcare provider about all your medical conditions, including if you: (continued)

Females who are able to become pregnant: (continued)

- Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with ORSERDU

Please see additional Important Safety Information throughout and accompanying full Prescribing Information, including Important Facts, for ORSERDU.

What is an **ESR1** mutation?

Nearly **1 out of 2**



If your cancer has this mutation, you are not alone

Nearly **1 out of 2** people with ER+/HER2- advanced or metastatic breast cancer may develop an *ESR1* mutation while on hormone therapy. The first treatment for most people is a hormone therapy, taken alone or in combination with a CDK4/6 inhibitor.

Some common hormone therapies include:

- exemestane
- letrozole
- anastrozole
- fulvestrant

Some common CDK4/6 inhibitors include:

- palbociclib
- abemaciclib
- ribociclib

ESR1 mutations may cause your treatment to stop working and cancer to progress, but they can also act as hints for what treatments may be effective in treating your cancer.



Consider asking your healthcare team:

- How do we determine my cancer's *ESR1* mutation status?

***ESR1* mutation status can help your healthcare team determine the next treatment for your cancer**

CDK4/6, cyclin-dependent kinase 4/6.

IMPORTANT SAFETY INFORMATION (continued)

Before taking ORSERDU, tell your healthcare provider about all your medical conditions, including if you: (continued)

Males with female partners who are able to become pregnant:

- You should use effective (contraception) birth control during treatment with ORSERDU and for 1 week after the last dose

Please see additional Important Safety Information throughout and accompanying full Prescribing Information, including Important Facts, for ORSERDU.

Getting tested for **ESR1** mutations



A blood test is the preferred way to detect **ESR1** mutations. A blood test:

- Can show why your disease may have progressed
- Can show why your current treatment may no longer be working
- Is fast and accurate, with results in about 1 week
- Can be done at your doctor's office or a lab



Tissue biopsies may not capture all **ESR1** mutations.

- Cancer can vary in different parts of the body. A tissue biopsy only samples a single spot, which means an **ESR1** mutation might be missed
- A liquid biopsy catches tiny pieces of cancer cells in your bloodstream. This gives your doctor a more complete view of your tumor and is the preferred way to find **ESR1** mutations



Disease progression is an important time to test for **ESR1** mutations.

- In cancer, small parts of your tumor, called cells, break apart and enter the bloodstream
- When cancer progresses, more of these cells can be found in the bloodstream
- This can make mutations, like **ESR1**, easier for doctors to detect



Consider asking your healthcare team:

- Has my cancer been tested for **ESR1** mutations?
- If my cancer has an **ESR1** mutation, what treatment may be right for me?

IMPORTANT SAFETY INFORMATION (continued)

Before taking ORSERDU, tell your healthcare provider about all your medical conditions, including if you: (continued)

- Are breastfeeding or plan to breastfeed. It is not known if ORSERDU passes into your breast milk. Do not breastfeed during treatment with ORSERDU and for 1 week after the last dose

Please see additional Important Safety Information throughout and accompanying full Prescribing Information, including Important Facts, for ORSERDU.

ORSERDU is a targeted treatment **for breast cancer that has an *ESR1* mutation**

A pioneer in treating *ESR1*-mutated advanced or metastatic breast cancer, ORSERDU:



Was the first FDA-approved treatment

for *ESR1*-mutated advanced or metastatic breast cancer.

ORSERDU was approved by the FDA in January 2023.



Is a targeted hormone therapy in 1 daily pill.

It's not chemotherapy.



Is backed by years of experience

and more than 16,000 prescriptions.



Consider asking your healthcare team:

- What could a treatment like ORSERDU mean for me?

IMPORTANT SAFETY INFORMATION (continued)

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. ORSERDU and other medicines may affect each other causing side effects. Know the medicines you take. Keep a list of them to show your healthcare provider or pharmacist when you get a new medicine.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information, including Important Facts, for ORSERDU.



How ORSERDU **was studied**

The ORSERDU clinical trial was designed to reflect real-world practice



- Studied in **228 people** to see if it would **delay disease progression in *ESR1*-mutated**, ER+/HER2- advanced or metastatic breast cancer
- ORSERDU was compared with common hormone therapies such as **exemestane, anastrozole, letrozole, and fulvestrant**. No one was given a placebo. ORSERDU was given alone and not in combination with other therapies

In the clinical trial:



- **100%** of participants had previously taken a type of treatment called a CDK4/6 inhibitor.* These included treatments such as Ibrance[®] (palbociclib), Kisqali[®] (ribociclib), and Verzenio[®] (abemaciclib)
- **72%** of participants had breast cancer that had spread to other organs, such as the liver and/or lungs[†]



Consider asking your healthcare team:

- What do I have in common with people from the clinical study?

Ibrance is a registered trademark of Pfizer Inc. Kisqali is a registered trademark of Novartis AG. Verzenio is a registered trademark owned or licensed by Eli Lilly and Company, its subsidiaries or affiliates.

*Patients whose cancer has progressed after prior hormone therapy and CDK4/6 inhibitor treatment may have a more limited response to subsequent hormone therapies.

†Patients may have had metastases in more than one location.

IMPORTANT SAFETY INFORMATION (continued)

The most common side effects of ORSERDU include (≥10% of patients):

- | | | |
|---|--|---------------------------------|
| • Muscle and joint (musculoskeletal) pain | • Decreased red blood cell counts | • Decreased appetite |
| • Nausea | • Vomiting | • Diarrhea |
| • Increased cholesterol and triglyceride levels in your blood | • Decreased salt (sodium) levels in your blood | • Headache |
| • Increased liver function tests | • Increased kidney function test | • Constipation |
| • Tiredness | | • Stomach-area (abdominal) pain |
| | | • Hot flush |
| | | • Indigestion or heartburn |

Please see additional Important Safety Information throughout and accompanying full Prescribing Information, including Important Facts, for ORSERDU.

ONCE-DAILY TABLET

ORSERDU[®]
elacestrant
345 mg • 86 mg tablets

Not actual
patients.

ORSERDU may offer **more time without disease progression**

ORSERDU helped people live 2x longer without their cancer spreading or getting worse



The median progression-free survival (mPFS) for people with *ESR1*-mutated cancer from the start of treatment was 3.8 months for ORSERDU vs 1.9 months for other commonly prescribed hormone therapies. Individual results may vary.

mPFS is a type of time measurement in a clinical trial. It measures the point in time when half of the people in the trial were living without their disease spreading or getting worse.

45%

In the clinical study, people who took ORSERDU had **45% less risk of disease progression or death** when compared with people who took fulvestrant or an aromatase inhibitor.



Consider asking your healthcare team:

- What could these results mean for me?

IMPORTANT SAFETY INFORMATION (continued)

Your healthcare provider may decrease your dose, temporarily stop, or completely stop treatment with ORSERDU, if you develop certain side effects.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information, including Important Facts, for ORSERDU.

What to expect while taking ORSERDU

Warnings and precautions



ORSERDU may cause serious side effects

- **Increased cholesterol and triglyceride levels in your blood (hypercholesterolemia and hypertriglyceridemia).** Your healthcare provider will do blood tests to check your lipid levels before and during your treatment with ORSERDU

Before taking ORSERDU, tell your healthcare provider about all your medical conditions, including if you:

- Have liver problems
- Are pregnant or plan to become pregnant. ORSERDU can harm your unborn baby

Females who are able to become pregnant:

- Your healthcare provider may do a pregnancy test before you start treatment with ORSERDU
- You should use effective (contraception) birth control during treatment with ORSERDU and for 1 week after the last dose
- Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with ORSERDU

Males with female partners who are able to become pregnant:

- You should use effective (contraception) birth control during treatment with ORSERDU and for 1 week after the last dose
- Are breastfeeding or plan to breastfeed. It is not known if ORSERDU passes into your breast milk. Do not breastfeed during treatment with ORSERDU and for 1 week after the last dose



Not actual
patients.

Warnings and precautions (continued)

Some of the most common side effects included ($\geq 10\%$):


- Muscle and joint (musculoskeletal) pain
- Nausea
- Increased cholesterol and triglyceride levels in your blood
- Increased liver function tests
- Tiredness
- Decreased red blood cell counts
- Vomiting
- Decreased salt (sodium) levels in your blood
- Increased kidney function test
- Decreased appetite
- Diarrhea
- Headache
- Constipation
- Stomach-area (abdominal) pain
- Hot flush
- Indigestion or heartburn

If you develop certain side effects, your healthcare provider may decrease your dose, temporarily stop, or completely stop treatment with ORSERDU.

These are not all of the possible side effects of ORSERDU. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

In the clinical study, most side effects were categorized as **mild to moderate**

Most people did not need to take anti-nausea medication while taking ORSERDU.



Of the people in the clinical study:

3%

needed to reduce the dose of ORSERDU due to side effects

6%

stopped taking ORSERDU due to side effects

15%

had to interrupt their treatment with ORSERDU due to side effects



Consider asking your healthcare team:

- What situations might require me to stop or change my dose of ORSERDU?

IMPORTANT SAFETY INFORMATION (continued)

ORSERDU may affect fertility in males and in females who are able to become pregnant. Talk to your healthcare provider if this is a concern for you.

ORSERDU is available as 345 mg and 86 mg tablets.

These are not all the possible side effects of ORSERDU. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088 or visit www.fda.gov/medwatch.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information, including Important Facts, for ORSERDU.

ONCE-DAILY TABLET


 **ORSERDU**[®]
elacestrant
345 mg • 86 mg tablets

Not actual
patients.

ORSERDU in the real world

ORSERDU was the first FDA-approved oral SERD for *ESR1*-mutated, ER+/HER2- advanced or metastatic breast cancer.* It has real-world data in more than 1,000 patients.

Real-world data:



- Show how **FDA-approved medicines work** outside of a clinical trial using information from real-world clinical practices
- Show how **ORSERDU worked** outside of the clinical trial[†]

Real-world data represent insights from everyday use. The information isn't intended to provide a final conclusion about a medicine's effectiveness, or be directly compared to findings from a clinical trial.



From the real world to your world, **ORSERDU has redefined treatment** for thousands of people with *ESR1*-mutated advanced or metastatic breast cancer.



Consider asking your healthcare team:

- What could real-world observations about ORSERDU mean for me and my treatment?

SERD, selective estrogen receptor degrader.

*ORSERDU was approved by the FDA in January 2023. More than 16,000 patients with *ESR1*-mutated metastatic breast cancer have been treated with ORSERDU in the US.

[†]Data come from different sources such as insurance claims for prescription medicines. Patient names are removed to protect privacy.

IMPORTANT SAFETY INFORMATION (continued)

ORSERDU may cause serious side effects, including:

- **Increased fat (lipid) levels in your blood (hypercholesterolemia and hypertriglyceridemia).** Your healthcare provider will do blood tests to check your lipid levels before and during your treatment with ORSERDU

Please see additional Important Safety Information throughout and accompanying full Prescribing Information, including Important Facts, for ORSERDU.

Once a day, every day: ORSERDU can help treatment fit into your routine



Pill not actual size.

Take ORSERDU at about the same time each day

- If a dose is missed, take your normal dose on the following day*
- Do not change your dose or stop taking ORSERDU unless your healthcare provider tells you to do so

*If you miss a dose by more than 6 hours or vomit after taking a dose of ORSERDU, do not take another dose of ORSERDU on that day. Skip the dose and take your next dose the following day at your regularly scheduled time.



Take ORSERDU with food

- This may help to reduce nausea and vomiting
- Pills should be swallowed whole. Do not chew, crush, or split pills
- **No fasting required**



ORSERDU may be taken anywhere (at home or on the go)

- ORSERDU is available as 345-mg and 86-mg tablets

One daily pill—No injections, no infusions

IMPORTANT SAFETY INFORMATION (continued)

Before taking ORSERDU, tell your healthcare provider about all your medical conditions, including if you:

- Have liver problems

Please see additional Important Safety Information throughout and accompanying full Prescribing Information, including Important Facts, for ORSERDU.

ONCE-DAILY TABLET

ORSERDU[®]
elacestrant
345 mg • 86 mg tablets

Not actual
patients.

Always take ORSERDU **exactly as prescribed** by your healthcare team



Before you start treatment

Before you start taking ORSERDU, **talk with your healthcare team** about any questions or concerns you may have. It's also important to read the **Important Facts** about ORSERDU.



After you start treatment

It's important to take ORSERDU as directed. Don't stop taking it without your doctor's direction.

Let your doctor or healthcare team know if you experience any side effects. Your healthcare provider may decrease your dose, temporarily stop, or completely stop treatment with ORSERDU if you develop certain side effects.

Tell your healthcare team about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. ORSERDU and other medicines may affect each other, causing side effects.

IMPORTANT SAFETY INFORMATION (continued)

Before taking ORSERDU, tell your healthcare provider about all your medical conditions, including if you: (continued)

- Are pregnant or plan to become pregnant. ORSERDU can harm your unborn baby

Please see additional Important Safety Information throughout and accompanying full Prescribing Information, including Important Facts, for ORSERDU.



Have questions about ORSERDU?

Here are some to start with. Ask your healthcare team:

-  Am I a candidate for ORSERDU? If so, why?
-  How can we tell if my current treatment is still working, or if it's time for a different treatment?
-  Can we schedule a blood test to check for *ESR1* mutations? When will we get results? How could this impact our next steps?
-  How well did ORSERDU work in the clinical trial? Were there results for my specific type of cancer?
-  If I start taking ORSERDU, what kind of lab work will we do? Will we do blood work and monitor lipid levels?
-  What should I know before taking ORSERDU? Could it impact my daily life? Should we discuss foods or other medications?
-  What are the most common side effects of ORSERDU?

If you are prescribed ORSERDU, take it every day as directed. If you experience any side effects, let your doctor or healthcare team know. They can help you manage certain side effects or provide other advice to help ensure you receive appropriate treatment.

IMPORTANT SAFETY INFORMATION (continued)

Before taking ORSERDU, tell your healthcare provider about all your medical conditions, including if you: (continued)

Females who are able to become pregnant:

- Your healthcare provider may do a pregnancy test before you start treatment with ORSERDU

Please see additional Important Safety Information throughout and accompanying full Prescribing Information, including Important Facts, for ORSERDU.



Not an actual
patient.

Helpful tips for your healthcare appointments



You play an important role in your care. Take part in your treatment decisions. Speak openly and honestly with your healthcare team so they understand your goals and preferences. Knowing what to expect and speaking up will help you get the most out of your treatment experience.

Additional tips:

- If possible, take a family member or friend to your appointments for support
- Come prepared for your appointments: Take notes and ask questions
- Ask your healthcare team members to repeat or write down information
- Before your next appointment, write down any changes to your overall health, treatment side effects you are experiencing, and/or changes in your daily routine

Talk to your healthcare team about any questions or concerns you may have

IMPORTANT SAFETY INFORMATION (continued)

Before taking ORSERDU, tell your healthcare provider about all your medical conditions, including if you: (continued)

Females who are able to become pregnant: (continued)

- You should use effective (contraception) birth control during treatment with ORSERDU and for 1 week after the last dose
- Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with ORSERDU

Please see additional Important Safety Information throughout and accompanying full Prescribing Information, including Important Facts, for ORSERDU.

Get support throughout treatment with Stemline ARC[®]

Stemline ARC provides access support, reimbursement assistance, and educational resources to help you focus on your health throughout treatment, including:



Understanding and navigating your insurance



Financial assistance programs and options for eligible patients



Educational resources for you and your healthcare team to help you access treatment with ORSERDU

Committed to providing helpful resources to support your treatment journey

IMPORTANT SAFETY INFORMATION (continued)

Before taking ORSERDU, tell your healthcare provider about all your medical conditions, including if you: (continued)

Males with female partners who are able to become pregnant:

- You should use effective (contraception) birth control during treatment with ORSERDU and for 1 week after the last dose

Please see additional Important Safety Information throughout and accompanying full Prescribing Information, including Important Facts, for ORSERDU.



Stemline ARC Patient Advocates provide a single point of contact

Stemline ARC Patient Advocates are here to provide ongoing support, connect you to helpful resources, and answer questions during your treatment. It's important to remember that support from Stemline ARC Patient Advocates is not intended to replace discussions between you and your healthcare team. Patient Advocates do not provide medical advice.



Here are some questions Stemline ARC Patient Advocates can help answer:

-  How can I get financial assistance during my treatment?
-  Will my insurance cover ORSERDU?
-  Where can I fill my ORSERDU prescription?
-  Where can I learn more about ER+/HER2- advanced or metastatic breast cancer?

Enroll in Stemline ARC today. Visit stemlineARC.com

IMPORTANT SAFETY INFORMATION (continued)

Before taking ORSERDU, tell your healthcare provider about all your medical conditions, including if you: (continued)

- Are breastfeeding or plan to breastfeed. It is not known if ORSERDU passes into your breast milk. Do not breastfeed during treatment with ORSERDU and for 1 week after the last dose

Please see additional Important Safety Information throughout and accompanying full Prescribing Information, including Important Facts, for ORSERDU.

ONCE-DAILY TABLET

ORSERDU[®]
elacestrant
345 mg • 86 mg tablets



ORSERDU ambassador, Sherri



ORSERDU ambassador, Bethany



The patients featured in these videos are ORSERDU ambassadors who have been compensated for sharing their story. Every patient's journey with metastatic breast cancer is unique. These patients' cancer treatment may have changed. Talk to your doctor to see if ORSERDU is right for you.

Real stories have real impact

A real-world view of the ORSERDU treatment experience



Together with their healthcare team, our ambassadors chose ORSERDU to help treat their cancer. Their stories reflect a kaleidoscope of moments of humanity, resilience, and joy.

If you or your loved one have a story, we'd love to hear from you.



Watch their stories at [ORSERDU.com/real-stories](https://www.orserdus.com/real-stories)
and learn how to share yours

Please see additional Important Safety Information throughout and accompanying full Prescribing Information, including Important Facts, for ORSERDU.

IF IT'S **ESR1m**,
COULD ORSERDU BE THE



FOR YOU?

Not an actual patient.



Talk to your doctor
about an *ESR1m*
blood test



Ask about ORSERDU,
a targeted hormone
therapy in 1 daily pill



It may offer more time
without your cancer spreading
or getting worse

mPFS: 3.8 months for ORSERDU vs 1.9 months
for other commonly prescribed hormone therapies



Learn more at ORSERDU.com

INDICATION

ORSERDU (elacestrant) is a prescription medicine to treat women who have gone through menopause and adult men with estrogen receptor (ER)-positive, human epidermal growth factor receptor 2 (HER2)-negative, ESR1-mutated advanced breast cancer or breast cancer that has spread to other parts of the body (metastatic), **and** whose disease has progressed after endocrine therapy.

Your healthcare provider will perform a test to make sure that ORSERDU is right for you. It is not known if ORSERDU is safe and effective in children.

IMPORTANT SAFETY INFORMATION (continued)

The most common side effects of ORSERDU include (≥10% of patients):

- Muscle and joint (musculoskeletal) pain
- Nausea
- Increased cholesterol and triglyceride levels in your blood
- Increased liver function tests
- Tiredness
- Decreased red blood cell counts
- Vomiting
- Decreased salt (sodium) levels in your blood
- Increased kidney function test
- Decreased appetite
- Diarrhea
- Headache
- Constipation
- Stomach-area (abdominal) pain
- Hot flush
- Indigestion or heartburn

Please see additional Important Safety Information throughout and accompanying full Prescribing Information, including Important Facts, for ORSERDU.

ONCE-DAILY TABLET

 **ORSERDU**[®]
elacestrant
345 mg • 86 mg tablets