

ONCE-DAILY TABLET

First and only treatment approved
for ER+/HER2- ESR1-mutated mBC

Getting the most
out of your treatment with

 **ORSERDU**[™]
elacestrant



Not an actual patient.

ER+, estrogen receptor-positive; ESR1, estrogen receptor 1;
HER2-, human epidermal growth factor receptor 2-negative;
mBC, metastatic breast cancer.

INDICATION

ORSERDU (elacestrant), 345 mg tablets, is a prescription medicine to treat women who have gone through menopause and adult men with estrogen receptor (ER)-positive, human epidermal growth factor receptor 2 (HER2)-negative, ESR1-mutated advanced breast cancer or breast cancer that has spread to other parts of the body (metastatic), **and** whose disease has progressed after endocrine therapy.

Your healthcare provider will perform a test to make sure that ORSERDU is right for you. It is not known if ORSERDU is safe and effective in children.

IMPORTANT SAFETY INFORMATION

ORSERDU may cause serious side effects, including:

- Increased fat (lipid) levels in your blood (hypercholesterolemia and hypertriglyceridemia). Your healthcare provider will do blood tests to check your lipid levels before and during your treatment with ORSERDU

Learn more at [ORSERDU.com](https://www.ORSERDU.com)

Please see additional Important Safety Information throughout and full Prescribing Information, including Patient Information, for ORSERDU.



Stemline[®]
A Menarini Group Company



Starting a treatment for ER+/HER2- *ESR1*-mutated mBC

When starting a new treatment for your type of metastatic breast cancer (mBC), knowing what to expect may impact your overall experience. You may also want to learn more about your type of cancer and why your doctor has prescribed ORSERDU. This guide contains information to help you understand more about ORSERDU.

Find information on:

- *ESR1* mutations
- How ORSERDU works
- ORSERDU results
- Taking ORSERDU
- What to expect with ORSERDU
- Communicating with your healthcare team
- Support resources for you

IMPORTANT SAFETY INFORMATION (continued)

Before taking ORSERDU, tell your healthcare provider about all your medical conditions, including if you:

- Have liver problems
- Are pregnant or plan to become pregnant. ORSERDU can harm your unborn baby

Females who are able to become pregnant:

- Your healthcare provider may do a pregnancy test before you start treatment with ORSERDU
- You should use effective (contraception) birth control during treatment with ORSERDU and for 1 week after the last dose

Please see additional Important Safety Information throughout and full Prescribing Information, including Patient Information, for ORSERDU.

“I’m taking an active role in treating my type of mBC when I take my ORSERDU pill every day.”



Not an
actual patient.


**“The more I learn
about metastatic
breast cancer, the
more empowered I
feel in my treatment.”**



Not an actual
patient.

About **ESR1** mutations

Tests done by your doctor showed that your type of mBC has a characteristic that makes it a target for treatment with ORSERDU. An *ESR1* mutation is a type of resistance mutation that sometimes develops after cancer is treated with certain cancer therapies. There is now a treatment specifically approved for people like you with *ESR1*-mutated mBC.



ORSERDU is a once-daily oral medication for women who have gone through menopause and adult men whose ER+/HER2- mBC has developed an *ESR1* mutation and whose tumor has shown signs of growth after treatment with endocrine therapy.

It's important to continue discussing your treatment with your healthcare team in order to take full advantage of their experience in treating your type of cancer.

IMPORTANT SAFETY INFORMATION (continued)

Females who are able to become pregnant (continued):

- Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with ORSERDU

Males with female partners who are able to become pregnant:

- You should use effective (contraception) birth control during treatment with ORSERDU and for 1 week after the last dose

Please see additional Important Safety Information throughout and full Prescribing Information, including Patient Information, for ORSERDU.

A different kind of oral treatment

ORSERDU is a type of hormone therapy called an *estrogen receptor antagonist/degrader*.

Hormone therapies work by helping to block estrogen from getting to the cells that drive breast cancer growth in patients with ER+ mBC. Without estrogen, the cancer cells may not grow, or they grow more slowly. Hormone therapies have been used to treat people with mBC for more than 2 decades.

ORSERDU is different and represents a significant development as the first oral treatment for people with ER+/HER2- *ESR1*-mutated mBC after progression on endocrine therapy.

No other treatment is specifically indicated for ER+/HER2- *ESR1*-mutated mBC.



Pill not actual size.

IMPORTANT SAFETY INFORMATION (continued)

Before taking ORSERDU, tell your healthcare provider about all your medical conditions, including if you (continued):

- Are breastfeeding or plan to breastfeed. It is not known if ORSERDU passes into your breast milk. Do not breastfeed during treatment with ORSERDU and for 1 week after the last dose

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. ORSERDU and other medicines may affect each other causing side effects. Know the medicines you take. Keep a list of them to show your healthcare provider or pharmacist when you get a new medicine.

Please see additional Important Safety Information throughout and full Prescribing Information, including Patient Information, for ORSERDU.

ORSERDU works better than certain commonly prescribed hormone therapies

Treatment with ORSERDU was evaluated in a 2-year clinical study. The study included a diverse group of people, with 228 people who had ER+/HER2- mBC with an *ESR1* mutation whose disease had progressed after prior endocrine therapy.



Median progression-free survival (mPFS) is a type of time measurement in a clinical trial. It measures the point in time when half of the people in the trial were living without their disease spreading, growing, or getting worse. The mPFS for people with *ESR1*-mutated cancer from the start of treatment was 3.8 months for ORSERDU vs 1.9 months for other commonly prescribed hormone therapies.

In the clinical study, ORSERDU was compared to traditional hormone therapies, including fulvestrant, exemestane, anastrozole, and letrozole. No one was given a placebo (sometimes called a *sugar pill*). These results were achieved with ORSERDU alone, not in combination with any other therapies.

IMPORTANT SAFETY INFORMATION (continued)

The most common side effects of ORSERDU include:

- Muscle and joint (musculoskeletal) pain
- Nausea
- Increased cholesterol and triglyceride levels in your blood
- Increased liver function tests
- Tiredness
- Decreased red blood cell counts
- Vomiting
- Decreased salt (sodium) levels in your blood
- Increased kidney function test

ORSERDU was designed to fit into your daily routine

Always take ORSERDU exactly as your healthcare team prescribes.



Take ORSERDU at about the same time each day.

- If a dose is missed, take your normal dose on the following day*
- Take ORSERDU every day until your doctor tells you to stop



Take ORSERDU with food.

- This may help to reduce nausea and vomiting
- Pills should be swallowed whole. Do not chew, crush, or split pills



ORSERDU may be taken anywhere (at home or on the go).



Before you start taking ORSERDU, be sure to read the Patient Information and Important Safety Information provided.

It's important to take ORSERDU as directed. Don't stop taking ORSERDU without your doctor's direction.

Let your doctor or healthcare team know if you experience any side effects. Tell your healthcare team about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. ORSERDU and other medicines may affect each other, causing side effects.

*If a dose is missed at the scheduled time, it can still be taken if less than 6 hours have passed. If more than 6 hours have passed after the scheduled dosing time, then the missed dose should be skipped, and regular dosing should resume the next day.

IMPORTANT SAFETY INFORMATION (continued)

The most common side effects of ORSERDU include (continued):

- Decreased appetite
- Diarrhea
- Headache
- Constipation
- Stomach-area (abdominal) pain
- Hot flush
- Indigestion or heartburn

Please see additional Important Safety Information throughout and full Prescribing Information, including Patient Information, for ORSERDU.

What to expect while taking ORSERDU

In the clinical study, most side effects were classified as mild and did not frequently require people to stop or reduce their treatment.

The most common side effects included ($\geq 10\%$):

- Muscle and joint pain
- Headache
- Vomiting
- Tiredness
- Hot flush
- Diarrhea
- Decreased appetite
- Nausea
- Constipation

Additional side effects ($\geq 10\%$) included increased cholesterol and triglyceride levels in your blood, increased liver function tests, decreased red blood cell counts, decreased salt (sodium) levels in your blood, increased kidney function test, stomach-area (abdominal) pain, and indigestion or heartburn.



Remember to let your doctor or healthcare team know if you experience any side effects during treatment.

It is important to continue taking your medication as prescribed until you speak with your healthcare team. Ask them about tools and strategies that may help manage certain side effects.

Not an actual patient.

IMPORTANT SAFETY INFORMATION (continued)

Your healthcare provider may decrease your dose, temporarily stop, or completely stop treatment with ORSERDU, if you develop certain side effects.

ORSERDU may affect fertility in males and in females who are able to become pregnant. Talk to your healthcare provider if this is a concern for you.

Warnings and precautions

ORSERDU may cause serious side effects, including:

- Increased fat (lipid) levels in your blood (hypercholesterolemia and hypertriglyceridemia). Your healthcare provider will do blood tests to check your lipid levels before and during your treatment with ORSERDU

Before taking ORSERDU, tell your healthcare provider about all of your medical conditions, including if you:

- Have liver problems
- Are pregnant or plan to become pregnant. ORSERDU can harm your unborn baby

Females who are able to become pregnant:

- Your healthcare provider may do a pregnancy test before you start treatment with ORSERDU
- You should use effective (contraception) birth control during treatment with ORSERDU and for 1 week after the last dose
- Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with ORSERDU

Males with female partners who are able to become pregnant:

- You should use effective (contraception) birth control during treatment with ORSERDU and for 1 week after the last dose

IMPORTANT SAFETY INFORMATION (continued)

These are not all the possible side effects of ORSERDU. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088 or visit www.fda.gov/medwatch.



Questions you may have



It's important to communicate with your healthcare team regularly. Remember to speak up and ask any questions as they come up. Here are some to get you started:

- ✓ How long can I expect to take ORSERDU?
- ✓ What are the possible side effects?
- ✓ Will I need to come into the clinic for any type of monitoring?
- ✓ Can I do anything to help manage side effects?
- ✓ When should I contact the doctor or nurse?
- ✓ What should I know or do before my next office visit?

Additional tips:

- If possible, take a family member or friend to your appointments for support
- Ask your healthcare team members to repeat or write down information
- Before your next appointment, write down any changes to your overall health, treatment side effects, and/or changes in your daily routine

Notes

You play a crucial role in your healthcare, and it's important to share your experience with your healthcare team. Write down questions as they come up so you can take them to your follow-up appointments.

Please see Important Safety Information throughout and full Prescribing Information, including Patient Information, for ORSERDU.

Stemline ARC® offers you comprehensive support

Stemline ARC provides access support, reimbursement assistance, and educational resources to help you focus on what matters most—your health.

Stemline ARC may help with:



Understanding and navigating your insurance



Financial assistance programs and options for eligible patients



Educational resources for you and your healthcare team to help you access treatment with ORSERDU

“It’s the ORSERDU team’s goal to create tools and resources that may help you have a positive treatment experience while living with *ESR1*-mutated mBC.”



Access these resources by scanning this QR code



Some helpful resources




CANCERcare®
Over 75 Years of Help and Hope
cancercares.org




**LIVING BEYOND
BREAST CANCER®**
lbbc.org




**Tigerlily
Foundation**
Beauty. Strength. Transformation.
tigerlilyfoundation.org

The above organizations provide support for cancer patients. This is not an exhaustive list. These organizations are independent from Stemline Therapeutics, Inc. ("Stemline"). Stemline has no financial interest in these organizations but may occasionally provide some financial support. All trademarks are property of their respective owners.

ONCE-DAILY TABLET



Learn more about your treatment at ORSERDU.com

It's important that your voice be heard throughout your treatment. Your healthcare team wants to hear about your experience. Remember to share how you're feeling, and take the time to ask questions.

INDICATION

ORSERDU (elacestrant), 345 mg tablets, is a prescription medicine to treat women who have gone through menopause and adult men with estrogen receptor (ER)-positive, human epidermal growth factor receptor 2 (HER2)-negative, ESR1-mutated advanced breast cancer or breast cancer that has spread to other parts of the body (metastatic), **and** whose disease has progressed after endocrine therapy.

Your healthcare provider will perform a test to make sure that ORSERDU is right for you. It is not known if ORSERDU is safe and effective in children.

IMPORTANT SAFETY INFORMATION (continued)

The most common side effects of ORSERDU include:

- Muscle and joint (musculoskeletal) pain
- Nausea
- Increased cholesterol and triglyceride levels in your blood
- Increased liver function tests
- Tiredness
- Decreased red blood cell counts
- Vomiting
- Decreased salt (sodium) levels in your blood
- Increased kidney function test
- Decreased appetite
- Diarrhea
- Headache
- Constipation
- Stomach-area (abdominal) pain
- Hot flush
- Indigestion or heartburn

Please see additional Important Safety Information throughout and full Prescribing Information, including Patient Information, for ORSERDU.