

# MATH 307

## CHAPTER 1

### SECTION 1.6: FURTHER PROPERTIES OF DETERMINANTS

CONTENTS
----------

---

<b>Determinants and Algebraic Operations</b>	<b>2</b>
Invertibility . . . . .	2
Matrix Multiplication . . . . .	3
Transpose . . . . .	3
<b>Ajoint of a Matrix</b>	<b>4</b>
Matrix of Cofactors . . . . .	4
Definition of the Adjoint . . . . .	5
Another Way to Find the Inverse . . . . .	5
<b>Cramer's Rule</b>	<b>6</b>

## Invertibility

A square matrix  $A$  is invertible if and only if  $\det(A) \neq 0$ .

Remark:

- If  $A$  is an invertible matrix, then  $\det(A^{-1}) = 1/\det(A)$ .
- Determinant can help to determine if a system of linear equations has a solution or not.

**EXAMPLE 1.** Which of the following matrices are invertible:

$$A = \begin{bmatrix} 1 & 2 \\ 2 & 3 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & 2 & 3 & 4 \\ -4 & -3 & -2 & 1 \\ 0 & 0 & 0 & 0 \\ 1 & 4 & 5 & 9 \end{bmatrix} \quad \text{and} \quad C = \begin{bmatrix} 1 & 2 & -1 \\ 0 & 3 & -1 \\ 0 & 0 & 4 \end{bmatrix}.$$

## Matrix Multiplication

If  $A$  and  $B$  are two  $n \times n$  matrices, then

$$\det(AB) = \det(A) \det(B).$$

**EXAMPLE 2.** Knowing that  $\det(A) = 2$  and  $\det(AB) = 32$ , find the determinant of the matrix  $B$ .

## Transpose

If  $A$  is a square matrix, then  $\det(A^\top) = \det(A)$ .

**EXAMPLE 3.** If  $A$  is the matrix

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 5 & -2 \\ 0 & 0 & 1 \end{bmatrix},$$

then find the determinant of  $AA^\top$ .

## Matrix of Cofactors

The cofactor matrix is the matrix  $C$  of all the cofactors of a given matrix  $A$ . If  $A$  has dimensions  $n \times n$ , then

$$C = \begin{bmatrix} C_{11} & C_{12} & \cdots & C_{1n} \\ C_{21} & C_{22} & \cdots & C_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ C_{n1} & C_{n2} & \cdots & C_{nn} \end{bmatrix}$$

where each  $C_{ij}$  is the cofactor obtained from  $A$ .

**EXAMPLE 4.** Find the matrix of cofactors of the following matrix

$$A = \begin{bmatrix} -2 & 3 & 0 \\ 4 & 10 & 2 \\ -5 & 7 & 0 \end{bmatrix}.$$

## Definition of the Adjoint

The adjoint of a matrix  $A$  of dimensions  $n \times n$  is the transpose of the cofactor matrix.

Explicitly, we denote the adjoint of  $A$  by  $\text{adj}(A)$  and its expression is

$$\text{adj}(A) = C^T = \begin{bmatrix} C_{11} & C_{21} & \cdots & C_{n1} \\ C_{12} & C_{22} & \cdots & C_{n2} \\ \vdots & \vdots & \ddots & \vdots \\ C_{1n} & C_{2n} & \cdots & C_{nn} \end{bmatrix}.$$

**EXAMPLE 5.** Find the adjoint of the matrix in Example 4.

## Another Way to Find the Inverse

If  $A$  is an invertible matrix, then

$$A^{-1} = \frac{1}{\det(A)} \text{adj}(A).$$

**EXAMPLE 6.** Find the inverse of the matrix in Example 4.

## CRAMER'S RULE

---

Suppose that  $AX = B$  is a system of  $n$  linear equations in  $n$  unknowns such that  $\det(A) \neq 0$ . Let

- $A_1$  be the matrix obtained from  $A$  by replacing the first column of  $A$  by  $B$ ;
- $A_2$  be the matrix obtained from  $A$  by replacing the second column of  $A$  by  $B$ .
- etc.

Then the solutions to the system are

$$x_1 = \frac{\det(A_1)}{\det(A)}, \quad x_2 = \frac{\det(A_2)}{\det(A)}, \quad \dots, \quad x_n = \frac{\det(A_n)}{\det(A)}.$$

**EXAMPLE 7.** Use Cramer's rule to solve the system

$$\begin{aligned} -2x + 3y &= 2 \\ 4x + 10y + 2z &= 3 \\ -5x + 7y &= 1. \end{aligned}$$