

UNIVERSITY OF HAWAI'I



Last name: Jones

First name: Lillie

Question:	1	2	3	4	Total
Points:	15	15	15	5	50
Score:	11	10	11	0	32

Instructions:

- Write your complete name on your copy.
- Answer all 4 questions below.
- Write your answers directly on the questionnaire.
- Show ALL your work to have full credit.
- Draw a square around your final answer.
- Return your copy when you're done or at the end of the 50min period.
- No electronic devices allowed during the exam.
- Scientific calculator allowed only (no graphical calculators).
- **Turn off your cellphone(s) during the exam.**
- Lecture notes and the textbook are not allowed during the exam.

Your Signature: Lillie Jones

$$\ker T = 0 \quad \dim \mathbb{R}^3 = 3$$

QUESTION 1

(15 pts)

Let $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be defined by $T(x, y, z) = (2y, z, z)$.

- (5 Pts) Show that T is a linear transformation.
- (5 Pts) Find the kernel of T and its dimension.
- (5 Pts) Using the Dimension Theorem, deduce the rank of T .

$$\ker + \text{Rank} = \dim \mathbb{R}^3$$

$$1) \textcircled{11} T(u+v) = T(u) + T(v)$$

$$T\left(\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} + \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} + \begin{bmatrix} z_1 \\ z_2 \\ z_3 \end{bmatrix}\right) = T$$

11/15

$$\textcircled{12} T(av) = aT(v)$$

$$b) T(x, y, z) = (2y, z, z) = 0$$

$$a(2y) + b(z) + c(z) = 0$$

$$c) \dim \ker T = 3$$

$$\dim \mathbb{R}^3 = 3$$

$$x(0) = 0$$

$$2a = 0$$

$$y(2a) = 0 \Rightarrow b + c = 0$$

$$b + c = 0$$

$$a = 0$$

$$b = -c$$

$$c = t$$

$$\begin{bmatrix} 0 \\ -t \\ t \end{bmatrix} \rightarrow t \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix}$$

$$z(b+c) = 0$$

$$\dim = 1$$

QUESTION 2

$\vec{b_1}$ $\vec{b_2}$ $\vec{b_3}$

(15 pts)

Let $V = \mathbb{R}^3$. Let B be the standard basis $\{(1, 0, 0), (0, 1, 0), (0, 0, 1)\}$ and let the set $D = \{(1, 2, 0), (0, -1, 2), (0, 2, 0)\}$ be another basis of \mathbb{R}^3 .

(a) (10 Pts) Knowing that

- $(1, 0, 0) = a(1, 2, 0) + b(0, -1, 2) + c(0, 2, 0)$, where $a = 1, b = 0, c = -1$;
- $(0, 1, 0) = a(1, 2, 0) + b(0, -1, 2) + c(0, 2, 0)$, where $a = 0, b = 0, c = 1/2$;
- $(0, 0, 1) = a(1, 2, 0) + b(0, -1, 2) + c(0, 2, 0)$, where $a = 0, b = 1/2, c = 1/4$;

Find the change matrix $P_{D \leftarrow B}$. Justify carefully your answer.

(b) (5 Pts) Let $T(x, y, z) = (2y, z, z)$. Find the matrix representation of T on the basis B , that is $M_B(T)$.

$$c_D(\vec{b_1}) = \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}$$

$$\begin{aligned} c_D(\vec{b_1}) &= 1(1, 2, 0) + 0(0, -1, 2) + (-1)(0, 2, 0) \\ &= (1, 2, 0) + (0, -2, 0) \\ &= (1, 0, 0) \end{aligned}$$

$$c_D(\vec{b_2}) = \begin{bmatrix} 0 \\ 0 \\ 1/2 \end{bmatrix}$$

$$\begin{aligned} c_D(\vec{b_2}) &= 0(1, 2, 0) + 0(0, -1, 2) + 1/2(0, 2, 0) \\ &= (0, 1, 0) \end{aligned}$$

$$c_D(\vec{b_3}) = \begin{bmatrix} 0 \\ 1/2 \\ 1/4 \end{bmatrix}$$

$$\begin{aligned} c_D(\vec{b_3}) &= 0(1, 2, 0) + 1/2(0, -1, 2) + 1/4(0, 2, 0) \\ &= (0, 0, 1) \end{aligned}$$

$$P_{D \leftarrow B} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1/2 \\ -1 & 1/2 & 1/4 \end{bmatrix}$$

$$c_D(\vec{b_1}(T)) = \begin{bmatrix} 0 \\ -1 \\ -1 \end{bmatrix}$$

$$c_D(\vec{b_2}(T)) = \begin{bmatrix} 0 \\ 1/2 \\ 1/2 \end{bmatrix}$$

$$c_D(\vec{b_3}(T)) = \begin{bmatrix} 1 \\ 1/4 \\ 1/4 \end{bmatrix}$$

$$M_B(T) = \begin{bmatrix} 0 & 0 & 1 \\ -1 & 1/2 & 1/4 \\ -1 & 1/2 & 1/4 \end{bmatrix}$$

9/5

Find $T(1, 0, 0)$
 $T(0, 1, 0)$
 $T(0, 0, 1)$

and write those
in the basis B .

10/15

$$\frac{6}{10} = \frac{3}{5} \quad \frac{3}{5} \cdot \frac{3}{1} = \frac{9}{5}$$

QUESTION 3

(15 pts)

Let $U = \text{span}\{(-1, 0, 3), (0, -3, 2)\}$.

- (a) (3 Pts) Are $(-1, 0, 3)$ and $(0, -3, 2)$ orthogonal?
- (b) (5 Pts) What is the dimension of U . Justify carefully your answer.
- (c) (5 Pts) Using the Gram-Schmidt process, transform the set of vectors $\{(-1, 0, 3), (0, -3, 2)\}$ in a set of orthogonal vectors F .
- (d) (2 Pts) Illustrate visually the Gram-Schmidt orthogonalization process for two vectors in \mathbb{R}^3 .

a) $(-1)(0) + (0)(-3) + (3)(2) = 6 \neq 0 \checkmark \rightarrow$ not orthogonal 3/3

b) $\dim U = 2 \checkmark$ More justification 3/5

c)
$$v_2 = (0, -3, 2) - \frac{(-1, 0, 3)(0, -3, 2)}{(-1)^2 + (0)^2 + (3)^2} \cdot (-1, 0, 3)$$



$$= (0, -3, 2) - \left(\frac{3}{5}\right) \cdot (-1, 0, 3)$$

$$= \left(0 + \frac{3}{5}, -3 - 0, 2 - \frac{9}{5}\right)$$

$$= \left(\frac{3}{5}, -3, \frac{1}{5}\right)$$
 $\frac{10}{5} = \frac{10}{5}$ 5/5

$\vec{F} = \left\{ (-1, 0, 3), \left(\frac{3}{5}, -3, \frac{1}{5}\right) \right\}$ ✓

d)



0/2

11/15

4

QUESTION 4

(5 pts)

Assume that $T : V \rightarrow \mathbb{R}^2$ is a linear transformation where V is a vector space and \mathbb{R} is the vector space of real numbers. Let $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ be a basis of V .

- (a) (1 Pt) If $T(\mathbf{v}_1) = (2, 2)$, $T(\mathbf{v}_2) = (1, 2)$ and $T(\mathbf{v}_3) = (2, 1)$, show that $-3\mathbf{v}_1 + 2\mathbf{v}_2 + 2\mathbf{v}_3 \in \ker T$. Explain carefully your answer.
- (b) (4 Pts) If $T(\mathbf{v}_1)$, $T(\mathbf{v}_2)$ and $T(\mathbf{v}_3)$ are defined as in part (a), find the nullity of T . Explain carefully your answer.

$$\begin{aligned} 9) \quad T(\mathbf{v}_1 + \mathbf{v}_2 + \mathbf{v}_3) &= T(\mathbf{v}_1) + T(\mathbf{v}_2) + T(\mathbf{v}_3) \\ T(\mathbf{v}_1 + \mathbf{v}_2 + \mathbf{v}_3) &= (2, 2) + (1, 2) + (2, 1) \end{aligned}$$

almost!
on the right track!

0/1

Assume linear independence
 $-3\mathbf{v}_1 + 2\mathbf{v}_2 + 2\mathbf{v}_3 = \mathbf{0}$

$$-3(\mathbf{v}_1) =$$

0/5