

MATH 311

LAST CHAPTER

SECTION 8.1: ORTHOGONAL COMPLEMENTS AND PROJECTIONS

CONTENTS

Gram-Schmidt Orthogonalization	2
The Gram-Schmidt Orthogonalization Algorithm	3

CREATED BY: PIERRE-OLIVIER PARISÉ
SPRING 2024

GRAM-SCHMIDT ORTHOGONALIZATION

EXAMPLE 1. Let $V = \mathbb{R}^2$ and $B = \{(1, -1), (2, 1)\}$. Notice that B is not an orthogonal basis. Using the vectors from B , construct an orthogonal basis F .

SOLUTION. Geometric intuition: <https://www.desmos.com/geometry/e9mrgozymb>.

The Gram-Schmidt Orthogonalization Algorithm

Let $B = \{\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_m\}$ be a basis of a subspace U of $V = \mathbb{R}^n$.

To transform B into an orthogonal basis $F = \{\mathbf{f}_1, \mathbf{f}_2, \dots, \mathbf{f}_m\}$, we define

- $\mathbf{f}_1 = \mathbf{b}_1$.
- $\mathbf{f}_2 = \mathbf{b}_2 - \frac{\mathbf{b}_2 \cdot \mathbf{f}_1}{\|\mathbf{f}_1\|^2} \mathbf{f}_1$.
- $\mathbf{f}_3 = \mathbf{b}_3 - \frac{\mathbf{b}_3 \cdot \mathbf{f}_1}{\|\mathbf{f}_1\|^2} \mathbf{f}_1 - \frac{\mathbf{b}_3 \cdot \mathbf{f}_2}{\|\mathbf{f}_2\|^2} \mathbf{f}_2$.
- \dots
- $\mathbf{f}_k = \mathbf{b}_k - \frac{\mathbf{b}_k \cdot \mathbf{f}_1}{\|\mathbf{f}_1\|^2} \mathbf{f}_1 - \frac{\mathbf{b}_k \cdot \mathbf{f}_2}{\|\mathbf{f}_2\|^2} \mathbf{f}_2 - \dots - \frac{\mathbf{b}_k \cdot \mathbf{f}_{k-1}}{\|\mathbf{f}_{k-1}\|^2} \mathbf{f}_{k-1}$.
- \dots
- $\mathbf{f}_m = \mathbf{b}_m - \frac{\mathbf{b}_m \cdot \mathbf{f}_1}{\|\mathbf{f}_1\|^2} \mathbf{f}_1 - \frac{\mathbf{b}_m \cdot \mathbf{f}_2}{\|\mathbf{f}_2\|^2} \mathbf{f}_2 - \dots - \frac{\mathbf{b}_m \cdot \mathbf{f}_{m-1}}{\|\mathbf{f}_{m-1}\|^2} \mathbf{f}_{m-1}$.